CM-P00057799

Ref. TH. 885

Archive

RHO MESON SHAPEJán Pišút ^{*)} and Matts Roos

CERN - Geneva

A B S T R A C T

A detailed description is developed for the invariant dipion mass distribution in the reaction $\pi p \rightarrow \pi\pi N$. Based on OPEM and modified phenomenologically this "generalized Breit-Wigner" formula accounts for peripheral production, t dependence of the ρ parameters, skewness of the ρ peak, energy dependence of the ρ width, energy-dependent departure from OPEM, background, and background interference. Contributions from other $\pi\pi$ partial waves are included and the possibility of ρ being a CDD pole considered. The formula is used on a data compilation of large statistics and conclusions concerning the presence of the various terms and the ρ parameters are reached. Finally, we compare the width of the pion produced ρ meson with the width of the e^+e^- produced ρ meson. In an Appendix we discuss the A_2 meson shape.

*) On leave from the University of Bratislava,
Katedra teoretickej fyziky, Bratislava,
Czechoslovakia.

1. INTRODUCTION

In a previous publication ¹⁾ (to be called "paper I" henceforth) one of us analysed a compilation of invariant dipion mass histograms in order to determine the mass and width of the ϱ meson. The distribution of ϱ meson events was analysed by a "conventional" Breit-Wigner formula, the width having a p wave energy dependence. The background was approximated by an empirical three-parameter formula. The experimental samples were enriched in ϱ mesons by selecting events of low momentum transfer $-t$ to the final baryon

$$-t_{\min} < -t < -T.$$

The analysis in paper I gave different values for the ϱ^+ and ϱ^- mass in samples of small $|T|$, and the results were also significantly T dependent. The ϱ^0 mass appeared to be T dependent when the ϱ^0 meson was produced together with Δ^{++} , but constant when produced with a neutron. The width of all samples appeared to be large, of the order of 140-170 MeV. When broadening due to the T dependence of the mass was taken into account, all samples yielded a consistent natural width, averaging 129 MeV. Paper I also concluded that ω mesons were present in the ϱ^0 samples with a relative amplitude different from zero by about three standard deviations.

The above results, in particular the ϱ mass T dependence and the $\varrho^- - \varrho^+$ mass difference, indicated an insufficient understanding of the production mechanism and the structure of the background. In the present paper we therefore undertake a renewed analysis of ϱ mesons of all charge states, produced in the reactions

$$\pi^\pm p \rightarrow \pi \pi N. \quad (1)$$

The main change in this renewed analysis, apart from some changes in the composition of our data compilation, is in the theoretical expressions for the shape of resonance peaks.

The reason why we analyse only invariant mass distributions and not angular distributions is because our data compilation in paper I happened not to include angular distributions. An ultimate understanding of the Q meson can, however, only be reached through analysing a simultaneous distribution in angles and invariant mass. The present paper therefore represents the first necessary step towards such a final analysis.

Usually the expression for the shape of a resonance peak is called a Breit-Wigner formula. There are, however, Breit-Wigner formulae and "Breit-Wigner formulae", as will be seen below. Let us write the invariant mass distribution ²⁾ for the production and subsequent decay of a resonance of energy ω_r

$$\frac{d\sigma}{d\omega} = d\sigma_s(\omega) \frac{2\pi^{-1} \cdot \omega \cdot \omega_r \cdot \Gamma(\omega)}{(\omega_r^2 - \omega^2)^2 + \omega_r^2 \Gamma^2(\omega)} \quad , \quad (2)$$

where $d\sigma_s(\omega)$ is the production cross-section of the resonance (treated as a stable particle of mass ω). We are now free to

- i) choose different parametrizations for $\Gamma(\omega)$
- ii) make different dynamical assumptions about $d\sigma_s(\omega)$.

The large variety of "conventional" Breit-Wigner formulae in use arises from different choices of $\Gamma(\omega)$ and different assumptions about $d\sigma_s(\omega)$. These assumptions are, it seems, among the most tacit ones in all of strong interaction physics. This is the primary reason why everybody reports such different Q masses and widths.

As we shall show later the success of some simplified "Breit-Wigner formulae" is entirely accidental and due to the neglect of a fortuitous combination of energy-dependent factors.

For narrow resonances the form of Eq. (2) is unimportant, but for the broad Q meson the assumptions under ii) become crucial. Both i) and ii) are important to consider carefully when the statistics become as large as in our samples. The freedom in choice i) leads to systematic uncertainties in the parameter values which we shall call "parametrization errors", and the freedom in ii) correspondingly to "dynamical errors". In general, the dynamical errors turn out to be much larger than the parametrization errors. The latter are of the same order as the statistical errors.

In Section 2 we derive a detailed, and as we hope, reasonable formula for the invariant mass distribution of the $\pi\pi$ in reaction (1). More attention is paid to the background than before ¹⁾ and, because of our large statistics, we can afford to introduce several small terms. The final formula is therefore quite general; it might be described as part OPEM, part phenomenological, part conventional.

The modifications to the ordinary two-parameter resonance shape are several. Explicit t dependence (peripheral production) is built in, in a phenomenological manner; thus one may have some hope of resolving the previously observed ¹⁾ T dependence of the Q mass. In Section 3 we discuss various choices of kind i). Part of that discussion is separated off into an Appendix section in order to facilitate reading.

In Section 4 we discuss phenomenologically some dynamical assumptions of kind ii) and the form of possible small terms: $\pi\pi$ scattering in other waves than $\ell = 1$, and background interference.

Section 5 then contains the detailed results of comparing the general shape formula with the compiled $\pi\pi$ invariant mass distributions: the parameters of the ρ mesons, their T dependence, and numerous conclusions concerning various modifications of the Breit-Wigner formula, the background, and the presence of the small terms.

In Section 6 we compare and discuss the shape of ρ^0 mesons produced in reaction (1) and in e^+e^- colliding beam experiments³⁾. Section 7 briefly summarizes the results of the previous Sections.

In Section A.3 of the Appendix we discuss the shape of the A_2 meson and derive an expression for the $\pi\rho$ scattering cross-section under the assumption of two resonances in the same partial wave.

2. DERIVATION OF THE RESONANCE SHAPE FORMULA

In this section we shall derive a general formula for the invariant mass distribution of the two outgoing pions in the reaction (1) dominated by the production and decay of ρ mesons

$$\pi p \rightarrow \rho N \rightarrow \pi \pi N \quad (3)$$

It is well known that events of reaction (1) populate the Chew-Low plot predominantly in the region of small momentum transfer $-t$ (to the proton). This is particularly true for events in the ρ meson mass band. It is therefore likely that peripheral meson production mechanisms, such as the OPE diagram, play an important rôle.

On the other hand, it is also known that the simple OPEM is in disagreement with experiment, and that many drastic modifications are needed to find a good, working model. One such model is OPEM with absorption ⁴⁾ which has had some surprising successes. Its agreement with experiment is, however, not general ⁵⁾ and it is certainly very cumbersome to use.

Our approach towards a working model is essentially heuristic. We take the basic framework of the formalism from the simple OPEM, in order to respect the dominantly peripheral features of the production process. We then modify this framework, whenever required by experiment, by admitting various empirical functions of energy or momentum transfer. In this way we believe one may have a manageable description of the process, and at the same time learn something about the requirements on a working model.

The distribution (2), although flexible, must be supplemented by a background term. Thus our general shape formula will be of the form

$$\frac{d\sigma}{d\omega} = |A_{\pi} + A_B|^2 \quad (4)$$

Here A_{π} corresponds to reaction (3) and A_B corresponds to the remainder of reaction (1). The division into A_{π} and A_B is, however, not unambiguous. Certainly A_{π} should contain the pion pole. But if A_{π} consists only of the pion pole, the experimental distribution would require A_B to have a very complicated form. We want to keep A_B simple and interpretable as a smooth background term. In consequence the pion pole term must suffer some modifications.

To be less tacit than has been the custom, we shall carry out below the derivation of the r.h.s. of Eq. (4) in somewhat pedantic detail.

Let us define the S matrix element for reaction (1) as

$$S_{fi} = \delta_{fi} + i(2\pi)^4 \delta^4(P_f - P_i) (2^5 \epsilon' E' \epsilon_1 \epsilon_2 E)^{-\frac{1}{2}} F_{fi} \quad (5)$$

where ϵ' , ϵ_1 , ϵ_2 denote the energies of the initial and final pions, respectively, and similarly E' and E for the nucleons. F_{fi} is the invariant amplitude for the process. Spin and isospin variables are suppressed. The total cross-section is then given by

$$\sigma = \frac{(2\pi)^4}{4 p_{iL} M_N} \int |F_{fi}|^2 \delta^4(P_f - P_i) \frac{d^3 q_1}{2\epsilon_1 (2\pi)^3} \frac{d^3 q_2}{2\epsilon_2 (2\pi)^3} \frac{d^3 P}{2E (2\pi)^3}, \quad (6)$$

where p_{iL} is the laboratory momentum of the incident pion, and the rest of the notation should be self-explanatory.

Let us introduce the square of the momentum transfer between the initial and final nucleon

$$t = (p_{iN} - p_{fN})^2,$$

the squared invariant mass of the two final pions

$$\omega^2 = (q_1 + q_2)^2,$$

and the constant

$$C_1 = 1/\pi M_N^2 (8\pi)^4.$$

We can then write down the differential cross-section

$$\frac{d\sigma}{d\omega^2 dt} = C_1 \frac{q}{p_{iN}^2 \omega} \overline{|F(t, \omega, s)|^2} \quad (7)$$

where

$$q = \frac{1}{2} (\omega^2 - 4m_\pi^2)^{\frac{1}{2}}$$

and $\overline{|F|^2}$ is the square of the invariant amplitude in Eq. (6) integrated over the angles in the final state. The quantity F is, in fact, proportional to the Feynman amplitude for reaction (1). Let us decompose F into its OPE part and its non-OPE background part,

$$F = F_{\text{OPE}} + F_{\text{B}} \quad (8)$$

8.

F_{OPE} represents the diagram in Fig. 1, in which the upper

Fig. 1

bubble corresponds to $\pi\pi$ scattering with the exchanged pion on the mass shell, and the lower bubble is the πNN vertex,

$$F_{\text{OPE}}(\omega, t, s, \cos \theta_{\pi\pi}) = A_{\pi\pi}(\omega, t, \cos \theta_{\pi\pi}) K'(t), \quad (9)$$

where $\theta_{\pi\pi}$ is the azimuthal angle in the c.m.s. of the final pions. $K'(t)$ depends only on t , and contains the vertex corrections in the lower bubble and in the propagator.

Let us assume, for simplicity, that $A_{\pi\pi}$ is factorizable into a t dependent factor and a $(\omega, \cos \theta)$ dependent factor. We can then absorb the t dependent part into $K'(t)$, writing

$$F_{\text{OPE}}(\omega, t, s, \cos \theta_{\pi\pi}) = K(t) A_{\pi\pi}(\omega, \cos \theta_{\pi\pi}), \quad (10)$$

where $K(t)$ is now completely phenomenological. Note that such a factorization is not true in general, not even if the Selleri correction ⁶⁾ and barrier penetration factors are included in the peripheral model. (These corrections do not influence significantly the invariant mass distribution. For a fuller discussion, see Appendix A.1.) We shall shortly return to a discussion of the form of the function $K(t)$.

With Eqs. (8) and (10), Eq. (7) can be written

$$\frac{d\sigma}{d\omega^2 dt} = \frac{C q}{P_{il}^2 \omega} \left| d(\cos \theta_{\pi\pi}) \left[K(t) A_{\pi\pi}(\omega, \cos \theta_{\pi\pi}) + F_B(\omega, t, \cos \theta_{\pi\pi}) \right] \right|^2, \quad (11)$$

where F_B is an as yet unspecified background term, assumed only to be independent of the polar angle in the c.m.s. of the final pions. The invariant Feynman amplitude for $\pi\pi$ scattering in the isospin T channel is now defined as

$$A_{\pi\pi}^T(\omega, \cos \theta_{\pi\pi}) = \frac{8\pi\omega}{q} \sum_{\ell} (2\ell+1) P_{\ell}(\cos \theta_{\pi\pi}) \frac{\exp(2i\delta_{\ell}^T) - 1}{2i}. \quad (12)$$

The amplitudes for $\pi^{\pm}\pi^0$ and $\pi^+\pi^-$ scattering may be obtained using the well-known formulae

$$\begin{aligned} A^{\pm 0} &= \frac{1}{2} (A^1 + A^2) \\ A^{+-} &= \frac{1}{3} A^0 + \frac{1}{2} A^1 + \frac{1}{6} A^2 \end{aligned} \quad (13)$$

Since the effective mass distribution in the Q region is dominated by Q meson production, we shall single out the terms in Eq. (11) with $\ell = 1$, and neglect interference terms in other waves. Thus

$$\begin{aligned}
& |K(t) A_{\pi\pi}(\omega, \cos \Theta_{\pi\pi}) + F_B(\omega, t, \cos \Theta_{\pi\pi})|^2 = \\
& |K(t) 12\pi \frac{\omega}{q} \cos \Theta_{\pi\pi} e^{i\delta_1'} \sin \delta_1' + F_B^{\ell=1}(\omega, t) \cos \Theta_{\pi\pi}|^2 + \quad (14) \\
& + |K(t) A_{\pi\pi}^{\ell \neq 1}(\omega, \cos \Theta_{\pi\pi})|^2 + |F_B^{\ell \neq 1}(\omega, t, \cos \Theta_{\pi\pi})|^2.
\end{aligned}$$

Lacking knowledge of the ω dependence of the F_B terms, we assume that there is no such dependence. On the other hand, we would like to include the t dependence explicitly, since something is known experimentally ^{5,7)}. We choose the empirical forms

$$\begin{aligned}
F_B^{\ell=1}(\omega, t) &= B e^{D_B t/2} e^{i\beta} \\
F_B^{\ell \neq 1}(\omega, t, \cos \Theta_{\pi\pi}) &= B' e^{D_B t/2} \quad (15)
\end{aligned}$$

Similarly, for the t dependence of $K(t)$ we take

$$K(t) = e^{D_g t/2} \quad (16)$$

where D_g is given by experiment ^{5,8)}. Note that the pure OPEM instead of Eq. (16) prescribes the form

$$K(t) = \sqrt{-t} (m_\pi^2 - t)^{-1} e^{D_g t/2} \quad (17)$$

with $D_g = 0$. However, it is well known that this form fails to account for the observed experimental distribution. We may then consider the empirical form (17) with $D_g \neq 0$. This combines the

virtue of explicitly introducing the pion pole at $t = m_\pi^2$, and the zero at $t = 0$, with a good phenomenological shape. On the other hand, the phenomenological shape in Eq. (16) is already very good. In what follows we shall take $K(t)$ to be given by Eq. (16), leaving the detailed arguments to the beginning of Section 5.

Inserting Eqs. (14), (15), and (16) into Eq. (11), and carrying out the integration over $\cos \Theta_{\pi\pi}$, we get

$$\begin{aligned} \frac{d\delta}{d\omega^2 dt} \propto & e^{D_\rho t} \frac{\omega}{q} \sin^2 \delta_1(\omega) + B e^{(D_\rho + D_B)t/2} \cos(\delta_1 - \beta) \sin \delta_1 + \\ & + e^{D_\rho t} \frac{\omega}{q} \Delta_{\pi\pi}(\omega) + \frac{q}{\omega} B_3 e^{D_B t} \end{aligned} \quad (18)$$

Here we have omitted an over-all constant as well as P_{11}^{-2} , which is a constant in any one experiment. The first term in Eq. (18) is the proper ρ meson term, the second term is the interference between the ρ amplitude and the non-OPE amplitude for producing the two final pions in a relative $\ell = 1$ state. The third term is the OPE contribution from partial waves different from P_1 integrated over the angular variables, and the fourth term contains all the non-interfering non-OPE background. For programming purposes, the second term in Eq. (15) may be rewritten as

$$B_1 e^{(D_\rho + D_B)t/2} (\sin^2 \delta_1 + B_2 \sin \delta_1 \cos \delta_1) \quad (19)$$

B_1 , B_2 , and B_3 in Eqs. (18) and (19) are free parameters.

The expression (18) is still not general enough, since all the ρ production is assumed to follow the OPEM, apart from the tribute to phenomenology in Eq. (16). Absorption, initial-state, and final-state interactions may destroy part of the OPE picture.

To account for such possibilities, we allow for unknown energy dependence, multiplying the OPE amplitude, Eq. (10), by an arbitrary function $\sqrt{f(\omega)}$. This corresponds closely to the phenomenological form factors used by Baton and Reignier⁹⁾.

To arrive at the form of Eq. (4), which is a distribution in effective mass, Eq. (18) has to be integrated over t , between the lower limit $-t = -t_{\min}(\omega)$ given by the phase-space contour in the Chew-Low plot [see, for example, Pilkuhn¹⁰⁾], and the upper limit $-t = -T$ of the experimental sample. Furthermore, one has to go from a distribution in ω^2 to a distribution in ω [this gave the factor 2ω in Eq. (2)]. We finally obtain

$$\begin{aligned} \frac{d\delta}{d\omega} \propto & \frac{\omega^2}{q} \left[f(\omega) \sin^2 \delta_1^1(\omega) + \Delta_{\pi\pi}(\omega) \right] \frac{1}{D_g} \int_{-T}^{t_{\min}(\omega)} e^{D_g t} + \\ & + \omega \sqrt{f(\omega)} B_1 \sin \delta_1^1 \left(\sin \delta_1^1 + B_2 \cos \delta_1^1 \right) \frac{2}{D_g + D_B} \int_{-T}^{t_{\min}(\omega)} e^{(D_g + D_B)t/2} + \\ & + B_3 q \frac{1}{D_B} \int_{-T}^{t_{\min}(\omega)} e^{D_B t} \end{aligned} \quad (20)$$

in which

$$\sin^2 \delta_1^1(\omega) = \frac{\omega_r^2 \Gamma_r^2(\omega)}{(\omega_r^2 - \omega^2)^2 + \omega_r^2 \Gamma_r^2(\omega)} \quad (21)$$

The expression (21) is the relativistic¹¹⁾ Breit-Wigner amplitude squared, ω_r is the ρ meson mass, and $\Gamma_r(\omega)$ is the energy-dependent width of the ρ meson, to be discussed in the next section.

It may be instructive to rewrite the expression (20) in the form

$$\frac{d\delta}{d\omega} \propto f(\omega) \frac{\omega^2}{\sqrt{\omega^2 - 4m_\pi^2}} \frac{\omega_r^2 \Gamma_r^2(\omega)}{(\omega_r^2 - \omega^2)^2 + \omega_r^2 \Gamma_r^2(\omega)} \int_{t_{\min}(\omega)}^T K^2(t) dt \quad (22)$$

+ background terms

which corresponds to Eq. (2). An example of a simplified Breit-Wigner formula is now

$$\frac{d\delta}{d\omega} \propto \frac{\Gamma_r(\omega)}{(\omega_r^2 - \omega^2)^2 + \omega_r^2 \Gamma_r^2(\omega)} + \text{empirical background}$$

used in paper I. This clearly implies the tacit assumption

$$f(\omega) \frac{\omega^2}{\sqrt{\omega^2 - 4m_\pi^2}} \cdot \Gamma_r(\omega) \cdot \int_{t_{\min}(\omega)}^T K^2(t) dt = \text{constant}.$$

In paper I this assumption happened to be good, judging from the goodness of the fits obtained (and as a consequence of the functions chosen for the width and the background).

Note that when Eq. (20) is compared with a distribution compiled from experiments done at different beam momenta P_{iL} , the integrals over t have to be weighted appropriately since $t_{\min}$ is a function of P_{iL} .

3. PARAMETRIZATIONS OF THE ENERGY-DEPENDENT WIDTH

Narrow resonances are generally well described by taking $\Gamma(\omega)$ in Eq. (21) to be a constant. This is certainly not true for the broad ρ meson. Here the energy dependence of $\Gamma(\omega)$ is an important factor which strongly affects the shape of the resonance peak in an invariant mass distribution ²⁾. The introduction of an energy dependence is actually equivalent to a phase shift analysis in the region near the resonance. The extent to which an invariant mass distribution can furnish information on the phase shifts, depends of course mainly on statistics. In the resonance region the signal-to-background ratio may be large, but the further one goes away from the resonance position, the more one depends on a good understanding of the background and the production process. A classical example of this procedure is given in the early work on the Δ_{33} resonance ¹²⁾, where the phase shift could successfully be fitted from threshold up beyond the resonance. In the case of the ρ meson we think that with the statistics present we will be able to extract useful information about 300 MeV on each side of the central mass.

Our parametrization of $\Gamma(\omega)$ can best be discussed in terms of the effective range expansion (for further details the reader is referred to the Appendix section). Let us write the relativistic amplitude $a_\ell(\omega)$ for the scattering of two spinless particles in a relative ℓ wave in the form

$$a_\ell(\omega) = \frac{\omega}{q} \frac{e^{2i\delta_\ell} - 1}{2i q^{2\ell}} \quad (23)$$

Since this amplitude is analytic in the q^2 plane, we may take the real part of the inverse amplitude in the physical region and write

$$\frac{q^{2\ell+1}}{\omega} \cot \delta_\ell(\omega) = C_0 + C_1 q^2 + C_2 q^4 + \dots, \quad (24)$$

valid in the region where the elastic unitarity condition holds. Let us define the central mass ω_r of a resonance in the ℓ^{th} wave as the point where the phase shift δ_ℓ goes through $\pi/2$ ¹³⁾. At such a point the expansion on the r.h.s. of Eq. (24) must have a zero. This becomes obvious if we rewrite the r.h.s.

$$a(\omega_r^2 - \omega^2) + b(\omega_r^2 - \omega^2)^2 + \dots \quad (25)$$

For an $\ell = 1$ resonance, keeping only the first two terms in the expansion (25), Eq. (24) becomes

$$\frac{q^3}{\omega} \cot \delta = a(\omega_r^2 - \omega^2) + b(\omega_r^2 - \omega^2)^2. \quad (26)$$

The energy-dependent width, $\Gamma(\omega)$, is most naturally introduced by the definition

$$\cot \delta = \frac{\omega_r^2 - \omega^2}{\omega_r \Gamma(\omega)} \quad (27)$$

In this notation, a scattering amplitude $\exp(i\delta) \sin \delta$, in a partial wave, takes the Breit-Wigner form

$$e^{i\delta} \sin \delta = (\cot \delta - i)^{-1} = \frac{\omega_r \Gamma(\omega)}{\omega_r^2 - \omega^2 - i\omega_r \Gamma(\omega)} \quad (28)$$

In the effective range approximation, for example, neglecting the second term on the r.h.s. of Eq. (26), $\Gamma(\omega)$ becomes

$$\Gamma(\omega) = \Gamma_0 \frac{q^3}{\omega} \left(\frac{\omega_r}{q_r} \right), \quad (29)$$

where Γ_0 is defined as the slope of δ at $\omega = \omega_r$. The constants in the parentheses have been chosen only to permit the interpretation of Γ_0 as a width. We shall, in fact, call Γ_0 the resonance width in what follows.

In the approximation (26), $\Gamma(\omega)$ becomes

$$\Gamma(\omega) = \Gamma_0 \frac{q^3}{\omega} \cdot \left(\frac{\omega_r}{q_r^3} \right) \cdot \frac{1 + R^2 q_r^2}{1 + R^2 q^2} \quad (30)$$

R corresponds to the radius of the region of interaction, well known in nuclear physics [see, for example, Refs. 10) or 14)]. We shall invoke this correspondence only to require R^2 to be positive.

Since higher order terms in the expansion (25) were not needed to fit the Δ_{33} resonance, we shall not introduce them here either.

Although the above parametrization of $\Gamma(\omega)$ seems straightforward due to the natural choice (23) of the partial wave amplitude, it is by no means unique. In fact, one could multiply Eq. (23) by any function which has only the left-hand cut and which does not spoil the threshold behaviour of $a_\ell(\omega)$. Thus an equally good choice for $a_\ell(\omega)$ would be

$$a_\ell(\omega) = \frac{e^{2i\delta_\ell} - 1}{2iq^{2\ell+1}} \quad (31)$$

Repeating the steps which led from Eq. (23) to Eqs. (29) and (30) we find that Eq. (31) leads to a parametrization of $\Gamma(\omega)$ which is a factor ω/ω_r different. Thus in the approximation of Eq. (29) we get

$$\Gamma(\omega) = \Gamma_0 \left(\frac{q}{q_r} \right)^3, \quad (32)$$

and in the approximation of Eq. (30) we get

$$\Gamma(\omega) = \Gamma_0 \left(\frac{q}{q_r} \right)^3 \frac{1 + R^2 q_r^2}{1 + R^2 q^2}, \quad (33)$$

The parameter R in Eq. (33) will, of course, be different from the parameter R in Eq. (30). In the following we shall denote $\Gamma(\omega)$ as given by Eqs. (29) and (30) "relativistic", and as given by Eqs. (32) and (33) "non-relativistic". We shall call the factor

$$\frac{1 + R^2 q_r^2}{1 + R^2 q^2}$$

in Eqs. (30) and (33) "penetration factor" and the parameter R "radius of interaction".

An alternative to the parametrization with a penetration factor is a parametrization in terms of a CDD pole¹⁵⁾. If, as suggested by some dynamical models, the ρ meson is a CDD pole, this implies that the phase shift δ goes through π at an energy ω_0 near the resonance. But when $\delta = \pi$, $\cot \delta$ is singular. Thus a CDD pole behaviour is obtained simply by adding a pole to the r.h.s. of Eq. (26). Exchanging the r.h.s. of Eq. (26) for the simplest algebraic expression with the required properties (zero at $\omega = \omega_r$, pole at $\omega = \omega_0$), we get

$$\frac{q^3}{\omega} \cot \delta = A \frac{\omega_r^2 - \omega^2}{\omega_0^2 - \omega^2} \quad (34)$$

or in next order in ω

$$\frac{q^3}{\omega} \cot \delta = A \frac{(\omega_r^2 - \omega^2)(B - \omega^2)}{\omega_c^2 - \omega^2} \quad (35)$$

Again we can introduce the energy-dependent width, $\Gamma(\omega)$, by the definition in Eq. (27).

Comparing the above expressions with Eq. (27), we see that the whole CDD pole behaviour may be contained in $\Gamma(\omega)$. Thus Eq. (34) is equivalent to Eq. (27) with the choice

$$\Gamma(\omega) = \Gamma_0 \frac{q^3}{\omega} \left(\frac{\omega_r}{q_r^3} \right) \frac{\omega_c^2 - \omega^2}{\omega_c^2 - \omega_r^2} \quad (36)$$

and Eq. (35) is equivalent to Eq. (27) with the choice

$$\Gamma(\omega) = \Gamma_0 \frac{q^3}{\omega} \left(\frac{\omega_r}{q_r^3} \right) \cdot \frac{\omega_c^2 - \omega^2}{\omega_c^2 - \omega_r^2} \cdot \frac{B - \omega_r^2}{B - \omega^2} \quad (37)$$

B is a free parameter. If B is positive the cross-section corresponding to the partial wave amplitude in Eq. (28) has a second maximum at $\omega = \sqrt{B} \equiv \omega_{rc}$. We shall denote ω_c the "CDD zero" and ω_{rc} the "CDD peak". For B infinite the expression (37) clearly simplifies to Eq. (36).

Note that $\omega_c > \omega_r$ corresponds to the case when the phase shift δ in Eq. (34) starts positive at threshold, goes through $\pi/2$ at ω_r and then through π at ω_c . In case $\omega_c < \omega_r$, δ starts negative at threshold, goes through 0 at ω_c and then through $\pi/2$ at ω_r . At ω_{rc} δ must go through $3\pi/2$ or $\pi/2$. In the fits discussed in Section 5 we shall always take

$$\omega_r < \omega_{rc}.$$

Actually it turns out that no acceptable fits with $\omega_c < \omega_r$ can be found. In general, we will therefore have

$$\omega_r < \omega_c < \omega_{rc}.$$

Note that a Breit-Wigner amplitude with a $\Gamma(\omega)$ of the form (37) is mathematically equivalent to a double pole, as well as to the coherent sum of two single poles. For a fuller discussion of this we refer the reader to Section A.3 of the Appendix.

4. DYNAMIC ASSUMPTIONS AND SMALL TERMS

The contribution from OPE $\pi\pi$ scattering in other partial waves than P_1 may easily be written out, using Eqs. (10) and (13). For charged ρ mesons we have

$$\Delta_{\pi\pi}^{\pm}(\omega) = \frac{1}{3} \left\{ \sum_{\ell \neq 1} (2\ell+1) \sin^2 \delta_{\ell}^1 + \sum_{\ell} (2\ell+1) \sin^2 \delta_{\ell}^2 \right\}, \quad (38)$$

where the superscript denotes isospin. However, all of $\pi\pi$ production is not $\pi\pi$ scattering, and therefore some modifications may be expected. In a phase shift analysis of peripherally produced dipions, Baton and Reignier⁹⁾ found it necessary to include phenomenological form factors for each partial wave, to account for the possibility of different attenuation of different partial waves in the production process. Including only S_2 , D_2 and F_1 partial waves, Eq. (38) must then be rewritten

$$\begin{aligned} \Delta_{\pi\pi}^{\pm}(\omega) = & \frac{1}{3} f_0^2(\omega) \sin^2 \delta_0^2 + \frac{5}{3} f_2^2(\omega) \sin^2 \delta_2^2 + \\ & + \frac{7}{3} f_3^1(\omega) \sin^2 \delta_3^1. \end{aligned} \quad (39)$$

In this way, the factor $f(\omega) \equiv f_1^1(\omega)$ appeared in front of the $\sin^2 \delta_1^1(\omega)$ term in Eq. (20).

In the same way the contribution of other partial waves to neutral ρ meson production may be taken into account. Assuming that only S_0 and D_0 waves need to be included, we have

$$\Delta_{\pi\pi}^0(\omega) = \frac{4}{27} f_0^0(\omega) \sin^2 \delta_0^0 + \frac{20}{27} f_2^0(\omega) \sin^2 \delta_2^0. \quad (40)$$

From here on one may proceed in many ways. Either one may set all the form factors $f_l^{\pi}(\omega)$ equal to 1, and use the experimental information about the phase shifts $\delta_l^{\pi}(\omega)$. This task seems hopeless, unless one decides immediately to disregard all partial waves but one, i.e. keeping only one term in each of the Eqs. (39) and (40). In the case of $Q^{\pm}$ the natural choice then is to keep the δ_0^2 term. For δ_0^2 we note two fits, both valid only in a relatively narrow region around the central mass of the Q . They are

$$\cotg \delta_0^2 = -\frac{1}{0.36q} + 7.14 q \quad (\text{in GeV/c}) \quad (41)$$

given by Baton et al.¹⁶⁾ and

$$\cotg \delta_0^2 \approx -0.17 \frac{\omega}{q} + \frac{2}{\pi} \ln \left(\frac{\omega + 2q}{2m_{\pi}} \right) \quad (42)$$

from Wolf¹⁷⁾.

The Q^0 case is somewhat more complicated. There is no doubt that the second term in Eq. (40) is present, due to the f meson. This we may take care of by the assumption that there are no further contributions to the D_0 wave than the f meson with well-known properties.

It is on the other hand well known that δ_0^0 may be large and positive in the Q^0 region. The phase shift analysis of Wolf¹⁷⁾ suggests a resonant δ_0^0 at 720 MeV with a 90 MeV width. More recently Schleim, Malamud¹⁸⁾ and Clegg¹⁹⁾ obtained much slower variation of δ_0^0 in the Q^0 region. Both these results

are consistent with the broad resonance (400 MeV or more) proposed by Lovelace et al. ²⁰⁾ on the basis of the πN backward dispersion relation. Thus we would be obliged to include the first term in Eq. (40) with some parametrization or other for δ_0^0 .

Equation (40) may then be written

$$\Delta_{\pi\pi}^0 = \frac{4}{27} \sin^2 \delta_0^0 + \frac{20}{27} \frac{\omega_f^2 \Gamma_f^2(\omega)}{(\omega_f^2 - \omega^2)^2 + \omega_f^2 \Gamma_f^2(\omega)}, \quad (43)$$

with the energy-dependent width $\Gamma_f(\omega)$ of the f meson given by the analogy to Eq. (29)

$$\Gamma_f(\omega) = \Gamma_{0f} \frac{\omega_f}{\omega} \left(\frac{q}{q_f} \right)^5, \quad (44)$$

and with δ_0^0 suitably parametrized. In Section 5 we shall specify what parametrizations we use for δ_0^2 in Eq. (39) and for δ_0^0 in Eq. (43).

A completely different approach is to disregard $\Delta_{\pi\pi}(\omega)$ altogether, and let all corrections due to other waves than P_1 be taken care of by the P_1 form factor $f(\omega)$ and by the background. In that case, $f(\omega)$ seems best fitted by the form

$$f(\omega) = c_1 + c_2 \omega^{-2}, \quad (45)$$

higher order terms not being required. We shall shortly return to this dynamic assumption.

To discuss the interference term, let us rewrite Eq. (18) in the form

$$\frac{d\sigma}{d\omega^2 dt} \propto \frac{\omega}{q} e^{D_\varphi t} \sin^2 \delta'_1 \left[1 + \frac{q}{\omega} B e^{(D_B - D_\varphi)t/2} \left[\sin \beta + \frac{\omega_r^2 - \omega^2}{\omega_r \Gamma(\omega)} \cos \beta \right] \right] \quad (46)$$

+ (non-interfering background).

Note that $D_\varphi > D_B$ and that therefore the interference is more pronounced at higher $|t|$ values. Furthermore, the term containing $\sin \beta$ has a weaker ω dependence than the $\cos \beta$ term. Thus the shape of the resonance will be more seriously modified by the interference term for smaller β . In this way, the φ meson parameters may be appreciably shifted by interference, as noted by Söding²¹⁾ in connection with photoproduction of φ .

If one expands the square bracket in Eq. (46) in terms of ω , the dominant terms are $c_1 + c_2 \omega^{-2}$ in the approximation $\Gamma(\omega) \approx q^3/\omega \approx \omega^2$. This shows that the whole bracket containing the interference term can be approximated by a function $f(\omega)$ of the form of Eq. (45) multiplying the φ meson term. This is an alternative interpretation of the function $f(\omega)$. Note, however, that if the interference term is appreciable it also adds some t dependence to $f(\omega)$, and ultimately to the φ parameters. Thus it is not very surprising that the models of Söding²¹⁾ and of Ross and Stodolsky²²⁾ were both able to account for the apparently low mass values of photo-produced φ^0 , although the former introduces an interference with background, and the latter shows, on the basis of the photon dissociation model, that a factor $1/\omega^4$ is present in the $f(\omega)$ function.

We may then conclude from Sections 3 and 4 that the function $f(\omega)$ effectively simulates

- i) final or initial state corrections to the P_1 partial wave in OPEM,
- ii) other partial waves in OPEM (when they are small and slowly varying),
- iii) interference terms between ϕ and background, the Söding model ²¹⁾,
- iv) energy-dependent factors of the kind used in the Ross and Stodolsky model ²²⁾,
- v) the CDD pole model, since the choice $c_1 < 0$, $c_2 > 0$ in Eq. (45) produces a zero in $f(\omega)$ at the positive energy

$$\omega = \sqrt{-\frac{c_1}{c_2}}.$$

Truly, a zero in $f(\omega)$ is something quite different from a zero in the cross-section $\sin^2 \delta_1^1$, but the net result is in both cases a zero in the invariant mass distribution, due to the presence of the term

$$f(\omega) \sin^2 \delta_1^1(\omega)$$

in Eq. (20).

5. ANALYSIS OF THE EXPERIMENTAL DISTRIBUTIONS

The compilation of experimental data to be used is much the same as in paper I, however with some important additions and changes. The addition is the very large Saclay experiment¹⁶⁾ with 2.77 GeV/c π^- . The change is in another large π^- experiment, as yet unpublished, which contained some errors in the version used in paper I. Altogether, we shall analyse seven different histograms in this section. Data on these histograms are given in Table 1 below. The statistical methods are the same as discussed in paper I.

The experimental distributions of events versus t may well be fitted by either one of the two phenomenological forms (16) and (17). Using Eq. (16) we find that D_0 is about 9 ± 1 (GeV/c)⁻² independently of beam momentum^{5,8,16)}. A variation in D_0 of about 1 (GeV/c)⁻² turns out to have a negligible effect on the Q parameters. The reason is of course that we do not use the full information in the Chew-Low plot, but only the projections on the mass axis. The strong t dependence of the integrand in

$$\int_{t_{\min}(\omega)}^T K^2(t) dt \quad (47)$$

is integrated out, leaving only a weak ω dependence and the T dependence to be sensed in the invariant mass distribution. If instead of Eq. (16) one would fit the experimental t distribution with Eq. (17), the integral (47) would differ by at most 3% throughout the t region of interest $\sqrt{s} - D_0$ in Eq. (17) is required to be slightly larger than D_0 in Eq. (16)]. The practical difference between Eqs. (16) and (17) is therefore completely negligible. We have chosen to use Eq. (16) for reasons of simpler programming, whereas the virtues of Eq. (17) are entirely aesthetical.

The constant D_B in Eq. (15) seems to be about $2 (\text{GeV}/c)^{-2}$, independently of beam momentum ⁵⁾. We have also found that a variation in D_B of $1 (\text{GeV}/c)^{-2}$ has a negligible effect on the parameters. The values we have chosen are $D_B = 2$, $D_{g^-} = 9.25$, $D_{g^0} = 10.25$, $D_{g^+} = 11.25$, all in units of $(\text{GeV}/c)^{-2}$.

5.1 Comparison of simplified Breit-Wigner functions

Since simplified Breit-Wigner functions have had much success in describing resonance peaks of low statistics, we have tried to use Eq. (20) neglecting several different combinations of energy-dependent factors. In Eq. (20) with $\Delta_{\pi\pi}(\omega) = B_1 = 0$ we set the product of the unknown $f(\omega)$ with every combination of the three factors

$$\omega, \quad \Gamma(\omega), \quad L(\omega, T) \equiv \frac{\omega}{q} \int_0^{t_{\min}(\omega)} \exp(D_g t) \quad (48)$$

equal to a constant. We further assume $\Gamma(\omega)$ to have the simplest possible energy dependence, Eq. (29), as is usually done in simplified Breit-Wigner formulae [unless $\Gamma(\omega)$ is taken to be a constant which is clearly wrong]. As a test sample we use SACL 13⁻ (see notation in Table 1) which has the advantage of large statistics at a unique $\pi\pi$ momentum.

Some of the eight resulting solutions are listed in Table 2. Those not listed have $\chi^2 > 150$ for about 77 degrees of freedom. Solution 1 (cf. Table 2) shows that the complete, unmodified distribution (20) [with $f(\omega) = 1$, $\Delta_{\pi\pi}(\omega) = B_1 = 0$ and $\Gamma(\omega)$ given by Eq. (29)] is unacceptable. Solutions 2 and 3 are at the

4% and 8% confidence level. If the simplified Breit-Wigner functions in solutions 2 or 3 (see Table 2) are combined with suitable empirical background functions, one can expect them to fit the data with even higher confidence. This was exactly the procedure in paper I, where a function of the kind of solution 3 was used [see also discussion after Eq. (22)]. For the sample SACL 13⁻, we then obtained a 47% confidence level.

Comparing solutions 2 and 3 we see that the systematic differences in ω_r and Γ_0 are of the order of 20 MeV.

In conclusion we find that energy-dependent modifications are required and that the success of some simplified Breit-Wigner functions is a consequence of ignoring a fortuitous combination of energy-dependent factors.

5.2 Comparison of different $\Gamma(\omega)$ and $\Delta_{\pi\pi}(\omega)$

Since the simplified functions above are approximate, we shall not use them anymore. It is, however, clear from the poor χ^2 of solution 1 that the complete, unmodified distribution (20) has to be modified in one way or another. We shall next test two expressions for $\Gamma(\omega)$: Eqs. (29) and (30); and four expressions for $\Delta_{\pi\pi}(\omega)$: 0, constant, Eq. (41) and Eq. (42).

It turns out that all solutions with the simplest $\Gamma(\omega)$, Eq. (29), are completely unacceptable with $\chi^2 > 330$. In Table 2 we therefore only list the solutions with the energy dependence of $\Gamma(\omega)$ given by the penetration factor formula, Eq. (30).

Clearly both the Baton¹⁶⁾ type $\Delta_{\pi\pi}^+(\omega)$, solution 4, and the Wolf¹⁷⁾ type $\Delta_{\pi\pi}^-(\omega)$, solution 5, are unacceptable. Not only is χ^2 much too large, but to achieve even this χ^2 the

background becomes negative. The trouble mainly appears in the wings, outside the region where the functions of Baton ¹⁶⁾ and Wolf ¹⁷⁾ were originally determined.

Solution 6 is the first good solution found. It requires a $\Delta_{\pi\pi}^-$ term corresponding to a constant phase shift

$$|\delta_0^2| = 0.05 \begin{matrix} +0.2 \\ -0.05 \end{matrix}, \text{ (in radians)}$$

consistent with zero. In solution 7 (see Table 2) we therefore set $\Delta_{\pi\pi}^- = 0$ and obtain a good fit. The R^2 value, in units of $(\text{GeV}/c)^{-2}$, corresponds to a radius of interaction of

$$R = 0.30 \begin{matrix} +0.034 \\ -0.024 \end{matrix} \text{ f.}$$

Thus we find that nothing can be gained in the description of the invariant mass distribution by parametrising $\Delta_{\pi\pi}^-(\omega)$ in terms of anything more complicated than a constant. This result is supported by the facts that no other charged $\pi\pi$ resonances are seen in invariant mass distributions in the ρ region, and that all phase shift analyses of angular distributions prefer slowly varying and small $\delta_0^2(\omega)$.

We also conclude that if $f(\omega)$ is assumed constant, the energy dependence of $\Gamma(\omega)$ cannot be given by Eq. (29) whereas Eq. (30) is acceptable.

5.3 Comparison of different $f(\omega)$ and CDD poles

The effect of including a CDD zero for which $\Gamma(\omega)$ is given by Eq. (36) is shown in solution 8. χ^2 is not significantly worse than for solution 7 (see Table 2) and the ρ parameters are essentially unchanged. The CDD zero wants to be located between 1.3 and 1.4 GeV (see Table 2) which is just above the region of the experimental histogram (the fit uses 10 MeV bins up to 1.3 GeV).

We have also tried to introduce a second peak (CDD peak) using Eq. (37) for $\Gamma(\omega)$. This did not change solution 8 at all and the CDD peak preferred to move out to infinity (at least 80 GeV). Note that this second peak could correspond to the second ρ (ρ') to account for the πp charge exchange polarization²⁴⁾. The present analysis which extends up to 1.3 GeV can of course neither prove nor disprove its existence, since data at much higher energies are needed. However, if the CDD pole is present then the mass of the ρ' must be higher than the position of the CDD zero (the phase shift must in this model go first through π and then reach a second resonance). If the ρ' exists and has a mass $\gtrsim 1.3$ GeV it is probably highly inelastic and our parametrization does not apply, since it is based on the elastic unitarity for $T = J = 1$.

We note that no acceptable solution could be found for $\omega_c < \omega_r$ [cf. discussion after Eq. (37)].

Next we have tried some one-parameter energy expansions of $f(\omega)$. With $\Gamma(\omega)$, again given by Eq. (29) and $B_1 = \Delta_{\pi\pi} = 0$ still, we found a good solution 9 and a very good solution 10 (see Table 2). Note that solution 10 corresponds to the cross-section (20) having a zero somewhere between 1.6 and 2.6 GeV, and solution 9 to a zero at about 1.4 GeV (error not calculated).

Since χ^2 is almost the same for solution 9 with $f(\omega) = f_0 + \omega^{-1}$ and solution 10 with $f(\omega) = f_0 + \omega^{-2}$, we also ran the program with $f(\omega) = f_0 + f_1 \omega^{-1} + \omega^{-2}$ to determine which of the terms was more important. The solution was identical to solution 10, and the value of f_1 came out to be

$$f_1 = \left(0 \pm \begin{matrix} 7 \\ -2 \end{matrix} \right) \text{ GeV}.$$

From the ten discussed solutions to the Saclay sample (SACL 13⁻) we conclude that

- a) $\Delta_{\pi\pi}^- = 0$, i.e. other partial waves than P_1 are negligible throughout the energy region analysed;
- b) interference term is not required ($B_1 = 0$);
- c) there are three equally favoured formulae:
 - i) penetration factor, Eq. (30), with $f(\omega)$ constant,
 - ii) CDD pole, Eq. (36), with $f(\omega)$ constant,
 - iii) simple $\Gamma(\omega)$, Eq. (29), with $f(\omega) = f_0 + \omega^{-2}$ (as noted in Section 4, this formula simulates a CDD pole).

Note that we still have not compared the different parametrizations (30) and (33) of $\Gamma(\omega)$ with penetration factor. As we shall do this on a larger sample than SACL 13⁻, we postpone the discussion to Section 5.5.

5.4 Determination of the systematic errors introduced by adding several experiments together

It is generally believed that by adding several experiments together one introduces some terrible biases or fudge factors. In paper I we subscribed to this view, trying to discredit wide ρ peaks

on the motivation that the broadening might be due to systematic errors in the energy scale of the individual experiments. The true state of affairs now appears from comparing the mono-energetic sample SACL 13⁻ with the independent sample COMP 10⁻, a compilation of seven experiments at thirteen different momenta, ranging from 1.59 GeV/c to 3.22 GeV/c. Solution 11 gives the penetration factor fit to COMP 10⁻ to be compared with solution 7 and solution 12 gives the CDD pole fit to be compared with solution 8 (see Table 2). The shift of the parameters in the two comparisons amounts to:

<u>Penetration factor, Eq. (30)</u>	<u>CDD pole, Eq. (36)</u>
$\Delta\omega_r = (0.9 \pm 2.8) \text{ MeV}$	$(0.9 \pm 2.8) \text{ MeV}$
$\Delta\Gamma_o = (2.0 \pm 8.0) \text{ MeV}$	$(3.0 \pm 8.0) \text{ MeV}$
$\Delta R^2 = (0.6 \pm 0.9) (\text{GeV}/c)^{-2}$	
$\Delta\omega_c =$	$(59.0 \pm 56.0) \text{ MeV}$
$\Delta(\text{background}) = (1.4 \pm 2.3) \%$	$(0.1 \pm 2.5) \%$

Since all these shifts are manifestly consistent with zero, evidently no systematic errors arise from the addition of the experiments. We then proceed to determine the ϕ^- parameters from the combined sample of Saclay and the previous compilation. This gives us TOT 10⁻, a sample of 12773 events.

5.5 Determination of $\Gamma(\omega)$ parametrization errors

In Eqs. (32) and (33) we showed that $\Gamma(\omega)$ only is defined up to a factor ω/ω_r . Actually Eqs. (29) and (30) are defined only up to a factor $(\omega/\omega_r)^n$, where n is any number. To determine the effect of this freedom we have studied the cases $n = -1, 0, 1$ on

the sample TOT 10⁻. Solution 13 gives the case $n = -1$, solution 14 the case $n = 0$ [corresponding to Eq. (30)], and solution 15 the case $n = 1$ [corresponding to Eq. (33)].

All the solutions are good and strikingly similar. The variation in ω_r and Γ_o over the three solutions is

$$\Delta \omega_r = (1.4 \pm 1.5) \text{ MeV}$$

$$\Delta \Gamma_o = (1.1 \pm 3.9) \text{ MeV.}$$

There is no acceptable solution with $R^2 = 0$, corresponding to Eqs. (29) and (32). As can be seen from Table 2 the variation in R^2 with n is very strong. Thus one may have the hope of determining the appropriate power n from some independent source of information on R^2 , for instance through the relation between R^2 and the scattering length at threshold which may be calculated using current algebra techniques or dispersion relations.

It is interesting to compare solution 14 to solution 16, which is a CDD pole fit to the same sample. The variation in the parameters is again quite small

$$\Delta \omega_r = (1.9 \pm 1.5) \text{ MeV}$$

$$\Delta \Gamma_o = (1.7 \pm 4.0) \text{ MeV.}$$

Thus we can conclude that the parametrization errors [defined in the Introduction as systematic errors due to the freedom in choice of parametrization for $\Gamma(\omega)$] are in general quite small, of the order of the statistical error in our samples. To be definite, the parametrization error δ_{par} is

$$\delta_{\text{par}} \approx 1 \text{ MeV} \quad (49)$$

if we take it as half of the variations $\Delta \omega_r$, $\Delta \Gamma_o$ above.

5.6 Determination of the ϕ^- parameters

For an absolute determination of the ϕ^- parameters it is necessary to fold in a 30-MeV wide Gaussian resolution into all the theoretical distributions. This we have not done until now because the folding increases considerably the time for computing MINUIT errors, and because we were only interested in comparing differences of parameter values in different solutions.

Solutions 17 and 18 (cf. Table 2) are identical, respectively, to solutions 14 and 16 except for the 30 MeV resolution function. If the ϕ^- meson had a pure Gaussian shape of width 150 MeV, one would expect a resolution of 30 MeV to decrease this width by

$$(150 - \sqrt{150^2 - 30^2}) \text{ MeV} \approx 3 \text{ MeV.} \quad (50)$$

Indeed, solution 17 yields a 4.3 MeV narrower width than solution 14 and solution 18 a 4.6 MeV narrower width than solution 16, in reasonable agreement with this.

The statistics of the sample TOT 10^- is now so large that it starts to be significant to compare the three best formulae, i), ii) and iii) of Section 5.3 on the basis of χ^2 . The penetration factor formula, $\Gamma(\omega)$ given by Eq. (30), solution 17, and the formula using Eqs. (29) and (45), solution 19, give about equally good fits (cf. Table 2), confidence level 51-53 %. On the other hand, the CDD pole hypothesis, solution 18, does not fit quite so well, confidence level 24 %. Of course, this in no way rules out the CDD pole hypothesis; one can only conclude that it is not necessary for the ϕ^- meson. In Fig. 2 we have drawn solutions 17, 18 and 19 on the experimental histogram TOT 10^- . In fact the three curves coincide within the width of the pencil !

Comparing the values of ω_r and Γ_0 in solutions 17 and 19 which differ in dynamical assumptions, we see that the dynamical error (defined in the Introduction as the systematic error due to difference in dynamical assumption) is of the order of 10 MeV in ω_r and 7 MeV in Γ_0 . A priori we have no way of deciding which hypothesis is right, the one leading to solution 17 or the one leading to solution 19. One procedure to determine the Q^2 parameters would be to take the average of these solutions and assign errors equal to half the dynamical errors (plus parametrization error and statistical error). A better procedure, we think, is the following.

We combine the hypotheses in solutions 17 and 19 taking $\Gamma(\omega)$ to be given by Eq. (30) and at the same time $f(\omega)$ to be given by Eq. (45). In addition, we admit background interference and finally we fold in a 30 MeV resolution. There are now so many free parameters that they become strongly correlated and their errors become large. This then has the same effect as to add systematic errors and statistical errors. To be sure that we do not fall in a local minimum and get too small errors, we have let the computer determine the two standard deviation errors (χ^2 increased by 4). The errors we quote in Table 2 for the resulting solution 20 are graphically interpolated one standard deviation errors, obtained using all the MINUIT output points. As could have been expected χ^2 as a function of each parameter is not at all parabolic but quite complicated. As an example we plot in Fig. 3 χ^2 as a function of ω_r . The curve clearly shows the two different minima due to solutions 17 and 19. The maximum between the minima is sufficiently high to force the one standard deviation error to be given by the local minimum due to solution 17. However, already for the 1.6 standard deviation error one is obliged to take the full latitude of the two minima into account.

When finally the parametrization errors from Eq. (49) are added quadratically to the errors from Fig. 3, we find the parameters

$$\begin{aligned} M_{\rho^-} &\equiv \omega_r = 764.3 \text{ MeV} \\ \Gamma_{\rho^-} &\equiv \Gamma_o = 147.3 \text{ MeV} \end{aligned} \quad (51)$$

with the one standard deviation errors

$$\delta_{\omega_r}^{(1)} = \begin{matrix} +1.9 \\ -1.8 \end{matrix} \text{ MeV}, \quad \delta_{\Gamma_o}^{(1)} = \begin{matrix} +4.0 \\ -3.9 \end{matrix} \text{ MeV}$$

and the two standard deviation errors

$$\delta_{\omega_r}^{(2)} = \begin{matrix} +19.2 \\ -3.3 \end{matrix} \text{ MeV}, \quad \delta_{\Gamma_o}^{(2)} = \begin{matrix} +8.2 \\ -7.8 \end{matrix} \text{ MeV}.$$

Note that the ω_r values of solutions 17 and 19, as well as that of paper I, are comfortably inside the errors.

The background interference term is not really necessary, as can be seen from the value of B_1 in Table 2. This is even more evident if one studies the values of B_1 inside the hypersurface $\chi^2(\omega_r, \Gamma_o, R, f_o, f_2, B_1, B_2, B_3) = \chi_{\min}^2 + 4$. The value $B_1 = 0$ is in many regions well inside this two standard deviation hypersurface.

5.7 T dependence of the ρ^- parameters

Although the more empirical approach in paper I seems to have well taken care of the energy-dependent shape of the resonance peak, it was not possible to account there for any t dependence. As a result, perhaps, we found that the $\rho^\pm$ mass was T dependent.

5.9 Determination of the ϕ^0 parameters

The ϕ^0 sample COMP 10⁰, Table 1 and Fig. 4, presents some new problems for the analysis. Firstly, this sample may contain the decay

$$\omega \rightarrow \pi^+ \pi^- \quad (53)$$

shown to be a one standard deviation effect in the reaction

$$\pi^+ p \rightarrow \pi^+ \pi^- \Delta^{++}$$

in paper I. The conclusions in paper I concerning the presence of the decay (53) in the reaction

$$\pi^- p \rightarrow \pi^+ \pi^- n \quad (54)$$

are no longer valid due to the error mentioned at the beginning of this section.

Secondly, the sample will certainly contain some decays

$$f^0 \rightarrow \pi^+ \pi^- \quad (55)$$

which we hope to account for by the second term in Eq. (43).

Thirdly, the sample COMP 10⁰, although mainly of reaction (54), also contains an important fraction of events of reaction

$$\pi^+ d \rightarrow \pi^+ \pi^- p p \quad (56)$$

Our background corresponds to phase space for reaction (1), or equivalently reaction (54). It is therefore certainly incorrect at the upper end of the mass distribution, in particular since the reaction (56) is represented by an experiment at 5 GeV/c, whereas all experiments of reaction (54) are at momenta not exceeding 4 GeV/c (see paper I, Tables 1 and 3).

Finally, Fig. 4 contains a strange bump at about 475 MeV, which we do not know what to do with. It may contain misidentified events of the types

and

For all these reasons, but the first, we have extended our ρ^0 analysis over a somewhat narrower energy region than the ρ^- analysis covering about 600 MeV.

Solution 24, Table 2, gives the best fit with $\Delta_{\pi\pi}^0(\omega) = B_1 = 0$ to the neutral sample COMP 10⁰. This solution uses a $\Gamma(\omega)$ of the form (30) and contains an incoherently added contribution from an ω meson Breit-Wigner formula with a constant width. Comparing this solution with solution 17, we make an interesting observation.

The presence of an incoherent peak at mass 783.4 MeV and of width 12 MeV in the ϕ^0 compilation does still have the significance reported in paper I, but it is evident from Fig. 2 that an even more significant peak is present in the charged sample. In this situation it is no longer possible to conclude that the peak is due to the decay (53). Thus the observation in paper I that there is some asymmetry in the isospin 1 channel in the peak region is still true. (Note that the excess in Figs. 2 and 4 is in the region 770-800 MeV and that there is a dip at 815 MeV in both. We do not try to analyse this asymmetry in the present paper.)

Next we must include $\Delta_{\pi\pi}^0(\omega)$, Eq. (43). Solution 25 assumes δ_0^0 to be a constant; the resulting value is

$$\delta_0^0 = 44^\circ \pm 2^\circ. \quad (57)$$

The solution wants a negative background and we have therefore fixed it at $B_3 = 0$. This seems rather artificial: it is obvious that no clear separation between background and constant $\Delta_{\pi\pi}^0$ term is possible.

In solution 26 we try a linearly energy-dependent phase shift δ_0^0 with the result

$$\delta_0^0 = \left(1.1^{+0.3}_{-0.2}\right) - \left(0.8^{+0.4}_{-0.5}\right) \frac{\omega}{[\text{GeV}]} \text{ [radians]}. \quad (58)$$

This solution is slightly better than the already good solution 25. The background is now positive but small, thus the same comment about the inseparability of $\Delta_{\pi\pi}^0$ and background applies. Another sign that the background is not taken properly care of is that Γ_0 is larger than for the ϕ^- meson.

As a third possibility we try a resonant S_0 term of the form

$$\sin^2 \delta_0^o(\omega) = \frac{\omega_\delta^2 \Gamma_\delta^2}{(\omega_\delta^2 - \omega^2)^2 + \omega_\delta^2 \Gamma_\delta^2}, \quad (59)$$

inserted into Eq. (43). The results are given in solution 27, Table 2. The resonance parameters become

$$\omega_\delta = (760 \pm 115) \text{ MeV}$$

$$\Gamma_\delta = (460 \pm 110) \text{ MeV.}$$

The ρ^0 width $\Gamma_\rho = (150 \pm 10) \text{ MeV}$ is now in agreement with the ρ^- width, but the background is still very small and consistent with zero.

We conclude that we cannot separate $\Delta^0 \pi\pi$ from background and we can therefore not decide whether a S_0 term is present or not. The invariant mass distributions can accommodate a resonant S_0 wave with a width of 350 MeV or more, but certainly no narrow resonance in the region studied. On the other hand, no such resonant S_0 wave is required.

Assuming that the background is similar in reactions (1) and (54) and that the $\Delta \pi\pi$ term is negligible in both cases, one can determine the mass difference between ρ^0 and ρ^- . Comparing the samples TOT 10^- and COMP 10^0 fitted with the same formula, as is done in solutions 17 and 24, we find

$$M_{\rho^0} - M_{\rho^-} = (2.4 \pm 2.1) \text{ MeV.} \quad (60)$$

Note that this error includes the systematic uncertainty whether the ω meson, reaction (53), is present or not. This uncertainty is already accounted for in the ω_r error in solution 24, Table 2.

The value (60) is different from the value in paper I mainly due to the changes and additions to the data compilation.

6. RHO MESON WIDTH AND THE PIONIC FORM FACTOR

In a pioneering experiment with colliding beams ³⁾ the ρ^0 meson has been produced in the reaction

$$e^+ e^- \rightarrow \rho^0 \rightarrow \pi^+ \pi^- \quad (61)$$

with a reported width of (93 ± 15) MeV. This is in serious conflict with our results, 150 ± 10 MeV for reaction (54) and $(146.7 \pm 4.9 - 4.1)$ MeV for reaction (1). In our opinion there is no hope for a major change in our results for the pion produced ρ mesons. Minor changes can of course still occur if one, for instance, makes some specific assumptions about the production processes leading to the background events. In this section we shall briefly discuss the pionic form factor which is the quantity measured by the Novosibirsk group ³⁾.

The pionic electromagnetic form factor F_π is usually described by formulae of two different kinds. The first kind expresses F_π in terms of the Omnès function ²⁶⁾ which is the most general formula with the correct analyticity properties. The second kind relates F_π directly to the P_1 wave scattering amplitude ^{27,28)} by a Breit-Wigner type expression. Here simplicity is achieved at the price of getting poles in the complex t plane. Common for both kinds is that they must be normalized to 1 at $t = 0$ (in this section t denotes the square of the four-momentum of the virtual photon). The analysis leading to the 93 MeV width for the Novosibirsk data ³⁾ is based on the latter type of formula, however, not normalized to unity at $t = 0$. In the present section we shall discuss the Omnès function ²⁶⁾ approach and compare it with the Novosibirsk data ³⁾.

The electromagnetic form factor F_π is an analytic function in the complex t plane with a cut extending from the threshold $t_1 = 4m_\pi^2$ to infinity. Defining the phase $\varphi(t)$ of F_π on the real axis by

$$F_\pi(t+i\epsilon) = |F_\pi(t+i\epsilon)| e^{i\varphi(t)}, \quad t > t_1, \quad (62)$$

F_π takes the form ^{26,29)}

$$F_\pi(t) = P_n(t) \exp \left[\frac{t}{\pi} \int_{t_1}^{\infty} \frac{\varphi(t') dt'}{t'(t'-t)} \right]. \quad (63)$$

$P_n(t)$ is an arbitrary polynomial subject to the requirement $P_n(0) = 1$.

In the region where only the $\pi\pi$ intermediate state contributes to the unitarity condition for $F_\pi(t)$ and where the $\pi\pi$ scattering is purely elastic

$$t_1 < t < t_H,$$

the phase $\varphi(t)$ is given by the P_1 wave $\pi\pi$ phase shift

$$\varphi(t) = \delta_1^1(t). \quad (64)$$

In the region $t > t_H$ very little can be said about $\varphi(t)$. Therefore, the Novosibirsk experiment ³⁾ can shed light on the P_1 phase shift $\delta_1^1(t)$ only if

- i) t_H is well above the φ region,
- ii) the behaviour of $\varphi(t)$ above t_H is not very dramatic,
- iii) something is assumed about $P_n(t)$. One usually assumes $P_n(t)$ to be equal to 1 although this assumption is not justified²⁹⁾.

Some support for condition i) is lent by the absence of inelastic effects in the φ channel: the φ meson does not decay into four pions and our analysis of pion produced φ mesons does not reveal any fast variations in $\delta_1^1(t)$ below 1.3 GeV. In the following we shall take t_H to be

$$t_H = (1.5 \text{ GeV})^2.$$

About condition ii) nothing is known. If we assume $\varphi(t)$ to be constant

$$\varphi(t) \rightarrow C \quad \text{for } t \rightarrow t_H \quad (65)$$

then

$$F_\pi(t) \rightarrow C' t^{-c/\pi} \quad (66)$$

Clearly C cannot be negative. If instead one wants $F_\pi(t)$ to decrease exponentially³⁰⁾ $\varphi(t)$ is necessarily an increasing function. We shall take $\varphi(t)$ to be given by (64) and (65). The parametrization we choose corresponds to Eqs. (27) and (30).

Finally, about $P_n(t)$ we make the customary assumption

$$P_n(t) = 1, \quad (67)$$

since one anyway cannot afford more parameters for the few experimental points known. As a result we find for the ρ^0 parameters

$$\Gamma_0 = (122 \pm 8) \text{ MeV}$$

$$\omega_r = (764 \pm 7) \text{ MeV}.$$

R prefers to move out to very large values, so that the penetration factor effectively becomes q_r^2/q^2 which corresponds to taking $n = -2$ in Section 5.5. The fit is, however, not very good with a $\chi^2/\langle\chi^2\rangle = 2.28$. This is entirely due to the first experimental point which contributes 1.68 to $\chi^2/\langle\chi^2\rangle$. The value of $F_\pi(t)$ at the first point is less than 1, which is very strange since $F_\pi(0) = 1$ and the first point is at $t = (0.58 \text{ GeV})^2$. Perhaps one has to do here with a statistical fluctuation which will disappear when the number of events at this energy increases from the present two or three. If not, one may have to consider less simple assumptions on $P_n(t)$ than Eq. (67) or one has to demand $\delta_1^1(t)$ to be negative in this region [which is contradicted by our study of CDD poles with $\omega_c < \omega_r$; cf. discussion after Eq. (37)]. Good fits to the Novosibirsk data ³⁾ using the Omnès function ²⁶⁾, Eq. (63), with this type of behaviour for $\delta_1^1(t)$ has been obtained by Bergia and Mignaco ³¹⁾. However, their parametrization of $\delta_1^1(t)$ corresponds to introducing odd powers of q in the expansion on the r.h.s. of Eq. (24) for which we see no physical justification.

Finally it should be mentioned that our fit is extremely insensitive to variations in c , the constant in expression (65). This is in contrast to the observation of Antoniou and Bowcock ³²⁾, perhaps because our choice of t_H is much higher than theirs.

The low value of $F_\pi(t)$ at the first point, $t = (0.58 \text{ GeV})^2$, may on the other hand be taken as an indication that t_H could be very small. If it turns out that experiment requires $F_\pi(0) < 1$ (in fits with the condition $F_\pi(0) = 1$ relaxed) this may indicate that reaction (61) proceeds through other intermediate states, as for instance the 4π state, in addition to the $\pi\pi P_1$ state. We are indebted to M. Jacob for this point.

7. CONCLUSIONS

We have derived a general expression for the invariant mass distribution of the two final pions in the reaction $\pi p \rightarrow \pi\pi N$, assumed to be dominated by the peripheral production of ϱ mesons. The t dependence of the ϱ meson and background production processes is accounted for by the phenomenological formulae (15) and (16). The possibility of interference between the ϱ meson amplitude and the non-OPE background term is considered. The general formula also contains terms due to $\pi\pi$ scattering in other waves than P_1 .

It is shown that this general formula (20) contains as special cases the simplified "Breit-Wigner formulae" commonly used in analyses of experiments of low statistics. Comparing our general formula with a large compilation of invariant mass histograms, we find that it cannot describe the experimental situation well unless some energy-dependent modifications are introduced. There are three main modifications of the ϱ meson amplitude which lead to successful descriptions. The first one consists in introducing a "radius of interaction" as a new parameter in the expression for the energy-dependent width $\Gamma(\omega)$. The second modification also introduces a new parameter into $\Gamma(\omega)$ by assuming the ϱ meson to be a CDD pole. The third modification assumes the ϱ amplitude to be multiplied by an energy-dependent form factor which need not have more than one free parameter. Finally, there is some further liberty in the parametrization of $\Gamma(\omega)$ which is related to the radius of interaction.

The $\pi^-\pi^0$ distributions require no S_2 wave term (this form of analysis is not sensitive to additional waves) and the $\pi^+\pi^-$ distributions require no S_0 wave term, when a D_0 wave corresponding to the f meson is included. On the other hand, an S_0 wave term

cannot be excluded and a resonance at (760 ± 115) MeV with a width of (460 ± 110) MeV can well be accommodated.

Comparing in detail a large sample of ϕ^- mesons from one experiment ¹⁶⁾ done at one momentum and an equally large compilation of data from several experiments done at several different momenta, we conclude that no systematic error is introduced by the addition of experiments in the range 1.6 to 3.2 GeV. In consequence, the procedure used in paper I ¹⁾ to evaluate the ϕ meson width is not needed now and we find the new parameter values

$$M_{\phi^-} = 764.3 \begin{matrix} + 1.9 \\ - 1.8 \end{matrix} \begin{matrix} (19.2) \\ (3.3) \end{matrix} \text{ MeV}$$

$$\Gamma_{\phi^-} = 147.3 \begin{matrix} + 4.0 \\ - 3.9 \end{matrix} \begin{matrix} (8.2) \\ (7.8) \end{matrix} \text{ MeV.}$$

The errors include the true statistical errors evaluated by program MINUIT as well as all systematic errors due to ambiguities in the choice of $\Gamma(\omega)$ parametrization or dynamic assumptions. The errors in brackets next to the 1 standard deviation errors are the 2 standard deviation errors.

The previously observed ¹⁾ dependence of the $\phi^\pm$ mass on T , the upper momentum transfer limit of the sample, is now entirely explained.

The $\phi^0 - \phi^-$ mass difference is found to be

$$M_{\phi^0} - M_{\phi^-} = (2.4 \pm 2.1) \text{ MeV.}$$

Finally we show that the width of the e^+e^- produced ϕ^0 meson, as evaluated from the Novosibirsk data ³⁾, is of the order of 120 MeV. The difference between this result and the published ³⁾ width 93 MeV is entirely to be sought in the difference in the theoretical formula used for the pionic form factor.

ACKNOWLEDGEMENTS

It is a pleasure to acknowledge the hospitality shown to one of us (J.P.) by Professor J. Prentki and the staff of the Theory Division, CERN. We have profited from stimulating discussions with and valuable criticism from Drs. M. Jacob, CERN, H. Pilkuhn, University of Karlsruhe, J.T. Donohue, J.A. Mignaco and C.B. Chiu, CERN. We also thank Drs. J.P. Baton, Saclay, V.A. Sidorov, V.L. Auslender and Iu.N. Pestov, Novosibirsk, for communicating experimental information prior to publication and for stimulating discussions.

A P P E N D I X

In the first part of this Appendix we shall discuss off the mass shell corrections due to treating the virtual exchanged pion in Fig. 1 as a real pion on the mass shell.

In the second part we shall briefly repeat the well-known argument which leads to the parametrization of $\Gamma(\omega)$ used in the text. In particular, we shall indicate possible reasons for the failure of Eq. (26). In the third part we shall show that two resonances in the same partial wave (two coherent elastic resonances) are conveniently described by a CDD pole and that this description in a natural way includes double poles discussed recently in connection with the A_2 meson.

A.1 Off the mass shell corrections

The expressions for the invariant mass distribution, given in Section 2, do not contain explicitly the corrections due to the virtuality of the exchanged pion. Similarly, the parametrization of the $\delta_1^1(\omega)$, given in Section 3, applies only to the elastic scattering for the equal mass particles. In this Appendix we wish to discuss these questions in more detail and to show that explicit inclusion of the off the mass shell effects cannot change the results obtained above.

In the following we shall denote the off the mass shell c.m.s. momentum, amplitude and cross-section as $q^{(o)}$, $a_\ell^{(o)}$, and $\sigma_\ell^{(o)}$, respectively. The on the mass shell notation is q , $a_\ell(\omega) = (\omega/q) \sin \delta_\ell \exp(i\delta_\ell)$, $\sigma_\ell(\omega)$. The relation between $q^{(o)}$ and the c.m.s. energy ω of the two pions is

$$q^{(o)} = \frac{1}{2\omega} \sqrt{[(\omega - m_\pi)^2 - t][(\omega + m_\pi)^2 - t]} \quad (\text{A.1})$$

where t is the momentum transfer in the reaction (1).

The relation between $a(\omega)$ and $a^{(o)}(\omega)$ has recently received much attention in connection with the current algebra calculations^{33,34)}. Even if the detailed features of this relation can be obtained only in a dynamical theory, one can get a reasonable form of this relation from general requirements. Thus, using the unitarity condition extrapolated in the mass of the exchanged pion and keeping only the 2π contribution in the intermediate state one obtains³⁴⁾

$$\text{Im } a_\ell^{(o)}(\omega) = q a_\ell^{(o)}(\omega, t)^* a_\ell(\omega), \quad (\text{A.2})$$

which shows that the phases of $a_\ell(\omega)$ and $a_\ell^{(o)}(\omega)$ are equal. Taking into account different threshold behaviour of $a_\ell^{(o)}$ and a_ℓ we can write

$$a_\ell^{(o)}(\omega, t) = \left(\frac{q^{(o)}}{q} \right)^\ell V_\ell(\omega, t) a_\ell(\omega) \quad (\text{A.3})$$

where $V_\ell(\omega, t)$ is finite at the threshold and real above it.

The simplest assumption $V \equiv 1$ corresponds to the Selleri ⁶⁾ correction, while an improved form

$$V_1^2(\omega, t) = \frac{1 + (qR)^2}{1 + (q^{(0)}R)^2} \quad (\text{A.4})$$

was proposed by Dürr and Pilkuhn ³⁵⁾ with $R \approx 1/3 m_\pi$.

More complicated prescriptions were used by Adler ³³⁾ and Katzin ³⁴⁾. The contribution of the peripheral production of the ρ to the effective mass distribution in the reaction (1) is proportional to the square of the $\pi\pi$ scattering amplitude. With the inclusion of the off the mass shell corrections we can therefore write this contribution

$$\frac{d\delta}{d\omega^2 dt} = K_\ell(t) f_\ell(\omega) \frac{q}{\omega} C(\omega, t) \delta_{\pi\pi}(\omega) \quad (\text{A.5})$$

where $K_\ell(t)$ and $f_\ell(\omega)$ do not include the off the mass shell corrections to $\pi\pi$ scattering,

$$C(\omega, t) = \left(\frac{q^{(0)}}{q} \right)^2 V^2(\omega, t), \quad (\text{A.6})$$

and $\delta_{\pi\pi}(\omega)$ is the on the mass shell $\pi\pi$ scattering. Off the mass shell corrections thus enter in a multiplicative way and this gives the reason why we have used a parametrization of $\Gamma(\omega)$ corresponding to the scattering of real pions.

As can be seen from Eqs. (A.6), (A.4) and (A.1) the function $C(\omega, t)$ is not factorizable into the product $h(\omega)g(t)$. This may introduce a dependence of the Q parameters on T (the maximal value of t in the sample). Integrating Eq. (A.5) over t as in Section 2 we obtain

$$\frac{d\delta}{d\omega^2} = f_e(\omega) \frac{q}{\omega} \sigma_{\pi\pi}(\omega) \int_{t_{\min}}^T K_e^2(t) C(\omega, t) dt$$

Using the mean value theorem we obtain

$$\frac{d\delta}{d\omega^2} = f_e(\omega) \frac{q}{\omega} \sigma_{\pi\pi}(\omega) C(\omega, \bar{t}(T)) \quad (A.7)$$

where, as indicated, $\bar{t}$ is a function of T . On the other hand, the phenomenological form of $K_e(t)$ is so much peaked at the lower end of the Chew-Low plot that $\bar{t}$ will practically be a constant. Then $C(\omega, \bar{t})$ can be included into the phenomenological function $f(\omega)$

$$f(\omega) = f_e(\omega) C(\omega, \bar{t})$$

and no T dependence will result.

A.2 $\Gamma(\omega)$ and dispersion relations

Let the invariant partial wave amplitude for two pions in a P wave, $a(\nu)$, be given ³⁶⁾ by Eq. (23) with $\ell = 1$. Since the phase shift δ is real in the region of elastic scattering it follows that

$$\operatorname{Re} \frac{1}{a(v)} = \frac{q^3}{\omega} \cot \delta, \quad (\text{A.8})$$

where $v = q^2$.

The dispersion relation for the inverse amplitude leads to ^{37,38)}

$$\begin{aligned} \operatorname{Re} \frac{1}{a(v)} = & \alpha + \beta v + \\ & + \operatorname{CM}(v) + L(v) + \operatorname{CDD}(v) + \operatorname{IN}(v), \end{aligned} \quad (\text{A.9})$$

where α, β are constants. Here $\operatorname{CM}(v)$ is given by ³⁶⁾

$$\operatorname{CM}(v) = \frac{1}{\pi} \sqrt{\frac{v^3}{v+1}} \ln \left(\sqrt{v+1} + \sqrt{v} \right),$$

$L(v)$ is the left-hand cut contribution

$$L(v) = \frac{v^2}{\pi} \int_{-\infty}^{-m_\pi^2} \frac{dv'}{v'^2} \frac{\operatorname{Im} \frac{1}{a(v')}}{v' - v}$$

and $\operatorname{IN}(v)$ is

$$\operatorname{IN}(v) = - \frac{v^2}{\pi} \mathcal{P} \int_{v_{\text{inel.}}}^{\infty} \frac{\sqrt{v'}}{\sqrt{v'+1}} [R(v') - 1] \frac{dv'}{v' - v}$$

$R(\nu)$ is the ratio of the total to elastic scattering cross-section in this partial wave. Finally $CDD(\nu)$, if present, is a sum of CDD poles¹⁵⁾

$$CDD(\nu) = \sum_i \frac{c_i}{\nu - \nu_i}$$

The formula (26) is evidently a good approximation near the resonance, provided that all terms in Eq. (A.9) are slowly varying functions. This is certainly true for the first two terms in Eq. (A.9) but possibly not for the last one. Therefore the parametrization used in Eq. (26) may break down if $\pi\pi$ scattering in the $T = J = 1$ partial wave becomes strongly inelastic close to the ρ resonance. Fortunately there are so far no indications of such a behaviour. Excluding inelasticity, the limitations of Eq. (26) will be due mainly to the third and fourth term in Eq. (26). The third term can be well approximated by a quadratic expression in ν over a broad region. The dynamics of the problem is hidden in the fourth term and any firm statements about it can hardly be made. Still, the quadratic expansion in ν should fit well at least in the limited region around a resonance.

A.3 CDD poles, double poles and nearby pairs of single poles

Let us now turn to CDD poles¹⁵⁾ and to the usual description of two nearby resonances and double poles in non-relativistic scattering. The similarity in results is not very surprising. If one wants to have two narrow resonances close together, the phase shift has to pass through π between the resonances because of the Wigner condition. This can be arranged in Eq. (A.9) only by a pole in the r.h.s.

In non-relativistic scattering a resonance is described by a pole in the S matrix element ³⁹⁾ in the complex k plane. Near the resonance

$$S(k) \equiv e^{2i\delta(k)} \approx \frac{k - \eta - ib}{k - \eta + ib} \quad (\text{A.10})$$

The variable k is related to the energy by $E = k^2$ and η and b are related to resonance position and width by $E_R = \eta^2$ and $\Gamma = 4\eta b$. From Eq. (A.10) one easily obtains

$$\cot \delta(E) = \frac{2}{\Gamma} (E_R - E) \quad (\text{A.11})$$

Eq. (A.11) is, in fact, a non-relativistic version of the Breit-Wigner formula, Eq. (27), used in the text.

The phase shift given by Eq. (A.11) has a wrong threshold behaviour. The reason is that the form for $S(k)$, given in Eq. (A.10), can be valid only in the region near the resonance. The simplest form consistent with analyticity and unitarity requirements on $S(k)$ is ⁴⁰⁾

$$S(k) = \frac{k - \eta - ib}{k - \eta + ib} \cdot \frac{k + \eta - ib}{k + \eta + ib} \quad (\text{A.12})$$

This leads to $\delta(k)$ in the form of Eq. (A.11) with $\Gamma \rightarrow \Gamma \frac{k}{\eta}$, which corresponds to an S wave threshold behaviour. In the following we shall discuss only the approximation (A.10) bearing in mind that the expressions corresponding to (A.12) can be obtained by the simple insertion $\Gamma \rightarrow \frac{k}{\eta} \Gamma$.

If there are two poles close together one has

$$S(E) = e^{2i\delta(E)} \approx \left(\frac{k - \eta_1 - ib_1}{k - \eta_1 + ib_1} \right) \left(\frac{k - \eta_2 - ib_2}{k - \eta_2 + ib_2} \right) = e^{2i(\delta_1 + \delta_2)} \quad (\text{A.13})$$

with

$$\begin{aligned} \cotg \delta_1 &= \frac{2}{\Gamma_1} (E_1 - E) , \\ \cotg \delta_2 &= \frac{2}{\Gamma_2} (E_2 - E) . \end{aligned} \quad (\text{A.14})$$

$\Gamma_1, \Gamma_2, E_1, E_2$ are related to η_1, η_2, b_1, b_2 by the same relations as those following Eq. (A.1). Using the addition theorem for $\cotg \delta = \cotg(\delta_1 + \delta_2)$ we immediately obtain from Eqs. (A.13) and (A.14)

$$\cotg \delta = A + B(E_c - E) + \frac{D}{E_c - E} \quad (\text{A.15})$$

with

$$A = \frac{2(\Gamma_2 - \Gamma_1)}{(\Gamma_1 + \Gamma_2)^2} (E_2 - E_1)$$

$$B = \frac{2}{\Gamma_1 + \Gamma_2}$$

$$E_c = \frac{\Gamma_2 E_1 + \Gamma_1 E_2}{\Gamma_1 + \Gamma_2}$$

$$D = - \left\{ \frac{2\Gamma_1\Gamma_2(E_2 - E_1)^2}{(\Gamma_1 + \Gamma_2)^3} + \frac{\Gamma_1\Gamma_2}{2(\Gamma_1 + \Gamma_2)} \right\}$$

A double pole corresponds to $E_1 = E_2 = E_c$ and $\Gamma_1 = \Gamma_2 = \Gamma$.

Then Eq. (A.15) simplifies to

$$\cot \delta(E) = \frac{1}{\Gamma} (E_c - E) - \frac{\Gamma}{4} \frac{1}{E_c - E} \quad (\text{A.16})$$

The difference between Eqs. (A.15) and Eq. (35) used in the text is only that in Eq. (A.16) the kinematical factor q^3/ω [or $(q^{2\ell+1})/\omega$ for the ℓ^{th} partial wave] is missing. Also, the variable E in Eq. (A.15) is replaced by ω^2 in Eq. (35). We therefore propose to use for two resonances in the same partial wave the relativistic version Eq. (35) rather than the non-relativistic one, Eq. (A.15). The scattering cross-section can easily be derived from Eq. (35)

$$\sigma = \frac{4\pi}{q^2} (2\ell + 1) \left| \frac{1}{\cot \delta - i} \right|^2 \quad (\text{A.17})$$

The differences between Eqs. (A.15) and (35) are of no importance for two very narrow resonances very close together. However, for an object extending over 100 MeV the shape may be significantly affected.

To make one "dynamical" remark on the plausibility of a double pole let us rewrite Eq. (A.15) in the form

$$\cot \delta = \frac{2}{\Gamma_1 + \Gamma_2} (E_0 - E) + \frac{D}{E_c - E}, \quad (\text{A.18})$$

where then

$$E_0 = \frac{2}{\Gamma_1 + \Gamma_2} (E_1 \Gamma_1 + E_2 \Gamma_2).$$

The last term in Eq. (A.18) then corresponds to $\text{CDD}(\sqrt{})$ in Eq. (A.9) and the first term to all other "dynamical" terms in Eq. (A.9). By "dynamical" we mean that all these terms are calculable, at least in principle, when the inelasticity and the discontinuity at the left-hand cut are given.

Equation (A.18) then demonstrates that the resonance at $E = E_0$ is produced by the "dynamical" terms in the absence of a CDD pole. The presence of a CDD pole, term 2 in Eq. (A.18), only forces the phase shift to go through π , and produces a second resonance at $3\pi/2$.

Comparing Eqs. (A.16) and (A.18) one sees that a true double pole would really appear as a miracle, since it is not only necessary to have the CDD pole sitting exactly at the dynamical resonance

$$E_c = E_0$$

but also the residue of the CDD pole must be precisely adjusted.

Let us, for completeness, write down explicitly the formula for πp scattering under the assumption of two resonances in the same partial wave. The parametrization can easily be obtained as the modification of Eq. (35) corresponding to an orbital momentum

$$l = 2$$

$$\frac{q^5}{k(\omega)} \cot \delta = \frac{(\omega_1^2 - \omega^2)(\omega_2^2 - \omega^2)}{\omega_d^2 - \omega^2} \frac{1}{P_n(\omega^2)} \quad (\text{A.19})$$

Here $k(\omega)$ is the kinematical factor taken to be ω in Eqs. (23)-(24) and 1 in Eq. (31). ω_1 is the mass of the first peak, ω_2 is the mass of the second peak and ω_d is the position of the dip (we called that before CDD zero position). $P_n(\omega^2)$ is a

polynomial in ω^2 corresponding to $(1+(Rq)^2)$ in the Q case and q is the c.m.s. momentum of outgoing particles related to ω by 10)

$$q = \frac{1}{2\omega} \sqrt{(\omega^2 - m_\pi^2 - m_\rho^2)^2 - 4m_\pi^2 m_\rho^2}.$$

The widths of the first and the second peak are given from (A.18) as [cf. Section 3, in particular the text before and after Eq. (27)]

$$\begin{aligned} \Gamma_1 &= \frac{q^5(\omega_1)(\omega_d^2 - \omega_1^2) P_n(\omega_1^2)}{k(\omega_1)(\omega_2^2 - \omega_1^2)} \\ \Gamma_2 &= \frac{q^5(\omega_2)(\omega_d^2 - \omega_2^2) P_n(\omega_2^2)}{k(\omega_2)(\omega_1^2 - \omega_2^2)} \end{aligned} \quad (\text{A.20})$$

In the actual fit it is advisable to take $P_n(\omega^2)$ to be constant and for $k(\omega)$ a simple factor ω or ω^2 . The polynomial $P_n(\omega^2)$ has anyway to be chosen so that it has no zeros in the A2 region, since these are accounted for explicitly in (A.19).

The cross-section for πQ scattering in the $\ell=2$ partial wave is then according to (A.17) and (A.19) given by

$$\sigma = \frac{20\pi (\omega_d^2 - \omega^2)^2 q^8 P_n^2(\omega)}{q^{10} (\omega_d^2 - \omega^2)^2 P_n^2(\omega) + k^2(\omega) (\omega_1^2 - \omega^2)^2 (\omega_2^2 - \omega^2)^2} \quad (\text{A.21})$$

This, however, is not directly applicable to the invariant mass distribution, since Eq. (A.21) corresponds only to the term $\delta_{\pi\pi}^{T=J=1}(\omega)$ used in Chapters 2 and 3 above and a more detailed description of the production mechanism is necessary.

The A_2 production is certainly a more complicated problem than that of the ρ meson, since

- i) the ρ meson decays exclusively to the $\pi\pi$ channel, but the A_2 meson decays to $\rho\pi$ in about 85 % ⁴¹⁾ of the cases; consequently the A_2 decay should rather be treated as a many channel problem;
- ii) in the $A_2 \rightarrow \rho\pi$ decay the ρ meson in the final state should be treated as an unstable particle while the Eq. (A.21) treats it as stable.

Table 1
Experimental Data

Sample name	Mass combination	Maximum $t [m_\pi^2]$	Events	Experiments
SACL 13 ⁻	$\pi^- \pi^0$	13	7666	2.77 GeV/c Saclay ¹⁶⁾ .
COMP 10 ⁻	$\pi^- \pi^0$	10	6205	Same as Fig. 14 in paper I ^{*)} .
TOT 10 ⁻	$\pi^- \pi^0$	10	12773	Above, and 2.77 GeV/c Saclay ¹⁶⁾ .
COMP 5 ⁻	$\pi^- \pi^0$	5	2038	Same as Fig. 11 in paper I ^{*)} .
TOT 5 ⁻	$\pi^- \pi^0$	5	5117	2.1 GeV/c Wisconsin ²³⁾ , 2.05-3.22 GeV/c Jacobs ⁵⁾ and 2.77 GeV/c Saclay ¹⁶⁾ .
COMP 10 ⁺	$\pi^+ \pi^0$	10	1831	Same as Fig. 13 in paper I ^{*)} .
COMP 10 ⁰	$\pi^+ \pi^-$	10	16109	Same as Fig. 8 in paper I ^{*)} .

*) Paper I, Tables 1 and 4, gives the breakdown into momenta and references.

Table 2
Best fit solutions

Solu- tion	Sample	$\Gamma(\omega)$	$f(\omega); \Delta_{\pi\pi}(\omega)$	$\chi^2 / \langle \chi^2 \rangle$	ω_Γ [MeV]	Γ_0 [MeV]	% backgr.	B_1	Other parameters
1	SACL 13 ⁻	Eq. (29)	1 ; 0	263/78	754	134	20	0	
2	"	"	ω^{-1} ; 0	100/78	766	152	20	0	
3	"	"	$(\omega\Gamma L)^{-1}$; 0	97/78	782	173	11	0	
4	"	Eq. (30)	1 ; Eq. (45)	218/77	772	202	-5	0	$R^2 = 6.3$
5	"	"	1 ; Eq. (46)	191/77	755	151	-72	0	$R^2 = -4.5$
6	"	"	1 ; C	79/76	767	154	23	0	$C = (0.1^{+2}_{-0.1})\%$, $R^2 = 2.3 \pm 0.5$
7	"	"	1 ; 0	79.1/77	767.0 ± 1.9	155 ± 5	23.0 ± 1.5	0	$R^2 = 2.3^{+0.5}_{-0.4}$
8	"	Eq. (36)	1 ; 0	83.7/77	765.1 ± 1.8	153 ± 5	27.7 ± 1.7	0	$\omega_0 = (1333^{+43}_{-37})$ MeV
9	"	Eq. (29)	$f_0 + \omega^{-1}$; 0	83.8/77	773.2	150	26	0	$f_0 = -0.70^*)$
10	"	"	$f_0 + \omega^{-2}$; 0	78.7/77	776.2 ± 2.4	161 ± 6	22.3 ± 1.6	0	$f_0 = -0.27^{+0.12}_{-0.10}^*)$
11	COMP 10 ⁻	Eq. (30)	1 ; 0	68.1/66	766.1 ± 2.1	153 ± 6	24.4 ± 1.8	0	$R^2 = 2.9 \pm 0.7$
12	"	Eq. (36)	1 ; 0	74.2/66	764.2 ± 2.1	156 ± 6	27.8 ± 1.9	0	$\omega_0 = (1274 \pm 42)$ MeV
13	TOT 10 ⁻	$(\omega_\Gamma/\omega) \times$ Eq. (30)	1 ; 0	69.1/68	767.9 ± 1.4	153.1 ± 3.9	20.7 ± 1.2	0	$R^2 = 0.36 \pm 0.09$
14	"	Eq. (30)	1 ; 0	70.7/66	766.6 ± 1.5	152.5 ± 3.9	21.4 ± 1.2	0	$R^2 = 2.7 \pm 0.5$
15	"	Eq. (33)	1 ; 0	70.1/68	766.5 ± 1.1	153.6 ± 3.9	17.9 ± 1.2	0	$R^2 = 384^{+200}_{-300}$
16	"	Eq. (36)	1 ; 0	82.3/66	764.7 ± 1.4	154.2 ± 4.0	24.7 ± 1.2	0	$\omega_0 = 1295 \pm 30$ MeV
17	"	Eq. (30)	1 ; 0	61.1/62	766.8 ± 1.5	148.2 ± 4.1	20.8 ± 1.3	0	$R^2 = 2.8 \pm 0.5$, res. 30 MeV
18	"	Eq. (36)	1 ; 0	69.6/62	765.0 ± 1.4	149.6 ± 4.2	24.3 ± 1.2	0	$\omega_0 = (1282 \pm 30)$ MeV, res. 30 MeV
19	"	Eq. (29)	$f_0 + \omega^{-2}$; 0	60.6/62	776.8 ± 1.9	155.4 ± 4.8	20.7 ± 1.5		$f_0 = -0.48 \pm 0.06^*)$, res. 30 MeV
20	"	Eq. (30)	$f_0 + f_2 \omega^{-2}$; 0	57.6/58	$764.3^{+1.6}_{-1.5}$	$147.3^{+3.9}_{-3.8}$	24.3	0.31	$f_0 = 0.82, f_2 = -0.17^*)$ $R^2 = 2.1^{+1.5}_{-0.8}$, $B_2 = 0$, res. 30 MeV
21	COMP 5 ⁻	"	1 ; 0	26.6/28	766.4 ± 3.2	162 ± 9	13.2 ± 2.4		$R^2 = 5.6^{+2.2}_{-1.5}$
22	TOT 5 ⁻	"	1 ; 0	65.1/78	764.5 ± 2.1	160 ± 6	12.8 ± 1.6	0	$R^2 = 1.9 \pm 0.4$
23	COMP 10 ⁺	"	1 ; 0	75.1/78	$757.3^{+4.3}_{-4.7}$	168 ± 14	32.5 ± 3.7	0	$R^2 = 5^{+3}_{-2}$
24	COMP 10 ⁰	"	1 ; 0	60.6/53	769.2 ± 1.5	163 ± 15	17.1 ± 2.0	0	$R^2 = 2.5 \pm 0.4$, inco- herent ω -meson with $\Gamma = 12$ MeV, $m_\omega = 783.4$ MeV. Res. 30 MeV
25	"	"	1 ; Eq. (43)	66.0/60	769.9 ± 1.4	164 ± 4	0	0	$R^2 = 2.1 \pm 0.3$, $\delta_0^0 = 44^\circ \pm 2^\circ$
26	"	"	1 ; Eq. (43)	61.5/58	768.7 ± 1.9	167^{+4}_{-3}	6.8 ± 4.0	0	$R^2 = 1.3 \pm 0.4$, $\delta_0^0 = (1.1^{+0.3}_{-0.2}) -$ $(0.8^{+0.4}_{-0.5})\omega^*)$
27	"	"	1 ; Eq. (43)	60.1/58	767.4 ± 1.7	150 ± 10	4 ± 9	0	$R^2 = 0.8 \pm 1.3$, $\omega_\Gamma = (760 \pm 115)$ MeV $\Gamma_\sigma = (460 \pm 110)$ MeV

*) ω is in units of GeV.

REFERENCES

- 1) M. Roos, Nucl. Phys. B2, 615 (1967).
- 2) J.D. Jackson, Nuovo Cimento 34, 1644 (1964).
- 3) V.L. Auslender, G.I. Budker, Iu.N. Pestov, V.A. Sidorov, A.N. Skrinsky and A.G. Khabakpashev, Phys. Letters 25B, 433 (1967), and private communications.
- 4) J.D. Jackson, J.T. Donohue, K. Gottfried, R. Keyser and B.E.Y. Svensson, Phys. Rev. 139, B428 (1965).
- 5) L.D. Jacobs, University of California, Lawrence Radiation Laboratory, Report UCRL-16877 (1966), unpublished.
- 6) F. Selleri, Phys. Letters 3, 76 (1962).
- 7) A.H. Rosenfeld, Oxford International Conference on Elementary Particles, 1965, Suppl. p.5.
- 8) D.H. Miller, L. Gutay, P.B. Johnson, F.J. Loeffler, R.L. McIlwain, R.J. Sprafka and R.B. Willmann, Phys. Rev. 153, 1423 (1967).
- 9) J.P. Baton and J. Reignier, Nuovo Cimento 36, 1149 (1965).
- 10) H. Pilkuhn, "The interactions of hadrons" (North-Holland, Amsterdam, 1967), Chapt. I.
- 11) The "non-relativistic" Breit-Wigner formula follows from Eq. (21) by the approximation $2\omega \approx \omega + \omega$. The r.h.s. of Eq. (21) then simplifies to $(\Gamma/2)^2 / [(\omega - \omega)^2 + (\Gamma/2)^2]$.
- 12) K. Brueckner, Phys. Rev. 86, 106 (1952).
- 13) We are leaving aside the question whether this is an appropriate definition from a fundamental point of view as discussed by G.F. Chew, University of California, Lawrence Radiation Laboratory, Report UCRL-16983 (1966), unpublished.
- 14) J.M. Blatt and V.F. Weisskopf, "Theoretical nuclear physics" (J. Wiley and Sons, New York, 1952), p. 312.

- 15) For a full discussion of CDD poles, see S.C. Frautschi "Regge poles and S matrix theory" (W.A. Benjamin, New York, 1964), which also contains references to the original literature. It is sufficient to consider our parametrization as a mere mathematical trick, allowing the phase shift to go through zero or π . We use "CDD poles" as a generic name for all zeros in the amplitude. Thus, in the N/D method, this includes zeros in N as well as poles in D.
- 16) J.P. Baton, G. Laurens and J. Reigner, Nucl. Phys. B3, 349 (1967);
J.P. Baton and G. Laurens (unpublished, submitted to Phys. Rev.)
and Phys. Letters 26B, 471 (1968);
- 17) G. Wolf, Phys. Letters 19, 328 (1965).
- 18) P.E. Schlein, Phys. Rev. Letters 19, 1052 (1967);
E. Malamud and P.E. Schlein, Phys. Rev. Letters 19, 1056 (1967).
- 19) A.B. Clegg, Phys. Rev. 163, 1664 (1967).
- 20) C. Lovelace, R.M. Heinz and A. Donnachie, Phys. Letters 22,
332 (1966);
C. Lovelace, Proc. of the Heidelberg International Conference
on Elementary Particles, 1967, North-Holland Publ. Co.,
(1968) p. 79.
- 21) P. Söding, Phys. Letters 19, 702 (1965).
- 22) M. Ross and L. Stodolsky, Phys. Rev. 149, 1172 (1966).
- 23) E. West, J.H. Boyd, A.R. Erwin and W.D. Walker, Phys. Rev. 149,
1089 (1966).
- 24) R.K. Logan, J. Beupre and L. Sertorio, Phys. Rev. Letters 18,
259 (1967);
See also L. Sertorio and M. Toller, Phys. Rev. Letters 19,
1146 (1967).
- 25) M.A. Jabiol, F.E. James and Nguyen Huu Khanh, Phys. Rev.
Letters 17, 1065 (1966).
- 26) R. Omnès, Nuovo Cimento 8, 316 (1958).
- 27) G.F. Chew, "S matrix theory of strong interactions"
(W.A. Benjamin, New York, 1962), p. 74.

- 28) W.R. Frazer and J.R. Fulco, Phys. Rev. Letters 2, 365 (1959)
and Phys. Rev. 117, 1609 (1960);
L.L. Vick, Nuovo Cimento 31, 643 (1964);
J.E. Bowcock, W.N. Cottingham and D. Lurié, Nuovo Cimento 16,
918 (1960).
- 29) P. Federbush, M.L. Goldberger and S.B. Treiman, Phys. Rev. 112,
642 (1958).
- 30) J. Harte, UCRL-preprint, August 1967 (and Phys. Rev. to be
published);
J. Stack, Phys. Rev. 164, 1904 (1967);
R. Hagedorn, CERN preprint TH. 853 (and refs. quoted there).
- 31) S. Bergia and J.A. Mignaco (preprint 1968), University of Bologna.
- 32) N.G. Antoniou and J.E. Bowcock, Phys. Rev. 159, 1257 (1967).
- 33) S. Adler, Phys. Rev. 140, B736 (1965).
- 34) J.C. Katzin, University of Maryland, preprint UMD-742 (1967).
- 35) H.P. Dürr and H. Pilkuhn, Nuovo Cimento 40A, 899 (1965),
- 36) G.F. Chew and S. Mandelstam, Phys. Rev. 119, 467 (1960).
- 37) J.W. Moffat, Phys. Rev. 121, 926 (1961).
- 38) K. Kang, Phys. Rev. 134, B1324 (1964).
- 39) M.L. Goldberger and K.M. Watson, Phys. Rev. 136, B1472 (1964);
J.S. Bell and C.J. Goebel, Phys. Rev. 138, B1198 (1965);
J.S. Bell, CERN preprints TH. 658 (1966) and TH. 784 (1967).
- 40) T. Regge, Lectures on High-Energy Physics, Herceg-Novi 1961,
Ed. B. Jaksic, Vol. I, p. 1;
A. Martin, *ibid*, Vol. I, p. 67.
- 41) A.H. Rosenfeld, N. Barash-Schmidt, A. Barbaroggaltieri, L.R. Price,
P. Söding, C.G. Wohl, M. Roos and W.J. Willis, Rev. Mod.
Phys. 40, 77 (1968).

FIGURE CAPTIONS

Fig. 1 : The OPE diagram.

Fig. 2 : Fits to the sample TOT 10^{-} (cf. Table 1). The error bars indicate the ± 1 standard deviation statistical error on the experimental histogram points. The curve represents solutions 17, 18 and 19 (cf. Table 2) which differ by less than the width of the line. The extension of the curve corresponds to the energy region used in the fits.

Fig. 3 : The χ^2 as a function of ω_r for solution 20 (cf. Table 2). The asymmetric ± 1 and ± 2 standard deviation errors are shown.

Fig. 4 : Fit to the sample COMP 10^0 (cf. Table 1). The error bars indicate the ± 1 standard deviation error on the experimental histogram points. The curve represents solution 24 (cf. Table 2). The extension of the curve corresponds to the energy region used in the fit.

DIA 33365