CM-P00058259

Ref.TH.1036-CERN

 $A_1 \rightarrow 3\pi$ DECAY

G.V. Dass ^{*)} and S. Papageorgiou
CERN - Geneva

A B S T R A C T

The coupling of the A_1 to a three-pion system is studied through the analysis of some simple two-body amplitudes where the A_1 appears as an either external or internal particle. The $A_1 \rightarrow 3\pi$ decay amplitudes to a pion and a P wave or S wave $\pi\pi$ system are related in the Veneziano model. The relevant ambiguities in the construction of dual amplitudes involving unnatural parity trajectories are discussed and some results on the strengths of the decays involving the two $\pi\pi$ systems are presented. Further experimental information of the four-pion decay of the f^0 would help to determine the strength of the A_1 decay involving the spinless dipion system.

^{*)} On attachment from Atomic Energy Research Establishment, Harwell and from (also, address after 16 May 1969) Rutherford Laboratory, Chilton, Didcot, Berks.

1. INTRODUCTION

In spite of doubts about its existence, the A_1 ($J^{PC} = 1^{++}$) meson has been discussed from many points of view. It can decay into a dipion system (of spin ℓ) and a pion with relative angular momenta $\ell \pm 1$ between the two. We shall be mainly concerned with $A_1 \rightarrow \epsilon\pi$ ($\ell=0$) and $\rho\pi$ ($\ell=1$) decays. Different theoretical models give different results for the characteristics of this decay. To quote a few: hard pion ¹⁾ or effective Lagrangian calculations predict a predominantly S wave $A_1 \rightarrow \rho\pi$ decay and a substantial $A_1 \rightarrow \epsilon\pi$ rate. Quark model calculations ²⁾ of pion emission by quarks, neglecting recoil corrections, predict a predominantly transverse ($1 \rightarrow 1$ helicity amplitude) $A_1 \rightarrow \rho\pi$ decay. Ademollo et al. ³⁾ predict a dominantly S wave $A_1 \rightarrow \rho\pi$ decay; other workers ⁴⁾ using current algebra and superconvergence ideas predict a predominantly longitudinal ($0 \rightarrow 0$ helicity amplitude) $A_1 \rightarrow \rho\pi$ decay; this implies there is more D wave than S wave contribution to this decay. The sum rules of Gilman and Harari ⁴⁾ gave a width $\Gamma(A_1 \rightarrow \pi\epsilon) \simeq 50$ MeV. A recent experiment ⁵⁾ which did not consider the possibility of a P wave decay ($A_1 \rightarrow \pi\epsilon$) indicates that the data are better fitted by a pure D wave than a pure S wave $A_1 \rightarrow \rho\pi$ decay.

There has been some interest ^{6), 7)} from the point of view of the Veneziano model ⁸⁾ also. De Alwis and Nutbrown and Brooker and Kosterlitz ⁶⁾ considered the reaction $\pi\pi \rightarrow \pi A_\mu$ where A_μ is the axial vector current. Their essential conclusion is that it is difficult to determine the relative strength of the S and D wave contributions in $A_1 \rightarrow \rho\pi$ decay without making further dynamical assumptions on some off-mass shell amplitudes. Fayyazuddin and Riazuddin ⁷⁾ considered the reaction $\pi\pi \rightarrow \pi A_1$ and by going to the ρ pole they tried to determine the ratio G_S/G_D of the S and D wave contributions defined by the Hamiltonian

$$\mathcal{H}(A_1 \rightarrow \rho\pi) = -G_S \epsilon^\rho \cdot \epsilon^{A_1} + 2G_D \epsilon^{A_1} \cdot p_\rho \epsilon^\rho \cdot p_A \quad (1)$$

where ϵ^{ρ, A_1} and p_{ρ, A_1} are the polarization vectors and four-momenta of the ρ and the A_1 . The ratio G_S/G_D can be expressed ⁷⁾ in terms of the anomalous magnetic moment δ of the A_1 meson as

$$\frac{G_S}{G_D} = \left(\frac{2+\delta}{\delta} \right) m^2 \quad (2)$$

where m is the ρ meson mass. The value $\delta = -2$ corresponds ⁷⁾ to the algebraic realization of chiral symmetry and the value $\delta \approx -1$ is "attractive" because it would give, in chiral symmetry, the right experimental ρ width. Actually, if one includes the possible decay $A_1 \rightarrow \pi \epsilon$ in a Veneziano amplitude for the process $\pi\pi \rightarrow \pi A_1$ (see next Section), where ϵ is the first daughter of the ρ -f trajectory at the ρ position, one gets a relation like

$$\left[G_S + (m^2 - 2\mu^2) G_D \right]^2 = (m^2 - 4\mu^2) g_{A_1 \epsilon \pi}^2 \quad (3)$$

where μ is the pion mass and $g_{A_1 \epsilon \pi}^2$ (defined in the Appendix) determines the $\epsilon\pi$ decay probability. The value $\delta = -1$ (in the approximation $\mu^2/m^2 \sim 0$) corresponds to a vanishing $\Gamma(A_1 \rightarrow \epsilon\pi)$. So, it is interesting to determine $g_{A_1 \epsilon \pi}^2$ in order to be able to choose between the various possibilities. For that purpose, it is natural to consider amplitudes (like, for example, $\pi\epsilon \rightarrow \pi\epsilon$, $\pi\epsilon \rightarrow \pi\rho$ and $A_1\epsilon \rightarrow \pi\epsilon$) where the ϵ occurs as an external particle rather than an internal particle because one does not, a priori, expect factorization to hold at the first daughter level in the usual Veneziano model. As we shall see in Section 3, the amplitudes involving unnatural parity trajectories are subject to some ambiguities. In order not to possibly multiply these ambiguities and add further spin complications, we choose to consider the simplest case: a $\pi\epsilon \rightarrow \pi\epsilon$ Veneziano amplitude and try to determine $g_{A_1 \epsilon \pi}^2$ by going to the A_1 pole.

Section 2 is devoted to a brief review of the Veneziano amplitude ⁷⁾ for the reaction $\pi\pi \rightarrow \pi A_1$ and a derivation of Eq. (3). In Section 3, $\pi\pi$ and $\pi\pi$ scattering and the ambiguities attending the relevant Veneziano amplitudes are discussed. The main conclusions are summarized in Section 4. Finally, some formulae are collected in the Appendix.

2. THE REACTION $\pi\pi \rightarrow \pi A_1$

The $\pi^+(p) \pi^-(q) \rightarrow \pi^-(q') A_1^+(k)$ Veneziano amplitude was considered in Ref. 7). For the sake of completeness, we review it here, mentioning also the structure of the amplitude ⁷⁾ A and all the three Adler conditions. Defining s, u and t as $s = (p+q)^2$, $u = (k-q)^2$ and $t = (q-q')^2$, one writes the amplitude as

(4)

$$T(s,t) = [A(s,t) \cdot (q-q')_\mu + B(s,t) \cdot (q+q')_\mu] \cdot \epsilon_\mu^A$$

where considerations of asymptotic behaviour of the invariant amplitudes A and B and crossing symmetry lead to one of the forms

$$A(s,t) = (\gamma_1 - \gamma_2 t + 2\gamma_2 s) \frac{\Gamma(1-\alpha(s))\Gamma(1-\alpha(t))}{\Gamma(2-\alpha(s)-\alpha(t))} \quad (5)$$

and

$$B(s,t) = (\gamma_1 + \gamma_2 t) \frac{\Gamma(1-\alpha(s))\Gamma(1-\alpha(t))}{\Gamma(2-\alpha(s)-\alpha(t))} \quad (6)$$

where α is the degenerate ρ, f trajectory and γ_1, γ_2 are constants. The Adler consistency condition that the full amplitude T vanishes when one of the pions has zero four-momentum gives the three requirements:

- a) $q_{\mu}^1 \rightarrow 0$: $(A+B)=0$ at $s=M^2$, $t=\mu^2$
 b) $q_{\mu} \rightarrow 0$: $(A-B)=0$ at $s=t=\mu^2$ and
 c) $p_{\mu} \rightarrow 0$: $B=0$ at $s=\mu^2$, $t=M^2$

where M is the A_1 mass. The condition c) was satisfied in Ref. 7) by requiring $\Gamma(2-\alpha(s)-\alpha(t))$ to have a pole at $s=\mu^2$, $t=M^2$. Choosing $\alpha(\mu^2)=\frac{1}{2}$ and $\alpha(M^2)=\frac{3}{2}$ with the trajectory slope $1/(2(m^2-\mu^2))$ gives the Weinberg mass formula for the A_1 . Condition b) is seen to be trivially satisfied because $(A-B)=0$ at $s=t$. Condition a) is also satisfied if condition c) is satisfied because A and B are separately zero for $s=M^2$, $t=\mu^2$. The structure in Eqs. (5) and (6), therefore satisfied all the three Adler conditions for arbitrary γ_1 and γ_2 . To determine these constants, one goes to $\alpha(s)=1$ and $\alpha(t)=1$ and compares the residua of (5) and (6) with those of the ρ and ϵ contributions. The details are given in Ref. 7) for the amplitude B . It turns out that doing the same thing for the A amplitude, one gets identically the same results as by considering the B amplitude. The net result is the relation

$$-g_{\rho\pi\pi} [G_S + (m^2 - 2\mu^2) G_D] = g_{\epsilon\pi\pi} g_{A_1\epsilon\pi} \quad (7)$$

from which, using the relation $\Gamma_{\epsilon}/\Gamma_{\rho} = 9/2$, one derives Eq. (3). If one knows $g_{A_1\epsilon\pi}^2$, Eq. (3) gives a relation between G_S and G_D . Another relation between G_S and G_D is obtained from a knowledge of the width $\Gamma(A_1 \rightarrow \rho\pi)$ in terms of G_S and G_D ; one can tentatively take $^{*)}$

$$\Gamma(A_1 \rightarrow \rho\pi) = \Gamma_{total}(A_1 \rightarrow 3\pi) - \Gamma(A_1 \rightarrow \epsilon\pi)$$

$^{*)}$ This relation, strictly speaking, is only approximate, in view of the rather large error on Γ_{total} and the large overlap due to the ρ and ϵ widths.

(neglecting overlap), the right-hand side being known if $g_{A_1 \pi \pi}^2$ is known. These two relations determine G_S and G_D apart from ambiguities arising from the large experimental error on Γ_{total} , the sign of $g_{A_1 \pi \pi}$ and from the solution of a quadratic equation. Lacking these ambiguities, this procedure would determine the complete Dalitz plot and angular distribution of $A_1 \rightarrow 3\pi$ decay. We now turn to $\pi \pi$ scattering which we use to determine $g_{A_1 \pi \pi}^2$.

3. $\pi \pi$ SCATTERING

The trajectories π and A_1 occur in the s and u channels and the f trajectory in the t channel. Symmetry between s and u channels kills odd partial waves (for example, the $J^P = 1^-$ pole at the ρ position) in the t channel. With the usual slope $b = 1/(2(m^2 - \mu^2))$, the π and the A_1 are degenerate in trajectories. Consideration of the $\pi \pi$ scattering alone does not rule out complete exchange degeneracy (in trajectory and in residue functions) between π and A_1 ; a consideration¹⁰⁾ of the $\pi \rho$ scattering, however, does. We shall consider the two cases separately, though we shall always assume πA_1 degeneracy in trajectories.

a. Non-degenerate case

The π and the A_1 trajectories are degenerate here, but the corresponding residues are not correlated. A simple Veneziano amplitude having, for each term, the correct asymptotic behaviour and pole structure in all channels is

$$\left[\left\{ (E_1 \alpha_\pi(s) + E_2) X(\alpha_\pi(s), \alpha(t)) + (s \rightarrow u) \right\} + F Y(\alpha_{A_1}(s), \alpha_{A_1}(u)) \right. \\ \left. + G Z(\alpha_\pi(s), \alpha_\pi(u)) + B \left\{ Y(\alpha(t), \alpha_{A_1}(s)) + (s \rightarrow u) \right\} \right. \\ \left. + \left\{ (H_1 \alpha_\pi(s) + H_2) X(\alpha_\pi(s), \alpha_{A_1}(u)) + (s \leftrightarrow u) \right\} \right]$$

where E_1 , E_2 , F , G , B , H_1 and H_2 are constants and
 $X(a,b) = \Gamma(-a) \Gamma(1-b) / \Gamma(1-a-b)$, $Y(a,b) = \Gamma(1-a) \Gamma(1-b) / \Gamma(1-a-b)$
and $Z(a,b) = \Gamma(-a) \Gamma(-b) / \Gamma(-a-b)$.

Before we turn to a determination of the various constants, we remark that the $\pi\pi \rightarrow \pi\pi$ Veneziano amplitude suffers from some ambiguities. In the $\pi\pi \rightarrow \pi\pi$ case ⁹⁾, for example, the possible ambiguities arise mainly from the addition of satellite terms which do not have the maximum possible asymptotic behaviour. Apart from these, the $\pi\pi \rightarrow \pi\pi$ case has another difficulty which arises mainly because on the natural parity trajectories ($\rho, f; \omega, A_2; \phi, f'$, etc.), one does not want a pole at $\alpha = 0$, but one does need the physical pole $\alpha_\pi(s) = 0$. This asymmetry between natural and unnatural parity trajectories (particularly the ones requiring pseudoscalar meson poles) is seen in the terms involving (E_1 and E_2) and (H_1 and H_2). Taking, for example, the (s,t) term, one sees that as long as one wants to write the amplitude in terms of Γ functions, one cannot have just one term having the correct maximum possible asymptotic behaviour and also the pion pole. The E_2 (E_1) term does (not) have the pion pole and the E_1 (E_2) term does (not) have the full asymptotic behaviour. The E_2 term does not contribute at the parent level in the t channel and so far as the t channel is concerned, it is only a satellite term; in the s channel, however, it is the only (s,t) term having the pion pole, and, therefore, is very important. This difficulty arises only when one wants the $\alpha = 0$ pole in only one of the relevant channels; in the terms involving F , G and B , this is not the case (as in $\pi\pi$ scattering).

One would be certainly "more general" by keeping both the E_1 and E_2 terms, but it is not obvious whether the full asymptotic behaviour should be imposed on each term separately or only on the complete amplitude. Moreover, information in the s channel about E_1 would arise from $J \geq 2$ poles and practically nothing is experimentally known about the quantitative strength of these poles. An extreme attitude would be to put $E_1 = 0$ because this term does not have the correct pole structure and $E_2 = 0$ because this term does not

have the full asymptotic behaviour. All the above remarks apply equally well to the $(H_1 \text{ and } H_2)$ term and doing the same thing for $H_1 = H_2 = 0$ would leave the G term as the only contribution from the π trajectory. It is clear from the above discussion that even the requirements of "simplicity" leave this Veneziano amplitude non-unique ^{*)}. We shall keep all the constants E_1 , E_2 , H_1 and H_2 non-zero and later consider special cases.

The Adler consistency condition that the full amplitude vanishes when the four-momentum of either of the pions goes to zero is satisfied for the above amplitude if one remembers ⁹⁾

$\alpha_p(p^2) = \frac{1}{2}$ and chooses $\alpha(s=m^2) = \alpha(u=m^2) = \frac{1}{2}$. Both of the Adler points are at $s=u=m^2$ and $t=p^2$ and all the Γ functions in the denominators in the amplitude (8) develop poles at this point.

In order to determine the various constants, we impose the requirement that there be no poles at $\alpha_\pi(s) = \text{odd integers}$ and $\alpha_{A_1}(s) = \text{even integers}$. This gives us the three conditions

$$\pi : \quad E_1 = H_1 \quad \text{and} \quad E_2 = G + H_2$$

$$A_1 : \quad B + F = H_1$$

Using these conditions in a comparison of the residues of the poles at $\alpha_\pi(s) = 0$ (pion pole), $\alpha_{A_1}(s) = 1$ (the $J=1$ part is the A_1 pole) and $\alpha_t = 2$ (the $J=2$ part is the f^0 pole), one gets (see the Appendix for definitions of coupling constants, etc.) respectively

$$g_{\pi\pi\pi}^2 = \frac{2E_2}{b} \quad (9)$$

^{*)} These remarks easily generalize to other similar amplitudes requiring $J=0$ poles.

$$g_{A_1 \epsilon \pi}^2 = -4B \quad (10)$$

$$g_{f \pi \pi} g_{f \epsilon \epsilon} = 8b(E_1 - B) \quad (11)$$

where $\alpha(t) = A + bt$ and $\alpha_\pi(s) = \alpha_{A_1}(s) = a + bs$. The aim to determine $g_{A_1 \epsilon \pi}^2$, therefore, cannot be achieved unless one uses some further information to determine E_1 and then B from Eq. (11). In the most general case, therefore, one is not able to determine $g_{A_1 \epsilon \pi}^2$. The following determinations arise only in special cases and serve to indicate the various possible relations between different (in some cases apparently unrelated) experimental numbers. If, however, one takes the attitude that only the complete amplitude (8) and not each of its individual terms need have the full asymptotic behaviour, one can use $E_1 = 0$ (therefore, $H_1 = 0$) and still have the full asymptotic behaviour $s^{\alpha(t)}$ for the whole amplitude. Then, $g_{A_1 \epsilon \pi}^2$ can be determined from a knowledge of $g_{f \pi \pi} g_{f \epsilon \epsilon}$ from Eq. (11). While $g_{f \pi \pi}$ is determined by the width $\Gamma(f \rightarrow \pi \pi)$, one still has to use as input some dynamical information for $g_{f \epsilon \epsilon}$. One can regard $f \rightarrow 4\pi^0$ decay to arise dominantly through the chain $f \rightarrow \epsilon \epsilon \rightarrow (\pi^0 \pi^0)(\pi^0 \pi^0)$ and use the width $\Gamma(f \rightarrow 4\pi^0)$ to get some estimate of $g_{f \epsilon \epsilon}$. Lacking an experimental knowledge of $\Gamma(f \rightarrow 4\pi^0)$ which does not get contributions from intermediate ρ^0 's, one might use $\Gamma(f \rightarrow 2\pi^+ 2\pi^-)$ and subtract out $\Gamma(f \rightarrow \rho \rho \rightarrow 2\pi^+ 2\pi^-)$ to get an upper limit for $g_{f \epsilon \epsilon}$. Unfortunately, $\Gamma(f \rightarrow \rho^0 \rho^0 \rightarrow 2\pi^+ 2\pi^-)$ is also not precisely known at the moment, though the data are consistent ¹¹⁾ with the assumption that the $\rho \rho$ intermediate state dominates the known branching ratio $\Gamma(f \rightarrow 2\pi^+ 2\pi^-) / \Gamma(f \rightarrow \pi^+ \pi^-) \approx 10\%$. We use

$$\frac{\Gamma(f \rightarrow \epsilon \epsilon \rightarrow 2\pi^+ 2\pi^-)}{\Gamma(f \rightarrow 2\pi)} = \chi \%$$

as a crude number for the contribution of the $\epsilon \epsilon$ state; perhaps it is reasonable to take $\chi = 1$ or $\frac{1}{2}$ say. This gives (see the Appendix for details) $g_{f \epsilon \epsilon} \simeq 130 \sqrt{\chi} \text{ GeV}^{-1}$. Taking the experimental

width of f decay as dominantly due to the 2π mode to get $g_{f\pi\pi}$, one determines B using the above value of $g_{f\epsilon\epsilon}$. One gets (for $\chi = \frac{1}{2}$), $\Gamma(A_1 \rightarrow 3\pi) \approx 90$ MeV, to be compared with the experimental total width $\Gamma(A_1 \rightarrow 3\pi) = 80 \pm 35$ MeV. Without a precise knowledge of χ , it is not very meaningful to use this $g_{A_1\epsilon\pi}^2$ in Eq. (3) and try to calculate G_S and G_D , etc.; it only suggests that $g_{A_1\epsilon\pi}^2$ could be substantial and should not be neglected in Eq. (3) above. It is on that ground that duality arguments, involving the A_1 can be used in $\epsilon\pi$ scattering. An interesting relation (for $E_1=0$) between $\Gamma(A_1 \rightarrow \epsilon\pi)$ and $\Gamma(f \rightarrow \epsilon\epsilon \rightarrow 2\pi^+ 2\pi^-)$ follows from Eqs. (10) and (11) as

$$g_{A_1\epsilon\pi}^2 = g_{f\pi\pi} g_{f\epsilon\epsilon} / 2b \quad (12)$$

It would be nice to have experimental information on $\Gamma(f \rightarrow 4\pi^0)$ or on the difference $\Gamma(f \rightarrow 2\pi^+ 2\pi^-) - \Gamma(f \rightarrow \rho\rho \rightarrow 2\pi^+ 2\pi^-)$ to get better limits on $g_{f\epsilon\epsilon}$ and then some number for $g_{A_1\epsilon\pi}^2$.

One might be tempted to exploit the presence of the satellite (so far as t channel is concerned) term E_2 to require factorization at the first daughter level, especially at the ϵ position ($\alpha_t=1$); this actually gives

$$g_{\epsilon\pi\pi} g_{\epsilon\epsilon\epsilon} = \frac{2E_2 + b m^2 (E_1 - B)}{b} \quad (13)$$

Using Eqs. (9), (11) and (13), one gets the condition

$$\frac{m^2}{8b} g_{f\pi\pi} g_{f\epsilon\epsilon} = g_{\epsilon\pi\pi} (g_{\epsilon\epsilon\epsilon} - g_{\epsilon\pi\pi}) \quad (14)$$

Furthermore, one can use a Veneziano amplitude of the form

$$[\gamma(\alpha(s), \alpha(t)) + \gamma(\alpha(u), \alpha(t)) + \gamma(\alpha(s), \alpha(u))]$$

for $\epsilon\epsilon$ scattering and going to the poles $\alpha(t)=2$ and 1 , get

a relation between $g_{f\epsilon\epsilon}^2$ and $g_{\epsilon\epsilon\epsilon}^2$ as

$$g_{\epsilon\epsilon\epsilon}^2 = g_{f\epsilon\epsilon}^2 \frac{2A + 3bm^2}{8b^2} \simeq 0.4 g_{f\epsilon\epsilon}^2 \quad (15)$$

(The unit of mass is 1 GeV.) This relation, combined with Eq. (14) and the positivity of $g_{f\epsilon\epsilon} g_{f\pi\pi}$ actually determines $g_{f\epsilon\epsilon} \simeq 15.6 \text{ GeV}^{-1}$ and thence, also $g_{\epsilon\epsilon\epsilon}$. This value of $g_{f\epsilon\epsilon}$ corresponds to a very small value of $x \simeq 1.4 \times 10^{-2}$. Imposing factorization at the ϵ pole by exploiting the presence of the (t channel) satellite term, therefore, one predicts the branching ratio for the mode $f \rightarrow \epsilon\epsilon \rightarrow 2\pi^+ 2\pi^-$ as $\simeq 0.015\%$; this is consistent with the experimental result that the $\rho\rho$ intermediate state dominates the decay mode $f \rightarrow 2\pi^+ 2\pi^-$.

Having thus determined $g_{f\epsilon\epsilon}$, one uses Eqs. (10) and (11) with $E_1 = 0$ to get $g_{A_1\epsilon\pi}^2$ and thence, $\Gamma(A_1 \rightarrow \epsilon\pi) \simeq 24 \text{ MeV}$. As a closer result one gets $\Gamma(A_1 \rightarrow 3\pi) \simeq 16.5 \text{ MeV}$, using the width $\Gamma(\epsilon \rightarrow \pi\pi) \simeq 540 \text{ MeV}$ from $\pi\pi$ scattering ⁹⁾ in order to carry out the final phase-space integrations.

Using this width $\Gamma(A_1 \rightarrow \epsilon\pi)$, one gets a relation between G_S and G_D from Eq. (3) and then writing $\Gamma(A_1 \rightarrow \rho\pi) = \Gamma_{\text{total}}(A_1 \rightarrow 3\pi) - \Gamma(A_1 \rightarrow \epsilon\pi)$ (neglecting overlap) in terms of G_S and G_D (see the Appendix for details), one gets

$$\frac{G_S}{G_D} = (-0.05 \pm 0.10) \quad \text{or} \quad (0.23 \pm 0.15) \quad (16)$$

The assigned errors attempt to account for the uncertainty in the experimental $\Gamma_{\text{total}}(A_1)$ and the two values correspond to the two solutions of the quadratic equation in G_S (or G_D) from the width $\Gamma(A_1 \rightarrow \rho\pi)$ equation. These values tend to agree with the experimental result ⁵⁾ that the D wave amplitude is more important.

Equation (2) in these units reads

$$\frac{G_S}{G_D} \simeq \frac{2+\delta}{\delta} (0.58)$$

so that the solution $\delta \leq -2$ is possible, but not any positive value as well as $\delta = -1$.

It is fair to remark that the results in Eq. (16) are not completely unique in view of the above mentioned ambiguities in the construction of the $\pi \epsilon \rightarrow \pi \epsilon$ Veneziano amplitude; these numbers arise in the case when : 1) $E_1 = 0$ and 2) the presence of the E_2 term is exploited to impose factorization at the ϵ pole in the t channel. It is encouraging to recall, however, that the predicted value of α for this case ($E_1 = 0$) is reasonable.

b. Degenerate case

Here, the π and A_1 are completely degenerate with each other (both in residue and trajectory). One can write the full $\pi \epsilon \rightarrow \pi \epsilon$ amplitude as

$$[(E'_1 \alpha(s) + E'_2) X(\alpha(s), \alpha(t)) + (s \rightarrow u)] + G' Z(\alpha(s), \alpha(u)) \quad (17)$$

The equations corresponding to (9), (10), (11) and (13) now are

$$E'_2 + G' = b g_{\epsilon \pi \pi}^2 \quad (18)$$

$$2(E'_2 + E'_1 - G') = g_{A_1, \epsilon \pi}^2 \quad (19)$$

$$E'_1 = \frac{g_{f \pi \pi} g_{f \epsilon \epsilon}}{8b} \quad (20)$$

$$2E'_2 + b m^2 E'_1 = b g_{\epsilon \pi \pi} g_{\epsilon \epsilon \epsilon} \quad (21)$$

It is important to have $E'_1 \neq 0$ in the degenerate case because, otherwise, there are no contributions at the parent level in the t channel. In general, however, one needs to know E'_1 ,

E'_2 and G' to determine $g_{A_1 \epsilon \pi}^2$; the input equations (18) and (20) are not sufficient to determine all these constants. However, because of the presence of the E'_2 term which is a satellite for the t channel, one may again use factorization at $\alpha_t = 1$ (ϵ pole) to get Eq. (21). One can now determine $g_{A_1 \epsilon \pi}^2$ using Eqs. (18) to (21) and the relation (15) between $g_{\epsilon \epsilon \epsilon}^2$ and $g_{\epsilon \epsilon \epsilon}^2$ from the $\epsilon \epsilon$ Veneziano model. One again gets a relation

$$g_{A_1 \epsilon \pi}^2 \cong 900 \sqrt{x} - 26$$

between the branching ratio x and $g_{A_1 \epsilon \pi}^2$. Using $x = \frac{1}{2}$, one finds a $g_{A_1 \epsilon \pi}^2$ that gives $\Gamma(A_1 \rightarrow \epsilon \pi) \simeq 65$ MeV which, again only indicates that $g_{A_1 \epsilon \pi}^2$ could be very big. A precise experimental knowledge of x would help to know $g_{A_1 \epsilon \pi}^2$. If one tentatively uses the value of $x \approx 0.015$ determined in the non-degenerate case above, one gets $\Gamma(A_1 \rightarrow \epsilon \pi) \approx 6$ MeV and therefore, $\Gamma(A_1 \rightarrow \rho \pi) \approx \Gamma_{\text{total}}$; in that case, Eq. (3) gives $G_S/G_D \simeq -m^2$ and $\delta \simeq -1$. We note that the degenerate case is really a special case in which the (πH) trajectory is exchange degenerate (in residue also) with the $(A_1 A_0)$ trajectory where H and A_0 are $I=0$ trajectories with odd and even signature respectively and having their lowest spin particles for $J^{PC} = 1^{+-}$ and 2^{--} . The degeneracy of the π and A_1 trajectories can be inferred from duality in KK^* charge exchange scattering.

c. Some parenthetical remarks

- 1) The full amplitude (8) of the non-degenerate case is still not determined because the constant G (or H_2) is still unknown. That would require, for example, a knowledge of the width of the $J^P = 0^-$ particle at the A_1 position, the relevant residue being $-\frac{1}{b}(H - \frac{m_B^2}{2M^2})$. In the degenerate case, one can determine this width which turns out to be about twenty times larger than the A_1 width for $x=1$ and ≈ 13 MeV for $x = 1.4 \times 10^{-2}$.

2) As expected, we do not get factorization for the $J^P=0^+$ particle at $\alpha_t=2$. In the ND case the corresponding residue further turns out to be negative if one uses the value of $g_{f\pi\pi}$ (corresponding to $x=1.4 \times 10^{-2}$) obtained above.

3) We have used the experimental ¹²⁾ f width. Our results, for example, Eq. (16), remain practically unchanged even if we used the width (≈ 100 MeV) given by the $\pi\pi \rightarrow \pi\pi$ Veneziano model ¹⁰⁾ for $\Gamma_{\text{total}}(f \rightarrow 2\pi) = 120$ MeV.

4. CONCLUSIONS

In order to know all the characteristics of $A_1 \rightarrow 3\pi$ decay, one needs to know G_S , G_D and $g_{A_1\pi\pi}$ separately. By considering the processes $\pi\pi \rightarrow \pi A_1$ and $\pi\pi \rightarrow \pi\pi$ where A_1 occurs as an external and internal particle respectively, one could determine G_S in terms of G_D knowing $g_{A_1\pi\pi}^2$ from $\pi\pi \rightarrow \pi\pi$ scattering. Combining this with a knowledge of the A_1 width, one can determine G_S/G_D . Unfortunately, even "simplicity" requirements are not enough to remove the ambiguities of the Veneziano amplitude for $\pi\pi$ scattering; these ambiguities have been discussed in detail in Section 3.a. With the most ambitious Veneziano amplitude (8), one cannot, in general, determine $g_{A_1\pi\pi}^2$ at present. By considering the case $E_1=H_1=0$ when only the complete amplitude (8) and not each separate term has the full asymptotic behaviour, we could determine $g_{A_1\pi\pi}^2$ only if we know the width $\Gamma(f \rightarrow 4\pi^0)$ or the contribution $\Gamma(f \rightarrow \pi\pi \rightarrow 2\pi^+2\pi^-)$ of the $\pi\pi$ intermediate state to $(f \rightarrow 2\pi^+2\pi^-)$ decay. Use of some reasonable numbers for this contribution indicates that $g_{A_1\pi\pi}^2$ could be large and should not be neglected in Eq. (3). An interesting outcome for the case $E_1=0$, is the relation (12) between the widths $\Gamma(A_1 \rightarrow \pi\pi)$ and $\Gamma(f \rightarrow \pi\pi \rightarrow 4\pi)$. It would be good to have experimental results on $(f \rightarrow 4\pi^0)$ or on $\Gamma(f \rightarrow \pi\pi \rightarrow 2\pi^+2\pi^-)$. Imposing the further requirement of factorization at the π pole in view of the presence of the satellite term E_2 , one could determine the branching ratio

$(f \rightarrow \epsilon \epsilon \rightarrow 2 \pi^+ 2 \pi^-)/(f \rightarrow \text{all}) \approx 0.015\%$ which is consistent with the experimental result that the decay $f \rightarrow 2 \pi^+ 2 \pi^-$ occurs mainly through the $\rho\rho$ intermediate state. This leads to the values G_S/G_D in Eq. (16).

In the degenerate case some of the above ambiguities are not present and one can write a simple Veneziano amplitude. However, we could determine $g_{A_1 \epsilon \pi}^2$ only by imposing factorization at the ϵ pole by exploiting the presence of the satellite term E_2' . Doing this, a knowledge of the contribution $(f \rightarrow \epsilon \epsilon \rightarrow 2 \pi^+ 2 \pi^-)$ to the decay $(f \rightarrow 2 \pi^+ 2 \pi^-)$ or of $\Gamma(f \rightarrow 4 \pi^0)$ would give $g_{A_1 \epsilon \pi}^2$. Taking this contribution to be very small ¹¹⁾ ($x = \frac{1}{2}$), one concludes, as in the non-degenerate case, that $\Gamma(A_1 \rightarrow 3 \pi)$ could be an appreciable part (≈ 65 MeV) of the total A_1 width. If, however, one tentatively uses $\alpha \approx 0.015$ as determined in the non-degenerate case, one gets a very small $\Gamma(A_1 \rightarrow \epsilon \pi)$ which means $\delta \simeq -1$ in Eq. (2).

In summary, we have noted the various ambiguities in the $\pi \epsilon \rightarrow \pi \epsilon$ Veneziano amplitude. If one adopts the attitude that only the complete amplitude and not each of the terms separately need to have the full asymptotic behaviour (therefore, we may put $E_1 = 0$) one relates the width $\Gamma(A_1 \rightarrow \epsilon \pi)$ to $\Gamma(f \rightarrow 4 \pi^0)$ or equivalently to $\Gamma(f \rightarrow \epsilon \epsilon \rightarrow 2 \pi^+ 2 \pi^-)$. If one further exploits the presence of the satellite term to impose factorization at the ϵ pole ^{*)}, one can even predict these widths and G_S/G_D as in Eq. (16) above. Experimental information on x or on $\Gamma(f \rightarrow 4 \pi^0)$ would be of further help to determining $g_{f \epsilon \epsilon}$.

*) In the degenerate case, this was essential even to get a relation between $\Gamma(A_1 \rightarrow \epsilon \pi)$ and $\Gamma(f \rightarrow 4 \pi^0)$.

ACKNOWLEDGEMENT

We are indebted to Professor M. Jacob for guidance, discussing and reading of the manuscript and we thank Dr. Ch. Schmid for useful discussions. One of us (GVD) is grateful to Dr. A.N. Kamal for collaboration in the early stage. He is also thankful for the hospitality of the CERN Theoretical Study Division.

A P P E N D I X

Our notation and some formulae are given below. Our metric differs from that of Ref. 7).

$$A_\mu A'_\mu = A_0 A'_0 - \vec{A} \cdot \vec{A}' ; \quad h = c = 1 .$$

The GeV is taken as the unit of mass. The various Lagrangians and widths, etc., are defined as

$$\begin{aligned} 1) \quad \varepsilon \pi \pi : \quad & g_{\varepsilon \pi \pi} \varepsilon (\vec{\pi} \cdot \vec{\pi}) \\ & \Gamma_{\text{total}} (\varepsilon \rightarrow \pi \pi) = \frac{3 g_{\varepsilon \pi \pi}^2 (m^2 - 4\mu^2)^{1/2}}{32 \pi m^2} \\ & = \frac{g}{2} \Gamma_{\text{total}} (\rho \rightarrow \pi \pi) = 540 \text{ MeV.} \end{aligned}$$

$$2) \quad f \varepsilon \varepsilon : \quad g_{f \varepsilon \varepsilon} \varepsilon_{\mu\nu} p_\mu^\varepsilon p_\nu^\varepsilon$$

$$\begin{aligned} 3) \quad f \pi \pi : \quad & g_{f \pi \pi} \varepsilon_{\mu\nu} p_\mu^\pi p_\nu^\pi (\vec{\pi} \cdot \vec{\pi}) \\ & \Gamma_{\text{total}} (f \rightarrow \pi \pi) = \frac{g_{f \pi \pi}^2 (m^2 - 4\mu^2)^{5/2}}{40 (32 \pi m_f^2)} = 145 \text{ MeV}^{(12)} \end{aligned}$$

$$\begin{aligned} 4) \quad f \rightarrow \varepsilon \varepsilon \rightarrow 2\pi^+ 2\pi^- : \\ \Gamma_{\text{total}} = \frac{g_{f \varepsilon \varepsilon}^2 g_{\varepsilon \pi \pi}^4}{(120)(128) m_f \pi^5} I \end{aligned}$$

where I is the integral over phase space, etc., defined as

$$\begin{aligned} I = & \int_{2\mu}^{M-2\mu} dx \frac{\sqrt{x^2 - 4\mu^2}}{(x^2 - m^2 + \frac{\Gamma^2}{4})^2 + m^2 \Gamma^2} \cdot \int_x^{(x^2 + M^2 - 4\mu^2)/2M} dQ_0 (Q_0^2 - x^2)^{5/2} \\ & \cdot \sqrt{\frac{M^2 + x^2 - 2MQ_0 - 4\mu^2}{M^2 + x^2 - 2MQ_0}} \cdot \frac{1}{(M^2 + x^2 - 2MQ_0 - m^2 + \frac{\Gamma^2}{4})^2 + m^2 \Gamma^2} \end{aligned}$$

Here Γ is the total width $\Gamma(\epsilon \rightarrow \pi\pi)$ and M in this formula (only) is the mass of the f meson. This form arises using the methods of Ref. 13). The integration was done numerically.

$$5) \quad A_1 \epsilon \pi : \quad g_{A_1 \epsilon \pi} (\vec{A}_1 \cdot \vec{\pi}) \epsilon_\mu^{A_1} p_\mu^\pi$$

$$\Gamma(A_1 \rightarrow \epsilon \pi) = \frac{g_{A_1 \epsilon \pi}^2}{24\pi} \cdot \frac{|\vec{p}|^3}{M^2}$$

where $\vec{p}$ is the $\epsilon \pi$ centre-of-mass momentum.

$$\Gamma(A_1 \rightarrow \epsilon \pi \rightarrow 3\pi) = \frac{g_{A_1 \epsilon \pi}^2 g_{\epsilon \pi \pi}^2}{192 \pi^3 M} J$$

where

$$J = \int_{\mu}^{\omega_0} p^3 d\omega \sqrt{\frac{Q^2 - 4\mu^2}{Q^2}} \cdot \left[\left\{ m^2 - \frac{\Gamma^2}{4} - Q^2 \right\}^2 + m^2 \Gamma^2 \right]^{-1}$$

where

$$\omega_0 = \frac{M^2 - 3\mu^2}{2M}$$

and

$$Q^2 = M^2 + \mu^2 - 2\omega M, \quad p^2 = \omega^2 - \mu^2, \quad \Gamma = \Gamma_{\text{total}}(\epsilon \rightarrow \pi\pi)$$

This form is obtained by the methods of Ref. 13) and the integration was done numerically.

$$6) \quad \epsilon \epsilon \epsilon : \quad g_{\epsilon \epsilon \epsilon}$$

$$7) \quad \rho \pi \pi : \quad g_{\rho \pi \pi} \epsilon^\rho (p_1^\pi - p_2^\pi)$$

$$\Gamma_{\text{total}} = \frac{g_{\rho \pi \pi}^2}{48 \pi m^2} (m^2 - 4\mu^2)^{3/2}$$

$$8) \quad A_1 \rho \pi : \quad -G_S \epsilon^{A_1} \cdot \epsilon^\rho + 2 G_D \epsilon^{A_1} \cdot p_\rho \epsilon^\rho \cdot p_{A_1}$$

$$\Gamma(A_1 \rightarrow \rho \pi) = \frac{|\vec{p}|}{24 \pi M^2} \cdot$$

$$\cdot \left\{ G_S^2 \left(3 + \frac{\vec{p}^2}{m^2} \right) - 4 G_S G_D \frac{M \omega_\rho \vec{p}^2}{m^2} + 4 G_D^2 \frac{M^2 \vec{p}^4}{m^2} \right\}$$

where $\vec{p}$ is the $\rho \pi$ centre-of-mass momentum and $\omega_\rho^2 = \vec{p}^2 + m^2$.

R E F E R E N C E S

- 1) See, for example, L. Brown and H. Munczek, Phys.Rev.Letters 20, 680 (1968);
H.J. Schnitzer and S. Weinberg, Phys.Rev. 164, 1828 (1967);
I. Gerstein and H.J. Schnitzer, Phys.Rev. 170, 1638 (1968);
R. Arnowitt, M.H. Friedman and P. Nath (to be published).
- 2) See, for example, H.J. Lipkin, Phys.Rev. 159, 1303 (1967);
J.G. Kuriyan and M. Suzuki, Phys.Rev. 169, 1385 (1968).
- 3) M. Ademollo et al., Nuovo Cimento 51A, 227 (1967).
- 4) See, for example, F.J. Gilman and H. Harari, Phys.Rev. 165, 1803 (1968);
P.H. Frampton and J.C. Taylor, Nuovo Cimento 49A, 152 (1968).
- 5) J. Ballam et al., Phys.Rev.Letters 21, 934 (1968).
- 6) S.P. de Alwis and D.A. Nutbrown, Cambridge preprint DAMTP 69/40 entitled "The algebra of currents and the Veneziano model - the $A_1 \rho \pi$ system";
C.J. Goebel, M.L. Blackmon and K.C. Wali, Wisconsin preprint, entitled "The Veneziano model for $\pi\pi \rightarrow \pi S$ where S has arbitrary spin and parity";
P. Brooker and J.M. Kosterlitz, Oxford preprint, entitled "Hard pion results from the Veneziano model".
- 7) Fayyazuddin and Riazuddin, Phys.Letters 28B, 561 (1969);
Rawalpindi, preprint entitled "Veneziano model for meson systems".
- 8) G. Veneziano, Nuovo Cimento 57, 190 (1968).
- 9) C. Lovelace, Phys.Letters 28B, 265 (1968).
- 10) M. Jacob, CERN preprint, Ref. TH. 1010, entitled "Duality in strong interaction physics".
- 11) G. Ascoli, H.B. Crawley, D.W. Mortara and A. Shapiro, Phys.Rev. Letters 21, 1712 (1968).
- 12) A.H. Rosenfeld et al., Berkeley report UCRL-8030 Part I, entitled "Review of particle properties".
- 13) R.H. Dalitz, Phys.Rev. 99, 915 (1955);
L.B. Okun and E.P. Shabalin, Soviet Phys. - JETP 10, 1252 (1960).