CM-P00058194

Archives

Ref.TH.1074-CERN

FINAL STATE INTERACTIONS AND CHARGE ASYMMETRY IN $K_{\ell 3}^L$ DECAYS

J.S. Bell and R.K.P. Zia *)

CERN - Geneva

A B S T R A C T

It is argued that final state interaction effects can become appreciable only if certain CP violating amplitudes are unexpectedly large. If such large hitherto hidden violations exist, then variation of the asymmetry over the Dalitz plot can yield otherwise unobtainable information about them.

---

*) NATO Postdoctoral Fellow ;
Address after October 15th : University of Birmingham,
Birmingham, England.

1. - INTRODUCTION

This work is concerned with corrections to the formula ¹⁾

$$\delta = (|p|^2 - |q|^2) \frac{1 - |x|^2}{1 + |x|^2} \quad (1)$$

relating the fractional charge asymmetry in the decays

$$K_L \rightarrow \pi^\pm \ell^\mp \nu \quad (2)$$

to the CP violation parameter ($|p|^2 - |q|^2$) and the $\Delta S = \Delta Q$ violation parameter x , with CPT assumed. The derivation of this formula requires the neglect of electromagnetic final state interactions. Especially if the electromagnetic interaction itself violated C, it might be feared ²⁾ that the correction would be of order of magnitude α , comparable to the observed ³⁾ $\delta \approx 3 \cdot 10^{-3}$. In this connection, Ryan ²⁾ has made essentially the following points.

- (a) - Final state interaction effects cancel when the asymmetry in total semi-leptonic decay rates is considered, i.e., taking along with (2) modes with extra photons and/or pions.
- (b) - These other modes are relatively rare. Those with extra photons are suppressed by powers of α , and those with extra pions by phase space. Moreover, asymmetries in these modes also arise (assuming CPT) only from electromagnetic final state interactions, or involve the small quantity $|p|^2 - |q|^2$.
- (c) - In view of (b) the cancellation (a) does not permit a large electromagnetically induced asymmetry in (2). The correction to (1), even with large CP violation, would be expected to be of order only

$$\alpha \Gamma(K_{\ell 4}) / \Gamma(K_{\ell 3}) \approx 10^{-3} \alpha .$$

As stressed by Ryan, these arguments apply only to the asymmetry in the total rates for (1), and leave open the question of what happens locally in the Dalitz plot. Now it seems that results so far available ³⁾ are based on an unspecified selection of the various modes, not necessarily uniform over the Dalitz plot of (2). So we have thought

it worthwhile to make more detailed considerations ^{*}). The effects remain negligible, unless the CP violation in certain weak amplitudes is grossly larger than any hitherto observed. This is just because even a maximal violation of C in electromagnetism should not show up ⁴⁾ in what would otherwise be the dominant final state interaction, namely πl scattering with one photon exchange. Nevertheless, as our calculations will illustrate, study of the variation of the asymmetry over the Dalitz plot can in principle yield otherwise inaccessible information on symmetry violating phases in weak decay amplitudes.

2. - GENERALITIES

Let F be an l^+ type final state and $\bar{F}'$ its CPT conjugate ^{**}). Suppose that in the experiment different modes are weighted (voluntarily or involuntarily) with weight w_F such that

$$w_F = w_{\bar{F}'} . \quad (3)$$

Then we are concerned with the asymmetry

$$\delta = \frac{\sum w_F \{ |K(F|T|L)|^2 - |K(\bar{F}'|T|L)|^2 \}}{\sum w_F \{ |K(F|T|L)|^2 + |K(\bar{F}'|T|L)|^2 \}} . \quad (4)$$

When final state interactions are ignored, CPT gives

$$(\bar{F}'|T|L) = (F|T|\bar{L}')^* \quad (5)$$

where

$$\begin{aligned} |L\rangle &= p|K\rangle + q|\bar{K}\rangle \\ |\bar{L}'\rangle &= q^*|K\rangle + p^*|\bar{K}\rangle . \end{aligned} \quad (6)$$

---

^{*}) Some of these considerations have been made already by C. Bouchiat and J. Prentki (unpublished).

^{**}) The notation is that of J.S. Bell and J. Steinberger ⁵⁾.

With the definition

$$x_F = \frac{q(F|T|\bar{K})}{P(F|T|K)} \quad (7)$$

we have

$$\frac{(F|T|\bar{L}')}{(F|T|L)} = \frac{\frac{q^*}{p} + \frac{p^*}{q} x_F}{1 + x_F} \quad (8)$$

Then to lowest order in $|p|^2 - |q|^2$

$$\delta = (|p|^2 - |q|^2) \frac{\sum w_F |(F|T|L)|^2 (1 - |x_F|^2) (1 + |x_F|)^{-2}}{\sum w_F |(F|T|L)|^2} \quad (9)$$

This reduces to (1) if x_F is supposed independent of F .

Allowing now for final state interactions we have instead of (5) *)

$$(\bar{F}'|T|L) = (F|S'T|\bar{L}')^* \quad (10)$$

where S is the scattering matrix of final state interactions. In estimating the additional effects we will ignore the small quantity $|p|^2 - |q|^2$. Then up to a possible phase factor

$$|\bar{L}'\rangle = |L\rangle$$

---

*) This is just the generalization of the more familiar formula, quoted for example in ⁵⁾ $(\bar{F}'|T|L) = (F|T|\bar{L}')^* e^{2i\delta}$, to the case in which F is not an eigenstate of the scattering matrix.

and (4) becomes

$$\delta = \frac{\sum w_F \{ |F|T|L|^2 - |F|S^*T|L|^2 \}}{\sum w_F \{ |F|T|L|^2 + |F|S^*T|L|^2 \}}. \quad (11)$$

This is what must be added to the $|p|^2 - |q|^2$ term (9).

When the summation over F includes multipion states, not only electromagnetic but also strong final state interactions enter. The effects of the latter would cancel out, because of the unitarity of the S matrix, if all states scattered into one another by strong interactions are equally weighted - e.g., all states with given final hadron mass and G parity (if G parity is conserved by strong interactions). But even without this it seems that such modes are unlikely to contribute significantly in experiments not specially designed to select them. They are expected to be quite rare, because of phase space limitation. It seems that these modes have not in fact been seen for neutral kaons. However, if the $\Delta I = \frac{1}{2}$ rule is not quite wrong one can anticipate

$$\frac{\Gamma(K_L \rightarrow \pi^0 \pi^- e^+ \nu)}{\Gamma(K_L \rightarrow \pi^- e^+ \nu)} < \frac{\Gamma(K^+ \rightarrow \pi^+ \pi^- e^+ \nu)}{\Gamma(K^+ \rightarrow \pi^0 e^+ \nu)} \approx 7 \times 10^{-4}. \quad (12)$$

The three pion mode, energetically just possible, is expected to be much rarer still, by several orders of magnitude ⁶⁾. Thus if the experiment does not especially select the multipion modes, only a gross asymmetry therein (more than 13%) will contribute to δ at the level 10^{-4} . We tentatively ignore this possibility, and omit multipion modes in (11) ^{*}).

---

^{*}) Note, by the way, that these modes can be excluded by the kinematic restriction $-2n_\mu K_\mu > (m_K^2 - 4m_\pi^2)$, where n_μ and K_μ are neutrino and kaon four-momenta.

We are then concerned only with final state interactions that arise electromagnetically. With

$$S^{-1} = 1 + iV$$

and to first order only in α

$$\delta = \frac{\sum_F w_F 2 \operatorname{Im} (F|T|L)^* (F|V|T|L)}{\sum_F w_F 2 |(F|T|L)|^2} . \quad (13)$$

Moreover to this order the summation can now be restricted to the $\pi l \nu$ states. For the presence in F of a photon would imply an extra factor α , in addition to that from the final state interaction.

There is still implicit in (13) a sum over all possible final states F' :

$$\delta = \frac{\sum_F \sum_{F'} w_F 2 \operatorname{Im} (F|T|L)^* (F|V|F')(F'|T|L)}{\sum_F w_F 2 |(F|T|L)|^2} . \quad (14)$$

Again states F' with photons will enter only at the next order in α , and so will be ignored. Multipion states can contribute in lowest order of α , but again should be greatly suppressed by phase space. In fact if we had $w_F = 1$, so that (14) were essentially antisymmetric between intermediate and final states, the contribution of multipion intermediate states F' would be like the contribution of multipion final states F - which we supposed above to be negligible. We have made more detailed considerations for simple models (Appendix), and it seems indeed unlikely that multipion states F' in (14) are significant. We now leave them out ^{*)}.

---

^{*)} Note that multipion intermediate states cannot contribute in the region below threshold, $-2n_{\mu} K_{\mu} > (m_K^2 - 4m_{\pi}^2)$.

Note now that with only the three-body ($\pi l \nu$) states involved, the scattering matrix element in (14) cannot exhibit time reversal failure. For we are concerned simply with one photon exchange between electrons and pions on their mass shells. Hermiticity and the continuity equation are then sufficient to forbid the non-invariant terms ⁴⁾. Now of course if time reversal (or CP, assuming CPT) is good also for the weak matrix elements, δ will vanish ^{*}). For the time reversed configuration is equivalent to the original when spins are summed over, and the matrix elements become complex conjugated, so that the imaginary part must vanish. Thus

$$\delta \sim \alpha x (\text{degree of time reversal violation in } K_{l3} \text{ amplitude})$$

In the absence of a quite gross violation this should be small compared with 10^{-3} .

3. - EXAMPLES

For explicit calculation it is convenient to define the local asymmetry

$$\delta_F = \frac{\text{Im} \sum_F (LITIF)(FIVIF')(F'ITIL)}{(LITIF)(FITIL)} \quad (15)$$

in which summation over spins is understood. Let the four-momenta be labelled

K_μ	kaon	n_μ	neutrino
p_μ	final pion	p'_μ	intermediate pion
k_μ	final lepton	k'_μ	intermediate lepton

---

^{*}) The converse is not necessarily true.

It is convenient to work in the centre-of-mass system of the charged lepton pair. Then

$$\delta_F = \alpha \int \frac{d^3 \vec{k}'}{4\pi k'_0 p'_0} \delta(k'_0 + p'_0 - k_0 - p_0) \frac{f_\pi((k_\mu - k'_\mu)^2)}{(k_\mu - k'_\mu)^2} I$$

$$I = \frac{\text{Im Trace } \bar{W}(m - i\gamma k) i\gamma_\mu (p + p')_\mu (m - i\gamma k') W'(-i\gamma n)}{\text{Trace } \bar{W}(m - i\gamma k) W(-i\gamma n)} \quad (16)$$

where m is lepton mass, $f_\pi(q^2)$ is the pion electromagnetic form factor, and the weak matrix element is

$$W = f(q^2) \{ i\gamma(K+p) + \xi(q^2) i\gamma(K-p) \} (1 + \gamma_5) \quad (17)$$

with $q = K - p$. From W , W' is obtained by $p \rightarrow p'$, and $\bar{W} = \gamma_4 W^\dagger \gamma_4$.

Define angles θ, θ', ϕ' so that

$$\begin{aligned} \hat{n} \cdot \hat{k} &= \cos \theta \\ \hat{k} \cdot \hat{k}' &= \cos \theta' \\ \hat{n} \cdot \hat{k}' &= \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos \phi' \end{aligned} \quad (18)$$

where a circumflex (^) indicates a unit vector. Then (16) becomes

$$\delta_F = \alpha \frac{k}{\omega} \int \frac{d\cos \theta' d\phi'}{4\pi} \frac{f_\pi(2k^2(1 - \cos \theta'))}{2k^2(1 - \cos \theta')} I \quad (19)$$

where

$$\omega = k_0 + p_0, \quad k = (k_0^2 - m^2)^{1/2} = (p_0^2 - m_\pi^2)^{1/2}.$$

As a first example suppose that the form factors in (18) are approximated by constants. Then the only way in which the upper trace in (16) can develop an imaginary part is by interference between the two terms in (18), involving $\text{Im} \xi$. Picking out this contribution we find

$$I = -\text{Im} \xi \frac{8\omega k m^2 p_0 (\hat{k} \cdot \hat{n} - \hat{k}' \cdot \hat{n})}{A(\omega) + B(\omega) \cos \theta} \quad (20)$$

$$\begin{aligned} A(\omega) &= k_0 \{ 4\omega^2 + m^2 |1 - \xi|^2 \} - 4\omega m^2 (1 - \text{Re} \xi) \\ B(\omega) &= k \{ 4\omega^2 - m^2 |1 - \xi|^2 \} \end{aligned} \quad (21)$$

whence

$$\delta_F = -\alpha \text{Im} \xi \frac{C(\omega) \cos \theta}{A(\omega) + B(\omega) \cos \theta} \quad (22)$$

$$C(\omega) = 4m^2 p_0 \frac{1}{2} \int d\cos \theta' f_\pi(2k^2(1 - \cos \theta'))$$

With the rough approximation $f_\pi(q^2) \approx f_\pi(0) = 1$, then $C(\omega) \approx 4m^2 p_0$, and ^{*})

$$\delta_F = - \frac{4m^2 p_0 \alpha \text{Im} \xi \cos \theta}{A(\omega) + B(\omega) \cos \theta} \quad (23)$$

There is a slight problem in discussing the magnitude of (23) if

$$2 \left(\frac{m_g + m}{m} \right) > 1 - \text{Re} \xi > 2 \left(\frac{\omega}{k_0 + k} \right)_{\min} \approx 2.2$$

---

^{*}) An equivalent formula was extracted by Ryan ²⁾ from Ginsberg's paper on radiative corrections ⁷⁾. His equation (4.13) has a misprint unfortunately.

This is perhaps not entirely excluded in the present confused experimental situation ^{*)}. Then there would be a point on the boundary of the physical region where the denominator of (23) would fail to vanish only in virtue of $\text{Im } \xi$. Thus large asymmetries would be possible in a small region of the Dalitz plot ^{**)}, and conceivably this could become an interesting way to study $\text{Im } \xi$. However this region is one in which the mean density in the Dalitz plot, precisely the denominator in (23), becomes very small - i.e., where there are few counts anyway. So it would seem that only by very bad luck, or deliberate design, would this region be weighted appreciably in an experiment on global asymmetry. Suppose for example we integrate the numerator and denominator in (23), separately, over negative values of $\cos\theta$ - i.e., replace $\cos\theta$ by $-\frac{1}{2}$. Then the resulting expression does not anywhere exceed, for $\text{Re } \xi$ negative,

$$(0.4) \propto \text{Im } \xi .$$

For this to reach the 10^{-4} level would require $|\text{Im } \xi| \geq 0.03$. To reach the level (1.35×10^{-3}) of the error in the reported experiment would require $\text{Im } \xi \gtrsim \frac{1}{2}$.

The currently quoted experimental value for $\text{Im } \xi$, from measurements of transverse muon polarization, is ^{***)}

$$\text{Im } \xi = (-1.25 \pm 6.6) \times 10^{-2}.$$

For K_{e3} decay one expects terms containing the lepton mass m to be quite negligible. Setting $m = 0$ makes (23) zero. Then to obtain an effect some allowance must be made for non-constancy of the form factor f in (17). With $m = 0$,

$$I = -2\omega k_0 \frac{1 + \hat{n} \cdot \hat{k} + \hat{n} \cdot \hat{k}' + \hat{k} \cdot \hat{k}'}{1 + \hat{n} \cdot \hat{k}} \text{Im} \left(\frac{f'}{f} \right) . \quad (24)$$

*) Values of $\text{Re } \xi$ range from -0.4 to -3.9 ⁸⁾.

**) Of course (22) would not be adequate right at the maximum.

***) Here also, when the accuracy improves considerably, allowance for final state interaction effects will be necessary ^{9), 10)}.

To permit explicit integration we suppose

$$f(q^2) = (1 - \lambda q^2) f(0) \quad (25)$$

with λ small. To first order in λ

$$\begin{aligned} (f'/f) &= 1 - \lambda(q'^2 - q^2) \\ &= 1 + 2\lambda(\hat{n} \cdot \hat{k} - \hat{n} \cdot \hat{k}') n_0 k_0. \end{aligned}$$

Then

$$\begin{aligned} \delta_F &= -2\alpha n_0 k_0 \operatorname{Im} \lambda \left(\frac{3}{2} \cos \theta - \frac{1}{2} \right) \\ &\quad \times \frac{1}{2} \int d\cos \theta' (1 + \cos \theta') f_\pi(2k^2(1 - \cos \theta')). \end{aligned} \quad (26)$$

If we again set $f_\pi \simeq 1$, the last factor is unity :

$$\delta_F \simeq -(3\cos \theta - 1) \alpha n_0 k_0 \operatorname{Im} \lambda. \quad (27)$$

Now $4k_0 n_0 < (m_K - m_\pi)^2 \approx 6.5 m_\pi^2$. So for (27) to reach the 10^{-4} level would require

$$m_\pi^2 \operatorname{Im} \lambda \gtrsim 0.0021$$

If this is compared with the measured

$$m_\pi^2 \operatorname{Re} \lambda = 0.019 \pm 0.008$$

or with the inverse mass squared of K^* as "natural unit"

$$m_\pi^2 m_{K^*}^{-2} \approx 0.025$$

it can be regarded as a rather large T (or CP) violation. There is of course no experimental information on $\operatorname{Im} \lambda$, and it is difficult to envisage how else such a phase variation of the weak amplitude could be studied *).

---

*) Of course if the linear form (25) is taken quite literally, $\operatorname{Im} \lambda$ contributes to the q^4 terms in the square of the modulus. But this could not be distinguished from the presence of real q^4 terms in the amplitude.

Note that the $\cos\theta$ dependent part of (26) can be rewritten, to correspond in form to (22), as

$$\frac{3\cos^2\theta + 2\cos\theta - 1}{\cos\theta + 1} \quad (28)$$

In both cases the denominators, coming from the lower trace in (17), are essentially the $\cos\theta$ dependent factors in the unperturbed Dalitz densities ρ . Then in both cases $\delta_F \rho$ averages to zero over $\cos\theta$ ^{*)}. Let us introduce as more conventional Dalitz variables the energies of neutrino and lepton in the kaon system

$$\begin{aligned} E_\nu &= -n_\mu K_\mu m_K^{-1} \\ E_l &= -k_\mu K_\mu m_K^{-1} \\ &= m_K^{-1} (k_0 K_0 - k n_0 \cos\theta) \end{aligned}$$

where the components k_0, K_0, n_0, k (in the charged pair centre-of-mass system) are functions of E_ν . Thus the asymmetry $\delta_F \rho$ averages to zero over E_l at fixed E_ν . This feature of the examples is in fact a more general result. It follows simply from the unitarity of S in (11), and from the fact that S does not connect states with different E_ν , because neutrino interactions are neglected.

---

^{*)} In the reported K_{e3} experiments ³⁾ it turns out that the detection efficiency is rather uniform with respect to $\cos\theta$, so that the final state interaction effect (27) is largely suppressed (K. Kleinknecht and J. Steinberger, private communication).

4. - CONCLUSION

The effect of final state interactions is the redistribution of events within kinematical regions bounded by constant values of $E_\nu = -n \mu_K \mu_K^{-1}$, the neutrino energy in the kaon system. They can be eliminated by integrating uniformly over any such region, for ℓ^+ and ℓ^- separately, and forming a fractional asymmetry from the results. However, we have found that even without such careful weighting, final state interaction effects can become appreciable only if certain CP violating amplitudes are large to a quite unexpected degree - unexpected even in a theory with "maximal" C violation in the electromagnetic current. If such large hitherto hidden violations exist then variation of the asymmetry over the Dalitz plot can yield information about them. In particular we saw, in (26), how a CP violating phase in the K_{e3} amplitude could show up. It is difficult to envisage how else it could be studied.

We are indebted for useful discussions especially to Drs. M. Jacob, C. Llewellyn Smith, J. Prentki, E. de Rafael and J. Steinberger. One of us (JSB) was interested in this problem by Dr. C. Ryan during a period at the Dublin Institute for Advanced Studies, and thanks his colleagues there for that pleasant time, and especially Dr. Ryan for several conversations on this topic. The other (RZ) is grateful to Drs. J. Prentki and W. Thirring for the hospitality extended to him in the Theoretical Study Division.

APPENDIX

We will here try to estimate the effect of including the two pion intermediate state in the summation over F' in (15). There is a slight complication from the fact that these states can suffer strong interaction scatterings. We have in fact for the electromagnetic increment V in S^{-1} :

$$iV = \delta S^{-1} = -S_{st}^{-1} \delta S_{em} S_{st}^{-1} .$$

So

$$\langle F | V | F' \rangle = \langle F | +i \delta S_{em} | F' \rangle e^{-2i\delta(F')} \quad (29)$$

where $\delta(F')$ is the strong scattering phase of the state F' , supposed to be an eigenstate of the strong S matrix. The exponential in (29) just cancels against the "Watson phase factors" $e^{i\delta(F')}$ in $\langle F | +i \delta S | F' \rangle$ and $\langle F' | T | L \rangle$. So we have simply to omit these strong interaction phases.

We will only consider the electron case, for which the phase space is less restrictive than for the muon. Setting the lepton mass zero and supposing helicity conservation simplifies things a lot. To begin with there is no spin summation implicit in (15), and we can divide out :

$$\delta_F = \sum_{F'} \text{Im} \langle F | V | F' \rangle \langle F' | T | L \rangle / \langle F | T | L \rangle . \quad (30)$$

The structure of the numerator is indicated in the diagram :

This also indicates how we label four-momenta.

In the weak matrix element we retain only the presumably dominant axial form :

$$(G/\sqrt{2}) \tilde{g} i\gamma_\mu (1 + \gamma_5) (r_\mu - s_\mu) \quad (31)$$

and we take $\tilde{g}$ constant ; the tilda is the conventional indication that the final state interaction phase has been removed ^{*)}.

The parity conserving electromagnetic vertex has the form

$$4\pi\alpha \tilde{h} \epsilon_{\mu\nu\rho\sigma} p_\nu q_\rho r_\sigma i\gamma_\mu q^2 \quad (32)$$

and we take $\tilde{h}$ constant.

In the system $\vec{r} + \vec{s} = 0$ the product of matrix elements is

$$q^2 (4\pi\alpha G/\sqrt{2}) \tilde{g} \tilde{h} \{ +i(p_0 + q_0) i\vec{\gamma} \times \vec{q} \cdot \vec{r} \} i\gamma_{k'} 2i\vec{\gamma} \cdot \vec{r} (1 + \gamma_5) \quad (33)$$

Performing the integration

$$(2\pi)^3 \int \frac{d^3\vec{r}}{2r_0 2s_0} 2\pi \delta(r_0 + s_0 - \omega') \quad (34)$$

---

^{*)} For a recent account of K_{e4} decay, and earlier references, see Berends et al. ¹¹⁾

leads to

$$\bar{q}^2 (4\pi\alpha G/\sqrt{2}) \hat{g} \hat{h} F (i\vec{\gamma}_x \vec{q}) (i\gamma k') \cdot 2i\vec{\gamma} (1+\gamma_5) \{ + i(p_0 + q_0) \} \quad (35)$$

where

$$F = \frac{\omega'^2}{96\pi} \left(1 - \left(\frac{2m_0}{\omega'} \right)^2 \right)^{3/2} \quad (36)$$

and

$$\omega'^2 = -(p+q)^2 = -(p+k-k')^2 = \omega^2 + 2(p+k) \cdot k'. \quad (37)$$

The expression (35) has the invariant form

$$\bar{q}^2 (4\pi\alpha G/\sqrt{2}) \hat{g} \hat{h} F \epsilon_{\mu\nu\rho\sigma} i\gamma_\mu p_\nu q_\rho i\gamma k' 2i\gamma_5 (1+\gamma_5). \quad (38)$$

The spinor factor reduces further to

$$4(k' q_\mu - k' p_\mu) i\gamma_\mu (1+\gamma_5) \quad (39)$$

Recalling that $q = k - k'$, and $i\gamma k \equiv 0$, this can be replaced by

$$4(k' (p+k)_\mu + k' k (p+k)_\mu - k' k k'_\mu) i\gamma_\mu (1+\gamma_5) \quad (40)$$

Go now to the system

$$\vec{p} + \vec{k} = \vec{K} - \vec{n} = 0.$$

We can average immediately over the angle θ' defined in (18) :

$$k'_\mu = (k'_0, \vec{k}')$$

$$\rightarrow (k'_0, k'_0 \hat{k} \cos \theta')$$

$$\equiv (k'_0 (1 - \cos \theta'), 0)$$

(after subtracting a multiple of k)

$$\rightarrow (k'_0(1-\cos\theta'), \vec{0}) = \bar{w}' k'_0(1-\cos\theta')(p+k)_\mu.$$

So (40) becomes

$$4([k' \cdot (p+k) - k' \cdot k] \bar{w}' k'_0(1-\cos\theta') + k' \cdot k) i\delta(k+p)(1+\gamma_5). \quad (41)$$

As was to be expected (for there is only one helicity conserving coupling) the spinor factor now cancels against the corresponding K_{e3} factor

$$\begin{aligned} & \int i\delta(K+p)(1+\gamma_5) G/\sqrt{2} \\ & \equiv 2 \int i\delta(k+p)(1+\gamma_5) G/\sqrt{2}. \end{aligned}$$

Collecting terms we then have, for the two-pion induced asymmetry in the one pion final state,

$$\begin{aligned} \delta'' &= -\text{Im} \ 4\pi\alpha \ (\hat{g}\hat{h}/f) \int \frac{d^3\vec{k}'}{(2\pi)^3 2k'_0} F(w') \\ & \quad \frac{1}{k_0} (k'_0 - \frac{k_0 k'_0}{\omega} (1-\cos\theta') + k_0) \\ &= -\frac{\alpha}{\pi} \text{Im} \left(\frac{\hat{g}\hat{h}}{f} \right) \int dk'_0 \frac{d\cos\theta'}{2} F(w') k'_0 \left(1 + \frac{k'_0}{k_0} + \frac{k'_0}{\omega} (\cos\theta' - 1) \right). \end{aligned} \quad (42)$$

With

$$\begin{aligned} F(w') &= \frac{w'^2}{96\pi} \left(1 - \left(\frac{2m_\pi}{w'} \right)^2 \right)^{3/2} \\ w'^2 &= \omega(\omega - 2k'_0) \\ \omega &= k_0 + \sqrt{m_\pi^2 + k_0^2}, \end{aligned} \quad (43)$$

$$\delta'' = -\frac{\alpha}{96\pi} \text{Im} \left(\frac{\hat{g}\hat{h}}{f} \right) \int dk'_0 k'_0 w'^2 \left(1 - \left(\frac{2m_\pi}{w'} \right)^2 \right)^{3/2} \left(1 + \frac{k'_0(\omega - k_0)}{k_0 \omega} \right). \quad (44)$$

This is nowhere bigger than

$$(\alpha/48\pi^2) \operatorname{Im}(f'\tilde{g}\tilde{h}) m_K^4 \int dx \, x (1-2x) \left(1 - \frac{4m_\pi^2/m_K^2}{1-2x}\right)^{3/2}$$

$$< (\alpha/48\pi^2) \operatorname{Im}(f'\tilde{g}\tilde{h}) m_K^4 \int dx \, x (1-2x - 4m_\pi^2/m_K^2).$$

So

$$|\delta''| < 3 \cdot 10^{-5} \alpha |\operatorname{Im} f'\tilde{g}\tilde{h} m_K^4| \quad (45)$$

Experiments on K_{e4}^{L} decay have not been done. On the basis of the $\Delta I = \frac{1}{2}$ rule and charged kaon decays

$$f'\tilde{g} m_K \approx 10. \quad (46)$$

To get some notion of a reasonable magnitude for $\tilde{h}$ we use vector dominance, giving

$$\tilde{h} m_K^3 \approx 9. \quad (47)$$

With these magnitudes, and with arbitrary T violating phases in the couplings,

$$|\delta''| \lesssim 2 \cdot 10^{-5} \quad (48)$$

which is negligible.

Actually, if we take account of crossing symmetry, hermiticity requires $\tilde{h}$ to be real. To obtain a T violating (and C violating) vertex we have then to introduce an additional factor odd under exchange of positive and negative pion momenta, i.e., $-p$ and r :

$$q^2 i\tilde{h}' i\gamma_\mu \epsilon_{\mu\nu\tau\sigma} p_\nu q_\tau r_\sigma q \cdot (p+r).$$

This leads to (42) with $\tilde{h} \rightarrow i\tilde{h}'$, and with the extra factor in the integrand

$$\frac{1}{2} k'_0 k_0 (\cos \theta - 1) - \frac{3}{2} \omega (k_0 - k'_0).$$

The modulus of this extra factor is bounded by m_K^2 , so now instead of (45)

$$|\delta''| < 3 \cdot 10^5 \alpha | \operatorname{Re} f' \tilde{g} \tilde{h}' m_K^6 |.$$

Then we have the problem of guessing what the value might be of the highly dimensional coupling constant $\tilde{h}'$ in a theory with "maximal CP violation" in electromagnetism. If we give $\tilde{h}' m^5$ the same value as the "reasonable" CP conserving $\tilde{h} m^3$, namely (47), then we again have (48).

REFERENCES

- 1) T.D. Lee, R. Oehme and C.N. Yang - Phys.Rev. 106, 340 (1957) ;
S. Bennett, D. Nguyen, H. Saal, J. Steinberger and J. Sunderland -
Phys.Rev.Letters 19, 997 (1967).
- 2) C. Ryan - V.C.D. Preprint, and Trieste Preprint IC/69/92.
- 3) D. Dorfan, J. Enstrom, D. Raymond, M. Schwarz and S. Wojcicki -
Phys.Rev.Letters 19, 987 (1967).;
S.Bennett, D. Nguyen, H. Saal, J. Steinberger and J. Sunderland -
Phys.Rev.Letters 19, 993 (1967) ; Phys.Letters 27B, 239
(1968) ;
J. Steinberger - Report at Topical Conference on Weak Interactions,
CERN, January 1969.
- 4) J. Bernstein, G. Feinberg and T.D. Lee - Phys.Rev. 139B, 1850 (1965).
- 5) J.S. Bell and J. Steinberger - Proceedings of Oxford International
Conference on Elementary Particle Physics (1965).
- 6) V.A. Kolkunov and I.V. Lyagin - Soviet Phys. JETP 18, 1379 (1964) ;
P. McNamee and R.J. Oakes - Phys.Letters 24B, 1379 (1964).
- 7) E. Grinsberg - Phys.Rev. 171, 1675 (1968) ; Phys.Rev. 174, 2169
(1968).
- 8) C. Rubbia - Report at Topical Conference on Weak Interactions, CERN
pp. 240-248, January 1969.
- 9) K. Young, M. Longo and J. Helland - Phys.Rev.Letters 18, 806 (1967).
- 10) N. Byers, S. McDowell and C.N. Yang - Seminar on High Energy Physics
and Elementary Particles, Trieste, p. 953 (1965).;
L.B. Okun and I.B. Khriplowich - Phys.Letters 24B, 672 (1967).
- 11) F.A. Berends, A. Donnachie and G.C. Oades - Phys.Rev. 171, 1457 (1968).