CM-P00058170

Ref.TH.1145-CERN

FROISSART BOUNDS WITH NO ARBITRARY CONSTANTS

A.K. Common *)

CERN - Geneva

A B S T R A C T

A method is described for calculating rigorous upper bounds to averaged total cross-sections $\overline{\sigma}_{\text{tot}}(s)$ for $\pi\pi$ scattering defined by

$$\overline{\sigma}_{\text{tot}}(s) = \frac{(m+1)}{(s-4\mu^2)^{m+1}} \int_{4\mu^2}^s (s'-4\mu^2)^m \sigma_{\text{tot}}(s') ds'$$

for all energies and all $m \gg 1$, from the D wave scattering length α_2^t in the t channel or alternatively from a rigorous bound for the D wave amplitude $a_2(t_0)$ at one value of t_0 between 0 and $4\mu^2$.

It is shown that as $s \rightarrow \infty$, the bound to $\overline{\sigma}_{\text{tot}}(s)$ is equivalent to the Froissart bound

$$\overline{\sigma}_{\text{tot}}(s) \lesssim \frac{4\pi(m+1)}{t_0^m} [\log(s/s_0)]^2$$

where the scaling factor s_0 can be calculated from the given information and $t_0 = 4\mu^2$ when the scattering length is used. By letting $m \rightarrow \infty$ the usual bound is obtained.

Similar results are derived for the fixed angle, fixed t and forward $\pi\pi$ scattering amplitudes, and again the bounds are asymptotically equivalent to the corresponding Froissart bounds and contain no unknown constants.

*) On leave of absence from the University of Kent at Canterbury, England.

1. - INTRODUCTION

In a classic paper ¹⁾ Froissart showed that two-body reaction amplitudes satisfying Mandelstam's representation have well determined asymptotic properties. For instance the total cross-section $\sigma_{\text{tot}}(s)$ must be less than $A[\log(s/s_1)]^2$ as $s \rightarrow \infty$, where s is the square of the centre-of-mass energy and A and s_1 are constants. Martin and co-workers ²⁾ have been able to derive such bounds from the much weaker assumptions of unitarity, crossing and analyticity in a domain derived from axiomatic field theory, and also obtained a value for the constant A for certain scattering states.

It is not possible to really test such a bound against experiment since there is the uncertainty due to the unknown constant s_1 . More important, however, is the problem of knowing when the asymptotic region has been reached. Both of these difficulties have been removed recently by Yndurain ³⁾ who has obtained using the same assumptions as Ref. 2), a bound for a weighted integral of the total cross-section over all energy ranges given by

$$\begin{aligned} \bar{\sigma}_{\text{TOT}}(s) &\equiv \frac{2}{(s-4\mu^2)^2} \int_{4\mu^2}^s (s'-4\mu^2) \sigma_{\text{TOT}}(s') ds' \\ &\leq 2 \left[\frac{n+1}{n} \right]^2 \frac{\pi}{\mu^2} [\log(s/\mu^2)]^2 + \frac{8\pi}{s} \left(\frac{n+1}{n} \right) \sqrt{\frac{s}{\mu^2}} \log\left(\frac{s}{\mu^2}\right) + \frac{60\pi^2}{\mu M_n} \alpha_2^t; n=1,2,\dots \end{aligned} \quad (1.1)$$

where α_2^t is the D wave scattering length in the t channel and μ is the mass of the pion. The M_n 's are constants defined by

$$M_n = \frac{(2n)!}{2^n (n!)^2 \left[\frac{(2n+1)}{n+1} \right]^n} \sim \left[\frac{e}{\pi n} \right]^{1/2} \text{ as } n \rightarrow \infty \quad (1.2)$$

The bound (1.1) holds for all $s > 4\mu^2$ and as $s \rightarrow \infty$ it becomes equivalent to the Froissart bound

$$\overline{\sigma}_{\text{Tot}}(s) \lesssim \frac{2\pi}{\mu^2} [\log(s/\mu^2)]^2 \quad (1.3)$$

This result may be compared with the asymptotic behaviour of the actual cross-section which has been given by Jin and Martin ^{*}), i.e.,

$$\sigma_{\text{Tot}}(s) \lesssim \frac{\pi}{\mu^2} [\log(s/s_0)]^2 \quad (1.4)$$

It can be seen that although the scaling factor is now known, there is an extra over-all factor 2 in (1.3).

This factor can be removed by defining $\overline{\sigma}_{\text{tot}}(s)$ by

$$\overline{\sigma}_{\text{Tot}}(s) \equiv \frac{(m+1)}{(s-4\mu^2)^{m+1}} \int_{4\mu^2}^s (s'-4\mu^2)^m \sigma_{\text{Tot}}(s') ds', \quad m \geq 1, \quad (1.5)$$

and in place of (1.1) we have using the methods of Ref. 3),

$$\begin{aligned} \overline{\sigma}_{\text{Tot}}(s) &\leq \left[\frac{m+1}{m} \right] \left[\frac{n+1}{n} \right]^2 \frac{\pi}{\mu^2} [\log(s/\mu^2)]^2 + \left[\frac{m+1}{m-\frac{1}{2}} \right] \left[\frac{n+1}{n} \right] \frac{4\pi}{s^{\frac{1}{2}}\mu} \log\left(\frac{s}{\mu^2}\right) + \\ &+ (m+1) 30\pi^2 M_n^{-1} \alpha_2^+ \end{aligned} \quad (1.6)$$

By taking $m = \log(s/\mu^2)$ and letting $n \rightarrow \infty$ we get the asymptotic bound (1.4).

^{*}) See the second paper of Ref. 2).

In this paper we will give a refined technique for obtaining bounds to $\overline{\sigma}_{\text{tot}}(s)$ which uses methods developed by Lukaszuk and Martin ⁴⁾ for getting rigorous bounds to $\pi\pi$ scattering amplitudes in the unphysical region. We will again only need a bound to α_2^t or alternatively an upper bound to the D wave $a_2(t_0)$ for one value of t_0 between 0 and $4\mu^2$. The latter bound can be derived from the basic assumptions of unitarity, crossing, etc.

From the same information, we also derive similar bounds for the averages of the fixed angle, fixed t and forward amplitudes. The asymptotic behaviour of these bounds will be investigated and it will be proved that they correspond exactly to the Froissart bounds for these quantities, but with all their constants determined.

We consider in this paper $\pi^0\pi^0 \rightarrow \pi^0\pi^0$ but it is not difficult to extend our work to cover other charge states. In Section 2 we take the fixed angle amplitude $F(s,t)$ with $t = (2\mu^2 - s/2)(1 - \cos\theta)$ at $\theta \neq 0$ and show how, from the given information on the D wave amplitude, bounds may be derived on the averages

$$\bar{F}[s, (2\mu^2 - s/2)(1 - \cos\theta)] = \frac{(m+1)}{(s-4\mu^2)^{m+1/4}} \int_{4\mu^2}^s |F[s', (2\mu^2 - s'/2)(1 - \cos\theta)]| (s' - 4\mu^2)^{m-3/4} ds' \quad (1.7)$$

where $m \geq 3/4$, through the Froissart-Gribov formula

$$a_2(t_0) = \frac{4}{\pi(4\mu^2 - t_0)} \int_{4\mu^2}^{\infty} Q_2\left[\frac{2s}{(4\mu^2 - t_0)} - 1\right] \text{Im} F(s, t_0) ds \quad (1.8)$$

which holds for $0 < t_0 < 4\mu^2$ and the corresponding expression for α_2^t which comes from letting $t_0 \rightarrow 4\mu^2$ in (1.8). Similar results will be derived for the averages of the modulus of the forward scattering amplitude $F(s, 0)$ and the optical theorem is then used to get the bounds to $\overline{\sigma}_{\text{tot}}(s)$.

In Section 3, the asymptotic behaviour of these bounds will be investigated and we will find first of all that as $s \rightarrow \infty$ and for $\theta \neq 0$,

$$\bar{F}[s, (2\mu^2 - s_1)(1 - \cos\theta)] \leq \frac{2}{3\sqrt{\pi} \sin\theta} \left(\frac{s}{t_0}\right)^{\frac{3}{4}} \left[\log(s/s_0)\right]^{\frac{3}{2}} \left(\frac{m+1}{m+4}\right) \quad (1.9)$$

where t_0 is the value between 0 and $4\mu^2$ where we have the bound on $a_2(t_0)$. If the D wave scattering length a_2^t is used then $t_0 = 4\mu^2$. The scaling factor s_0 is given by

$$s_0 = \frac{s_1^{3/2}}{4\sqrt{t_0}} \log \left\{ s / \left[\frac{3}{8} \sqrt{\frac{s_1^3}{t_0}} \log(s/s_1) \right] \right\} \quad (1.10)$$

where *)

$$s_1^{3/2} = \frac{e(4\mu^2 - t_0)^2}{15\pi^2 \sqrt{t_0} a_2(t_0)} \quad (1.11)$$

Similarly defining the average of the fixed t amplitude for $t \neq 0$ by

$$\bar{F}(s, t) \equiv \frac{(m+1)}{(s - 4\mu^2)^m} \int_{4\mu^2}^s |F(s, t)| (s - 4\mu^2)^{m-1} ds \quad (1.12)$$

with $m \gg 1$, we prove that asymptotically it is bounded by

*) $a_2(t_0)$ may be replaced by its upper bound in this expression.

$$\bar{F}(s,t) \lesssim \frac{2}{3\sqrt{\pi}} \frac{s}{[-4t+t_0^3]^{1/4}} \left[\log(s/s_0) \right]^{\frac{3}{2}} \frac{(m+1)}{(m+1/4)} \quad (1.13)$$

with s_0 defined as previously.

The average for the forward scattering amplitude is defined as in (1.12) and we prove that

$$\bar{F}(s,0) \lesssim \frac{s}{4t_0} \left[\log(s/s_0) \right]^2 \left[\frac{m+1}{m} \right] \quad (1.14)$$

and finally using the optical theorem and defining $\bar{\sigma}_{\text{tot}}$ by (1.5) we find that

$$\bar{\sigma}_{\text{tot}}(s) \lesssim \frac{4(m+1)}{m} \frac{\pi}{t_0} \left[\log(s/s_0) \right]^2 \quad (1.15)$$

where s_0 is again defined by (1.10).

Letting $m \rightarrow \infty$ in (1.15), we have the bound (1.4) but with s_0 known. In fact the scaling factor s_0 is an increasing function of s for large s . The bounds (1.9), (1.13), (1.14) and (1.15) are therefore better than the corresponding Froissart bounds with constant scaling factors as for example the bound (1.3). Similarly improvements have already been obtained by Singh and Roy ⁵⁾ who have derived asymptotic bounds for the total cross-section, etc., which are essentially equivalent to the Froissart bounds but which depend on the value of the elastic cross-section at the energy under consideration.

In the last Section we will compare the bounds of these authors with our bounds. Also we give the results of numerical calculations of our bounds to $\bar{\sigma}_{\text{tot}}(s)$ for various energies and compare

them with those obtained by using the techniques of Yndurain and we will see that our method does give improved bounds.

Finally before passing on to the next Section, we should remark that although we have obtained asymptotic bounds for average cross-sections, etc., these may be converted into asymptotic bounds for the actual quantities if certain assumptions are made concerning their oscillatory behaviour for large s ⁶⁾. The averages for the total cross-section, the fixed-angle amplitude, etc., are defined in the different ways so that this should happen.

2. - INTEGRAL BOUNDS

We define our normalization so that the $\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$ scattering amplitude is given by

$$F(s, t) = \frac{\sqrt{s}}{k} \sum_{\substack{\ell=0 \\ \text{even}}}^{\infty} (2\ell+1) f_{\ell}(s) P_{\ell}(\cos \theta) \quad (2.1)$$

where $f_{\ell}(s) = e^{i\delta_{\ell}} \sin \delta_{\ell}$ and k is the centre-of-mass momentum. The partial wave amplitude $a_{\ell}(t_0)$ given by the Froissart-Gribov formula,

$$a_{\ell}(t_0) = \frac{4}{\pi(4\mu^2 - t_0)} \int_{4\mu^2}^{\infty} Q_{\ell} \left[\frac{2s'}{4\mu^2 - t_0} - 1 \right] \text{Im} F(s', t_0) ds'; \ell=2, 4, \dots \quad (2.2)$$

is then equal to $\frac{\sqrt{s}}{k} f_{\ell}(s)$. The D wave scattering length $\alpha_2^t = \lim_{k \rightarrow 0} \frac{f_2(s)}{k^5}$ and from (2.2) it is also given by

$$\alpha_2^t = \frac{8}{15\pi\mu} \int_{4\mu^2}^{\infty} \frac{\text{Im} F(s', t_0) ds'}{s'^3} \quad (2.3)$$

We wish to find the maximum of

$$\int_{4\mu^2}^{\infty} (s' - 4\mu^2)^{m-\frac{3}{2}} |F[s', (2\mu^2 - s')/2, (1 - \cos\theta)]| ds'$$

for $m > \frac{1}{4}$ under the condition (2.2) with $\ell = 2$ or (2.3). To do this we first bound $|F(s, t)|$ by $\hat{F}(s, t)$ which is defined by the relations *)

$$\begin{aligned} |F(s', t)| &\leq \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{\infty} (2\ell+1) |f_{\ell}(s')| |P_{\ell}(\cos\theta)| \\ &\leq \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{\infty} (2\ell+1) |f_{\ell}(s')| B_{\ell}(\cos\theta) \equiv \hat{F}(s, t) \end{aligned} \quad (2.4)$$

where $B_{\ell}(\cos\theta)$ is a function such that $|P_{\ell}(\cos\theta)| \leq B_{\ell}(\cos\theta)$ for all even ℓ and which decreases with ℓ . We take

$$B_{\ell}(\cos\theta) = \sqrt{\frac{2}{\pi \ell |\sin\theta|}} \quad \ell = 2, 4, \dots; \quad B_0 = \sqrt{\frac{1}{|\sin\theta|}} \quad (2.5)$$

Now from unitarity

$$\text{Im } F(s', t_0) \geq \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{\infty} (2\ell+1) |f_{\ell}(s')| P_{\ell}\left[1 + \frac{2t_0}{s' - 4\mu^2}\right] \equiv G(s', t_0) \quad (2.6)$$

where this relation defines $G(s', t_0)$. Of great importance to our work is the fact that when $\hat{F}(s, t)$ has a given value, $G(s', t_0)$ has a well defined minimum. This fact is an immediate generalization of the contents of Ref. 4) **) to the case when $\theta \neq 0$. The distribution of partial wave amplitudes which gives this minimum value for $G(s, t)$ has moduli given by

*) Unless otherwise stated it will be taken as understood that all sums are over even ℓ .

**) We would like to thank Dr. A. Martin for informing us of this generalization.

$$|f_\ell(s')| = 1, \quad \ell = 0, 2, \dots, 2L$$

$$|f_\ell(s')| = \frac{c B_\ell(\omega\theta)}{P_\ell(x)}, \quad \ell = 2L+2, 2L+4, \dots \quad (2.7)$$

where c is independent of ℓ and $x = 1+2t_0/(s'-4\mu^2)$. L is defined by the inequalities

$$\frac{P_{2L}(x)}{B_{2L}(\omega\theta)} < c \leq \frac{P_{2L+2}(x)}{B_{2L+2}(x)} \quad (2.8)$$

It may be proved that L and c are uniquely defined when $\hat{F}(s', t)$ is given.

Let $\Phi[\hat{F}, s', t_0]$ be this minimum value of $G(s', t_0)$.

Then

$$\Phi = \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{2L} (2\ell+1) P_\ell(x) + \frac{\sqrt{s'}}{k'} c^2 \sum_{\ell=2L+2}^{\infty} (2\ell+1) \frac{[B_\ell(\omega\theta)]^2}{P_\ell(x)} \quad (2.9)$$

where

$$c = \left\{ \hat{F}(s', t) - \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{2L} (2\ell+1) B_\ell(\omega\theta) \right\} / \left\{ \frac{\sqrt{s'}}{k'} \sum_{\ell=2L+2}^{\infty} \frac{(2\ell+1) [B_\ell(\omega\theta)]^2}{P_\ell(x)} \right\} \quad (2.10)$$

From (2.2)

$$\frac{\pi}{4} (4\mu^2 - t_0) a_2(t_0) \geq \int_{4\mu^2}^s Q_2 \left[\frac{2s'}{4\mu^2 - t_0} - 1 \right] \Phi[\hat{F}, s', t_0] ds' \quad (2.11)$$

It is easy to see from (2.9) and (2.10) that Φ increases with $\hat{F}$ so that $\hat{F}$ cannot be made too large otherwise Φ becomes too large and violates (2.11). By the method of Lagrange multipliers, the maximum value of $I(s, \cos\theta)$ defined by

$$I(s, \cos\theta) \equiv \int_{4\mu^2}^s (s' - 4\mu^2)^{m-3/4} \hat{F}(s', t) ds' \quad (2.12)$$

under the constraint (2.11) occurs when

$$-\lambda Q_2 \left[\frac{2s'}{4\mu^2 - t_0} - 1 \right] \frac{\partial \Phi}{\partial \hat{F}} + (s' - 4\mu^2)^{m-3/4} = 0 \quad (2.13)$$

where λ is independent of s' but depends on s . From (2.9), (2.10) and (2.13),

$$C = \frac{(s' - 4\mu^2)^{m-3/4}}{2\lambda Q_2 \left[\frac{2s'}{4\mu^2 - t_0} - 1 \right]} \quad (2.14)$$

Substituting in (2.9) and (2.10)

$$\Phi(\hat{F}, s', t) = \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{2L} (2\ell+1) P_{\ell}(x) + \frac{\sqrt{s'}(s' - 4\mu^2)^{(2m-3/2)}}{k' [2\lambda Q_2 \left(\frac{2s'}{4\mu^2 - t_0} - 1 \right)]^2} \sum_{\ell=2L+2}^{\infty} (2\ell+1) \frac{[B_{\ell}(\cos\theta)]^2}{P_{\ell}(x)} \quad (2.15)$$

$$\hat{F}(s', t) = \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{2L} (2\ell+1) B_{\ell}(\cos\theta) + \frac{\sqrt{s'}(s' - 4\mu^2)^{(m-3/4)}}{k' [2\lambda Q_2 \left(\frac{2s'}{4\mu^2 - t_0} - 1 \right)]} \sum_{\ell=2L+2}^{\infty} (2\ell+1) \frac{B_{\ell}(\cos\theta)}{P_{\ell}(x)} \quad (2.16)$$

No explicit formula can be derived for the maximum of $I(s, \cos\theta)$ and it has to be calculated numerically. The method is to find the value of λ which when substituted in (2.15) gives equality in (2.11) ^{*}). This value of λ is unique and gives the maximum value of $I(s, \cos\theta)$ from (2.16) and hence an upper bound to $\bar{F}[s, (2\mu^2 - s/2)(1 - \cos\theta)]$.

Similarly if the D wave scattering length α_2^t is given, then instead of (2.11) one uses the inequality

$$\frac{15\pi\mu}{8} \alpha_2^t \geq \int_{4\mu^2}^s \frac{1}{s'^3} \Phi[\hat{F}, s', t_0] ds' \quad (2.17)$$

In the case of forward scattering we can likewise obtain a bound to $\bar{F}(s, 0)$, the only difference is that the B_ℓ 's are all replaced by unity. Finally to get a bound to $\bar{\sigma}_{\text{tot}}$ we use the optical theorem ^{**})

$$\sigma_{\text{TOT}}(s) \leq \frac{8\pi}{k\sqrt{s}} \text{Im} F(s, 0) \quad (2.18)$$

Then

$$\begin{aligned} \overline{\sigma_{\text{TOT}}}(s) &\equiv \frac{(m+1)}{(s-4\mu^2)^{m+1}} \int_{4\mu^2}^s (s'-4\mu^2)^m \sigma_{\text{TOT}}(s') ds' \\ &\leq \frac{32\pi(m+1)}{(s-4\mu^2)^{m+1}} \int_{4\mu^2}^s \frac{k'}{\sqrt{s'}} (s'-4\mu^2)^{m-1} \text{Im} F(s', 0) ds' \leq \frac{32\pi(m+1)}{(s-4\mu^2)^{m+1}} \int_{4\mu^2}^s \frac{k'}{\sqrt{s'}} (s'-4\mu^2)^{m-1} |F(s', 0)| ds' \end{aligned} \quad (2.19)$$

^{*}) We can obviously replace $a_2(t_0)$ by its upper bound in (2.11).

^{**}) There is an extra factor 2 on the right-hand side due to the fact that we are dealing with the scattering of identical particles.

A bound of the right-hand side can then be obtained in the manner described above. In Section 4 we will present some results on the numerical calculation of bounds to $\overline{\sigma}_{\text{tot}}(s)$ at a selection of energies.

3. - ASYMPTOTIC BOUNDS

Since our technique only gives numerical bounds to $\bar{F}(s,t)$ etc., and not a general formula for them, it may seem a little surprising that we can give an asymptotic bound for these quantities as $s \rightarrow \infty$. However, we will now proceed to do this and derive bounds which are equivalent to the Froissart bounds.

We consider first of all the fixed angle scattering amplitude and investigate the behaviour of $I(s, \cos\theta)$ for large s .

The starting point is the fact that L defined by (2.8) is a function of both s and s' since c is also a function of these two variables from (2.14). Also from (2.8) and (2.14)

$$\frac{P_{2L}(x)}{B_{2L}(\cos\theta)} < \frac{(s'-4\mu^2)^{m-3/4}}{2\lambda Q_2 \left[\frac{2s'}{4\mu^2 - t_0} - 1 \right]} \leq \frac{P_{2L+2}(x)}{B_{2L+2}(\cos\theta)} \quad (3.1)$$

and it is easily proved from these inequalities that for fixed s , L increases with s' . Therefore writing in the dependence of L on s and s' ,

$$L(s, s') \leq L(s, s) \equiv L_0 \quad (3.2)$$

where this defines L_0 .

The next step is to obtain an asymptotic bound for L_0 . We start from the following bounds for the Legendre polynomials which hold for $x \gg 1$ *)

$$M_n \left[x + \frac{n}{n+1} (x^2-1)^{1/2} \right]^{2L} \leq P_{2L}(x) \leq \left[x + (x^2-1)^{1/2} \right]^{2L}, \quad n=1,2,\dots \quad (3.3)$$

where M_n is defined in (1.2). Using these bounds in (3.1),

$$\frac{M_n \left[x + \frac{n}{n+1} (x^2-1)^{1/2} \right]^{2L}}{B_{2L}(\omega\theta)} \leq \frac{(s'-4\mu^2)^{m-3/4}}{2\lambda Q_2 \left[\frac{2s'}{4\mu^2-t_0} - 1 \right]} \leq \frac{\left[x + (x^2-1)^{1/2} \right]^{2L+2}}{B_{2L+2}(\omega\theta)} \quad (3.4)$$

Then from (2.11), (2.15) and (3.4),

$$\begin{aligned} & \left\{ \frac{Q_2 \left[\frac{2s}{4\mu^2-t_0} - 1 \right]}{B_{2L_0}(\omega\theta) (s-4\mu^2)^{m-3/4}} \right\}^2 M_n^2 \left[X + \frac{n}{n+1} (x^2-1)^{1/2} \right]^{4L_0} x \\ & \times \int_{4\mu^2}^{\infty} \left\{ (s'-4\mu^2)^{2m+3/2} Q_2 \left[\frac{2s'}{4\mu^2-t_0} - 1 \right] \right\}^{-1} \frac{\sqrt{s'}}{k'} \sum_{\ell=2L+2}^{\infty} \frac{(2\ell+1) [B_{\ell}(\omega\theta)]^2 ds'}{P_{\ell}(x)} \\ & \leq \frac{\pi}{4} (4\mu^2-t_0) a_2(t_0) \end{aligned} \quad (3.5)$$

where $X = 1 + 2t_0/(s-4\mu^2)$. Using (3.2) and substituting for the B_{ℓ} we get

$$\begin{aligned} & 4 M_n^2 L_0 \left\{ \frac{Q_2 \left(\frac{2s}{4\mu^2-t_0} - 1 \right)}{(s-4\mu^2)^{m-3/4}} \right\}^2 \left[X + \frac{n}{n+1} \sqrt{x^2-1} \right]^{4L_0} \int_{\alpha s}^s \frac{\left[x + (x^2-1)^{1/2} \right] (s'-4\mu^2)^{2m-3/2} \left\{ \left[x + (x^2-1)^{1/2} \right] - 1 \right\}^{-1} ds'}{Q_2 \left[\frac{2s'}{4\mu^2-t_0} - 1 \right]} \\ & \leq \frac{\pi}{4} (4\mu^2-t_0) a_2(t_0) \end{aligned} \quad (3.6)$$

*) These bounds are due to A. Martin. It should be noted that the lower bounds are saturated as $x \rightarrow \infty$ when $n = L$.

where $0 < \alpha < 1$. Then if $X_\alpha = 1 + 2t_0/(\alpha s - 4\mu^2)$, we find that as $s \rightarrow \infty$,

$$L_0 \leq \frac{3}{8} \sqrt{\frac{s}{t_0}} \left\{ \log \left[s / (s_2 L_0^{2/3}) \right] \right\} / \left[\frac{2n}{n+1} - \frac{1}{\alpha} \right] + O(1) \quad (3.7)$$

where

$$s_2^{3/2} = \frac{2M_n^2 (4\mu^2 - t_0)^2 (1 - \alpha^{2m+3})}{15\pi a_2(t_0) \sqrt{t_0} (2m+3)} \quad (3.8)$$

We now take $n = s^\epsilon$ and $\alpha = \{1 - (2m+3)s^{-\epsilon}\}^{1/(2m+3)} \simeq 1 - s^{-\epsilon}$, choosing ϵ so that $s^\epsilon \rightarrow \infty$ as $s \rightarrow \infty$ but $\epsilon \rightarrow 0$ as $s \rightarrow \infty$, e.g., we can put $\epsilon = [\log(s/\mu^2)]^{-\beta}$ with β a constant such that $0 < \beta < 1$.

Then as $s \rightarrow \infty$

$$L_0 \leq \frac{3}{8} \sqrt{\frac{s}{t_0}} \log \left[s / (s_1 L_0^{2/3}) \right] + O(s^{1/2-\epsilon}) \quad (3.9)$$

where s_1 is defined in (1.11). Finally from (3.9) we get that asymptotically

$$L_0 \leq \frac{1}{4} \sqrt{\frac{s}{t_0}} \log \left[s / s_0 \right] \quad (3.10)$$

with s_0 defined by (1.10).

From (2.12) and (2.16)

$$I(s, \cos \theta) \leq \int_{4\mu^2}^s (s' - 4\mu^2)^{m-3/4} \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{2L} (2\ell+1) B_{\ell}(\cos \theta) ds' +$$

$$+ \int_{4\mu^2}^s \left\{ \frac{(s' - 4\mu^2)^{2m-3/2}}{2\lambda Q_2 \left[\frac{2s'}{4\mu^2 - t_0} - 1 \right]} \frac{\sqrt{s'}}{k'} \sum_{\ell=2L+2}^{\infty} \frac{(2\ell+1) [B_{\ell}(\cos \theta)]^2}{P_{\ell}(x)} \right\} ds' \quad (3.11)$$

Let I_1 and I_2 be the first and second integrals respectively. Then from (3.9) and (3.11),

$$I_1 \leq \int_{4\mu^2}^s (s' - 4\mu^2)^{m-3/4} \frac{\sqrt{s'}}{k'} \sum_{\ell=0}^{2L_0} (2\ell+1) B_{\ell}(\cos \theta) ds' \leq \frac{16 s^{m+1}}{3\sqrt{\pi} \sin \theta (m+1/4) \left(\frac{1}{4\mu^2 - t_0} \right) \left(\frac{s}{s_0} \right)^{3/2}} \quad (3.12)$$

We will prove in the Appendix that as $s \rightarrow \infty$,

$$I_2 = O \left\{ s^{m+1} [\log(s/s_0)]^{1/2} \right\} \quad (3.13)$$

The bound for the averaged fixed angle amplitude is then for $\theta \neq 0$ and $s \rightarrow \infty$,

$$\bar{F} [s, (2\mu^2 - s/2)(1 - \cos \theta)] \leq \frac{(m+1)}{(s - 4\mu^2)^{m+1/4}} \int_{4\mu^2}^s (s' - 4\mu^2)^{m-3/4} \hat{F}(s', t) ds'$$

$$\leq \frac{2}{3\sqrt{\pi} \sin \theta} \left(\frac{s}{t_0} \right)^{3/4} \left\{ \log(s/s_0) \right\}^{3/2} \left[\frac{m+1}{m+1/4} \right] \quad (3.14)$$

which is the bound given in Section 1. In exactly the same way the asymptotic bounds for the averaged fixed t , forward amplitudes and total cross-section may be shown to be given by (1.13), (1.14) and (1.15) respectively.

4. - NUMERICAL RESULTS AND CONCLUSIONS

To make numerical estimates of our bounds to $\overline{\sigma}_{\text{tot}}(s)$, etc., we need to have a bound to $a_2(t_0)$ for some $0 < t_0 < 4\mu^2$ or a value for the D wave scattering length α_2^t . For $\pi^0 \pi^0 \rightarrow \pi^0 \pi^0$ scattering a bound for $a_2(2\mu^2)$ can be obtained from the known bounds

$$|F(2\mu^2, 2\mu^2)| < 37, \quad |F(2\mu^2, 3\mu^2)| < 150 \quad (4.1)$$

which have been derived from our basic assumptions of unitarity, crossing, etc. ⁴⁾. Since the Froissart-Gribov formula implies all the partial waves $a_\ell(2\mu^2)$ are positive for $\ell \geq 2$, ⁷⁾,

$$0 \leq a_2(2\mu^2) \leq \frac{F(2\mu^2, 3\mu^2) - F(2\mu^2, 2\mu^2)}{5[P_2(2) - P_2(1)]} \leq 8.31 \quad (4.2)$$

This bound has been used to calculate the bound to $\overline{\sigma}_{\text{tot}}(s)$ for $m = 1$ and for a range of momenta between 1.0 and 13.0 GeV/c. The results are given in column two of the Table and can be compared in column three with the asymptotic bound for $\overline{\sigma}_{\text{tot}}(s)$ given by (1.15) for the above value of m .

To test our bound when the D wave scattering length is used we take a value for α_2^t which has been evaluated from forward dispersion relations ⁸⁾. In units where $\mu = 1$, $\alpha_2^t = 16 \times 10^{-5}$. The calculated bound for $\overline{\sigma}_{\text{tot}}(s)$ is given in column four and the corresponding asymptotic bound is given in column five of the Table. Once again we have taken $m = 1$. Finally we compare these results with the bounds of Yndurain given by (1.1) which are given in column six and the corresponding asymptotic bounds obtained by taking the leading term in (1.1) which are given in column seven.

It can be seen that our bounds are better by roughly a factor five than those given in Ref. 3) when α_2^t has the value given above. If α_2^t is increased by a factor ten then our bounds are again better but now only by a factor two or three. It can be also seen that although the bound (1.1) is already very close to the asymptotic form for the energies considered, our bounds do not tend to the asymptotic bound which we have derived for them.

An estimate of $\pi\pi$ scattering cross-sections may be made from experimental information on pion-nucleon and nucleon-nucleon scattering using either the quark model or factorization⁹⁾. These estimates are of $\sim 16 \mu\text{b}$ in our momentum range so that even our best bound for $\overline{\sigma}_{\text{tot}}(s)$ from the value of α_2^t used above is ~ 200 times too large. The bounds obtained from our bound to $a_2(2\mu^2)$ which are completely rigorous since they require no input information beyond the given assumptions of crossing, unitarity and analyticity are, as one might expect worse than these derived from α_2^t , being in fact 10^3 times larger than the expected values.

This large discrepancy between the expected value for $\overline{\sigma}_{\text{tot}}$ and our bounds can be attributed partly to the fact that as explained in Ref. 3), we have taken $\pi\pi$ dynamics to be governed by 2π exchange whereas experimentally it seems to be governed by ρ exchange. Another reason could be the fact we have used formula (2.3) for the D wave $\pi\pi$ scattering length, but this only requires that $\text{Im}F(s, 4\mu^2)$ roughly speaking is of order $s^{2-\epsilon}$ as $s \rightarrow \infty$ with $\epsilon > 0$. However, it appears to be the case that $\text{Im}F(s, 4\mu^2)$ is of order s^* , so that we are not making full use of this behaviour in our work.

*) This statement is based on the assumption that the Pomeron trajectory is almost flat.

As mentioned earlier, Singh and Roy ⁵⁾ have already derived asymptotic bounds for the total cross-section, fixed angle amplitude, etc., using as we have done the techniques of Ref. 4). For example they find that

$$\sigma_{\text{TOT}}(s) \lesssim \frac{\pi}{\mu^2} \left\{ \log \left[s / (\mu^4 \sigma_{\text{el}}) \right] \right\}^2 \left[\frac{\sigma_{\text{el}}}{\sigma_{\text{TOT}}} \right] \quad (4.3)$$

and

$$|F(s,0)| \lesssim \frac{s \sqrt{\sigma_{\text{el}}}}{8\mu \sqrt{\pi}} \left\{ \log \left[s / (\mu^4 \sigma_{\text{el}}) \right] \right\} \quad (4.4)$$

These bounds depend on the elastic cross-section σ_{el} as do those for the fixed angle and fixed t amplitudes.

Replacing σ_{el} by its upper bound σ_{TOT} we get from (4.3) that,

$$\sigma_{\text{TOT}}(s) \lesssim \frac{\pi}{\mu^2} \log \left[s / (\mu^4 \sigma_{\text{TOT}}) \right] \quad (4.5)$$

We can "iterate" this inequality to obtain ^{*)}, in much the same way as we derived the bound (3.10) for L_0 from (3.9), the bound

$$\sigma_{\text{TOT}}(s) \lesssim \frac{\pi}{\mu^2} \log (s/s_4) \quad (4.6)$$

where

$$s_4 = \pi \mu^2 \log \left\{ s / [\pi \mu^2 \log (s/\mu^2)] \right\} \quad (4.7)$$

^{*)} To do this rigorously one needs information on the lower order terms in (4.5).

Comparing (4.6) with (1.15) we see that the bounds are equivalent as m becomes very large and the respectively scaling factors s_0 and s_4 have the same dependence on s except for the difference of certain constants.

Similarly for the forward scattering amplitude from (4.4) and (4.5),

$$|F(s,0)| \lesssim \frac{s}{8\mu\sqrt{\pi}} \sqrt{\sigma_{\text{tot}}(s)} \log [s/(4\mu^4\sigma_{\text{tot}})] \quad (4.8)$$

Using (4.6),

$$|F(s,0)| \lesssim \frac{s}{8\mu^2} \left\{ \log (s/s_4) \right\} \log \left\{ s/[\pi\mu^2 \log (s/s_4)] \right\} \quad (4.9)$$

This bound is almost equivalent to our bound (1.14) for the average forward amplitude when $t_0 = 4\mu^2$ except here the scaling factors in the two logarithms are now different and there is an over-all factor 2 in (4.9). This latter factor comes from the fact that we have used the symmetry of the $\pi^0\pi^0$ state to put all odd waves to zero, while Roy and Singh have considered the general case when the odd waves have to be included.

For the fixed angle and fixed t amplitudes our bounds have the same form as those given in reference in the sense described above, but are worse by an over-all scaling factor $\frac{3}{2}$, when due allowance is made for the fact that we have considered a symmetric $\pi\pi$ state.

To summarize briefly the results of our work, we can say that we have improved on the technique of Yndurain ³⁾ for obtaining rigorous bounds to the averaged total cross-sections over all energy ranges and extended the method to cover averages of the fixed angle, fixed t and forward amplitudes.

The asymptotic form of the bounds has been found and we have shown that they are closely related to the most recent form of the Froissart bounds ⁵⁾. This last fact is not too surprising since the latter bounds were also obtained by using the methods of Ref. 4).

ACKNOWLEDGEMENTS

We would like to thank Dr. A. Martin for many useful and stimulating discussions. Thanks are also due to Dr. F.J. Yndurain for a useful correspondence and in particular for informing us of the most recent form of his work.

APPENDIX

We prove in this Appendix that

$$I_2 \equiv \int_{4\mu^2}^s (s' - 4\mu^2)^{2m-3/2} \left\{ \frac{1}{2\lambda Q\left[\frac{2s'}{4\mu^2-t_0}-1\right]} \frac{\sqrt{s'}}{R'} \sum_{\ell=2L+2}^{\infty} \frac{(2\ell+1) [B_\ell(\omega\theta)]^2}{P_\ell(x)} \right\} ds' \quad (A.1)$$

$$= O \left\{ s^{m+1} [\log(s/s_0)]^{1/2} \right\} \quad \text{as } s \rightarrow \infty$$

Substituting a lower bound for λ from (3.1),

$$I_2 \leq \sqrt{\frac{\pi \sin \theta}{a}} s^{m-3/4} \int_{4\mu^2}^s \sqrt{2L+2} P_{2L+2}(x) \left\{ \sum_{\ell=2L+2}^{\infty} \frac{(2\ell+1) [B_\ell(\omega\theta)]^2}{P_\ell(x)} \right\} ds' \quad (A.2)$$

Then using the bounds (3.3) for $P_\ell(x)$ and the fact that $L \leq L_0$,

$$I_2 \leq c_1 s^{m-3/4} \int_{4\mu^2}^s \frac{\sqrt{L_0}}{M_n} \left[\frac{x+(x^2-1)^{1/2}}{x+\frac{n}{n+1}(x^2-1)^{1/2}} \right]^{2L_0+2} \left\{ \sum_{\ell=0}^{\infty} \left[x+\frac{n}{n+1}(x^2-1)^{1/2} \right]^{-\ell} \right\} ds' \quad (A.3)$$

where c_1 is a constant. As $s \rightarrow \infty$,

$$I_2 \lesssim c_1 s^{m-3/4} \int_{4\mu^2}^s \frac{\sqrt{L_1}}{M_n} \left[\frac{x+(x^2-1)^{1/2}}{x+\frac{n}{n+1}(x^2-1)^{1/2}} \right]^{2L_1+2} \sqrt{\frac{s'}{4t_0}} ds' \quad (A.4)$$

where $L_1 = \frac{1}{4} \sqrt{\frac{s}{t_0}} \log(s/s_0)$ is the asymptotic upper bound to L_0 .

Now

$$\left[\frac{x + (x^2 - 1)^{1/2}}{x + \frac{n}{n+1} (x^2 - 1)^{1/2}} \right]^{2L_1 + 2} \leq \left[\frac{1}{1 - \frac{1}{n+1} (1 - 1/x^2)^{1/2} - \frac{n}{n+1} (1 - 1/x^2)} \right]^{2L_1 + 2} \quad (A.5)$$

choose n so that

$$\frac{1}{n+1} (1 - 1/x^2)^{1/2} + \frac{n}{n+1} (1 - 1/x^2) = \frac{1}{2L_1 + 2} \quad (A.6)$$

and then the right-hand side of (A.5) $\rightarrow e$ as $L_1 \rightarrow \infty$ with s .

For n as given by (A.6) to be ≥ 1 , it is necessary and sufficient that

$$x^2 \geq (x^2 - 1) (2L_1 + 2) \quad (A.7)$$

i.e.,
$$s' \geq 4\mu^2 + 2t_0 / \left\{ \left[\frac{2L_1 + 2}{2L_1 + 1} \right]^{1/2} - 1 \right\}$$

Defining this lower bound to be s_3 , we split the bound to I_2 given by (A.4) into two integrals in the following way

$$\begin{aligned} I_2 \leq & c_1 s^{m-3/4} \int_{4\mu^2}^{s_3} \frac{\sqrt{L_1}}{M_n} \left[\frac{x + (x^2 - 1)^{1/2}}{x + \frac{n}{n+1} (x^2 - 1)^{1/2}} \right]^{2L_1 + 2} \sqrt{\frac{s'}{4t_0}} ds' + \\ & + c_1 s^{m-3/4} \int_{s_3}^s \frac{\sqrt{L_1}}{M_n} \left[\frac{x + (x^2 - 1)^{1/2}}{x + \frac{n}{n+1} (x^2 - 1)^{1/2}} \right]^{2L_1 + 2} \sqrt{\frac{s'}{4t_0}} ds' \quad (A.8) \end{aligned}$$

Let J_1 and J_2 be the two integrals on the right-hand side of (A.8). Then since $M_n < c_2 n^{-\frac{1}{2}}$ with c_2 a constant,

$$J_2 \lesssim \frac{c_1 s^{m-3/4}}{\sqrt{4t_0} c_2} \int_{s_3}^s \frac{\sqrt{L_1(2L_1+2)} (1-1/x^2)^{1/4} s'^{1/2}}{\{1 - (2L_1+2)(1-1/x^2)\}^{1/2}} ds' \quad (A.9)$$

$$\lesssim c_3 s^{m+1/2} L_1 = c_4 s^{m+1} \log(s/s_0)$$

where c_3 and c_4 are independent of s .

Similarly since $s_3 \sim 4\mu^2 + 4t_0(2L_1+1)$ as $s \rightarrow \infty$,

$$J_1 \lesssim c_5 s^{m-3/4} \int_{4\mu^2}^{[4\mu^2+4t_0(2L_1+1)]} n^{1/2} L_1^{1/2} s'^{1/2} \left[\frac{2}{2-\frac{1}{n+1}} \right]^{2L_1+2} ds' \quad (A.10)$$

Taking $n = L_1$ ^{*)},

$$J_1 \lesssim c_6 L_1^{5/2} s^{m-3/4} = O\{s^{m+1/2} [\log(s/s_0)]^{5/2}\} \quad (A.11)$$

*) It is permissible to take n to be different in the two integrals J_1 and J_2 , since we could first of all split the integration range in the expression for the bound to I_2 given by (A.2) into two parts divided by the point $s' = s_3$, before using the bounds (3.3) for $P_\ell(x)$.

where c_5 and c_6 are constants. Comparing (A.9) with (A.11) we see that J_2 is the leading term in I_2 so that

$$I_2 = O[s^{m+1} \log(s/s_0)] \quad (\text{A.12})$$

k (GeV/c)	Bounds to $\sigma_{\text{tot}}(s)$ from the bound to $a_2(2\mu^2)$		Bounds to $\sigma_{\text{tot}}(s)$ from the bound to α_2^t		Yndurain's bounds to $\sigma_{\text{tot}}(s)$	
	Actual (μb)	Asymptotic (μb)	Actual (μb)	Asymptotic (μb)	Actual (μb)	Asymptotic (μb)
1.0	1.56 E + 04	2.62 E + 04	7.18 E + 02	9.82 E + 02	4.28 E + 03	4.07 E + 03
3.0	2.18 E + 04	3.78 E + 04	1.60 E + 03	2.48 E + 03	7.99 E + 03	7.88 E + 03
5.0	2.52 E + 04	4.38 E + 04	2.20 E + 03	3.50 E + 03	1.02 E + 04	1.01 E + 04
7.0	2.76 E + 04	4.82 E + 04	2.66 E + 03	4.28 E + 03	1.18 E + 04	1.18 E + 04
9.0	2.95 E + 04	5.16 E + 04	3.04 E + 03	4.94 E + 03	1.31 E + 04	1.31 E + 04
11.0	3.11 E + 04	5.44 E + 04	3.36 E + 03	5.52 E + 03	1.42 E + 04	1.42 E + 04
13.0	3.24 E + 04	5.68 E + 04	3.66 E + 03	6.00 E + 03	1.52 E + 04	1.51 E + 04

REFERENCES

- 1) M. Froissart - Phys.Rev. 123, 1053 (1961).
- 2) A. Martin - Phys.Rev. 129, 1432 (1963) ;
Y.S. Jin and A. Martin - Phys.Rev. 153, B1375 (1964).
- 3) F.J. Yndurain - "Absolute Bounds on Cross-Sections at All Energies and Without Unknown Constants". To be published in Phys. Letters.

The bound quoted here in (1.1) is actually an improvement on the bound given in the original version of the above paper. We would like to thank Dr. F.J. Yndurain for informing us of his new bound.
- 4) L. Lukaszuk and A. Martin - Nuovo Cimento 52A, 122 (1967).
- 5) V. Singh and S.M. Roy - "Upper Bounds on the Elastic Differential Cross-Sections". Preprint of the Tata Institute of Fundamental Research, Bombay, India.
- 6) J.L. Gervais and F.J. Yndurain - Phys.Rev. 167, 1289 and 169, 1187 (1968).
- 7) A. Martin - Nuovo Cimento 47, 265 (1967).
- 8) D. Morgan and G. Shaw - Nuclear Phys. B10, 261 (1969).

Almost the same value for α_2^t has been obtained by J.C. Le Guillou, A. Morel and H. Navelet using a model for low energy $\pi\pi$ scattering incorporating unitarity and crossing. Private communication.
- 9) For a comparison of results from these techniques and further references, see : H.J. Lipkin - "Quarks and Symmetries", p. 253, Proceedings of the Heidelberg International Conference on Elementary Particles (1967).