CM-P00058646

Ref.TH.1215-CERN

A GENERAL APPROACH TO DUAL MULTILoop DIAGRAMS

V. Alessandrini *)

CERN - Geneva

A B S T R A C T

The general solution of the Fairlie-Nielsen analogue model for dual multiparticle theory is discussed. By using the theory of functionals on Riemann surfaces, it is shown that all the relevant Green functions for orientable and non-orientable multiloop diagrams can be obtained in terms of Burnside automorphic functions. As an example, the one-loop orientable and non-orientable diagrams are considered in detail from this point of view. A general discussion of N loop orientable and non-orientable diagrams in the operator formalism is also given.

*) On leave of absence from University of La Plata, Argentina.

1. INTRODUCTION

It was conjectured by Nielsen some time ago ¹⁾ that the calculation of the integrand of an N loop Veneziano formula ²⁾ is equivalent to the solution of a harmonic problem on a suitably chosen open Riemann surface. The duality diagram is made to correspond to a two-dimensional conducting domain by identifying the quark lines with its boundaries, and the external lines with electrodes carrying a current equal to the momentum of the corresponding particle. Then the integrand of the generalized Veneziano formula is given by the exponential of the heat generated by the steady current on the surface, assuming a constant resistivity. This conjecture was originally proved only in the case of zero and one orientable loop diagrams ¹⁾. An m loop diagram will give a $(m+1)$ -ply connected domain.

This work was initially motivated by an attempt to understand the one-loop non-orientable diagrams from the point of view of the analogue model. This requires the solution of a harmonic problem on a Möbius strip, and, to the best of my knowledge, there does not exist in the literature an explicit discussion of this problem. The investigation and solution of the one-loop non-orientable case turned out to be a profitable exercise: it gave the clue for the general solution of the analogue model; and suggested that the role played by elliptic functions in the one-loop case will be played by automorphic functions in the multi-loop case.

Nielsen's conjecture can be put in a neat mathematical form ³⁾ by stating that the Veneziano integrand for the duality diagram B is given by

$$e^{-\frac{\alpha'}{2} \sum_{i \neq j} p_i \cdot p_j N_B(z_i, z_j)} \quad (1.1)$$

where α' is the slope of the input Regge trajectories and $N_B(z_i, z_j)$ is the Neumann function taken at the points z_i, z_j on the boundary of B where the external lines with momenta p_i, p_j are attached. By the Neumann function we mean the usual Green function on the surface with the boundary condition that its normal derivative has a constant value on the boundaries. The reason why the Neumann function occurs in Eq. (1.1) is because the analogue model represents a boundary value problem of the second kind: the current - that is to say, the normal derivative of the potential - is given on the boundary. Therefore, from Eq. (1.1) one can see that the main problem is to obtain the Neumann function corresponding to any orientable or non-orientable Riemann surface.

I start by discussing in Section 2 the basic elements of the theory of functionals on Riemann surfaces that are needed in the rest of the paper, following Schiffer and Spencer ^{4)*}. In particular, it is shown how one constructs Green and Neumann functions on a Riemann surface in terms of the so-called third Abelian integrals ⁴⁾. This method treats both orientable and non-orientable surfaces in a unified way. In Section 3 I point out how one is naturally led to a general solution in terms of automorphic functions ⁵⁾. Indeed, in 1891 Burnside studied a class of automorphic functions ⁶⁾ that are precisely the ones that solve Nielsen's analogue model. A short account of Burnside's result is given, and it is also discussed how they must be slightly extended in order to construct the third Abelian integrals for the problem at hand.

The next section (Section 4) is devoted to the discussion of the one-loop orientable and non-orientable diagrams. This section serves several purposes: all the mathematics of the previous sections is illustrated with an example, the solution of harmonic problems on a Möbius strip is given, all the previous results on one-loops are derived with almost no calculational effort (because they are special cases of the general solution) and the differences between orientable and non-orientable diagrams is understood from very simple topological arguments.

*) I am grateful to Prof. C. Lovelace for pointing out this reference to me.

An elegant proof of Nielsen's conjecture has been given by Lovelace ³⁾, by showing that the solution of the analogue model in terms of Burnside automorphic functions is obtained by starting from his simple N Reggeon vertex in the operator formalism and sewing M pairs of Reggeons to construct an M loop diagram. Needless to say, he also obtains the measure with which the Veneziano integrand must be integrated, missing in the analogue model. I include a final section with a discussion of his results tidying up various technical details, because they are essential for a complete clarification of this field.

2. FUNCTIONALS ON RIEMANN SURFACES

As explained in the Introduction, the content of this section is a collection of classical results from the theory of functionals on finite Riemann surfaces, extracted mainly from Chapter I-IV of Ref. 4). I am including them in order to make the paper to a large extent self-contained, with the conviction that this section will be appreciated by anyone who tries to make use of Ref. 4).

The characteristic property of a Riemann surface is the fact that to each point $\xi (\xi_1, \xi_2)$ on the surface one can associate a point $z(x,y)$ in the complex $z=x+iy$ plane in such a way that a neighbourhood of ξ is mapped conformally onto a complex neighbourhood of z . Of course, in general a global conformal mapping onto the complex plane is not required. The complex variable z corresponding to a neighbourhood of a point ξ on the surface is called a "local uniformizer". Moreover, if z is a uniformizer, any analytic function of z and $\bar{z}$ is also a uniformizer. In order to define functionals like, for example, the Green or the Neumann functions, on a given Riemann surface B , one proceeds in several steps. First, one constructs another Riemann surface D , called the "double" of B . Next, one defines certain functionals

on the double D , the so-called Abelian integrals. Finally, one defines the functionals on B in terms of the Abelian integrals on D . These steps are independent of whether the original surface B is orientable or non-orientable, and I proceed now to discuss them. The perceptive reader will notice the deep analogy of this method with the well-known "method of images" frequently used to solve simple boundary value problems.

A) Double of a Riemann Surface

The important remark here is that the original Riemann surface B can have any number of boundary curves, and may be orientable or non-orientable, but the double D is always closed and orientable, as it will be clear from the definition. R is defined by the following requirements.

- i) Two points of D are associated with, or lie over, each interior point of B . These two points of D , say ξ and $\tilde{\xi}$, that correspond to the same interior point of B , are called "conjugate points".
- ii) One point of D is associated with each boundary point of B . In other words: on the boundary of B , conjugate points of D coincide.
- iii) Finally, two disjoint neighbourhoods of D lie over each neighbourhood of an interior point of B ; i.e., the neighbourhoods of ξ and $\tilde{\xi}$. Suppose that z is a uniformizer of an interior point of B . Then the convention is made that only analytic functions of z are used as uniformizers in ξ , and analytic functions of $\bar{z}$ are used as uniformizers in $\tilde{\xi}$.

Let us consider some simple examples. If the original Riemann surface B is orientable, then the construction of D is easy. One imagines another identical surface $\bar{B}$, a mirror image of B , and constructs a closed surface by identifying corresponding boundary points of B and $\bar{B}$. The surface so constructed is D .

Conjugate points are corresponding points in B and $\bar{B}$. In this way it is easy to see that the double of a disk is a sphere; the double of an annulus is a torus (or sphere with one handle) and that the double of a plane domain with m boundary curves is a connected sum of $(m-1)$ tori, or sphere with $(m-1)$ handles.

The construction of the double of a non-orientable Riemann surface requires more imagination. An elementary theorem in the topology of surfaces states that any non-orientable surface B has a "two-sheeted orientable covering", to be denoted by B' . Here two-sheeted means that there are two points of B' corresponding to each point of B (both for interior and boundary points). Therefore, if B has n boundary curves, its two-sheeted covering B' has $2n$ boundary curves. Moreover, the fact that B is non-orientable means that there are closed paths on it along which the orientation is reversed. The two-sheeted covering has the basic property that such paths along which the orientation is reversed lead from one sheet of B' to the other. In this case the double D is constructed by starting from the two-sheeted orientable covering B' and "closing" it by identifying the corresponding boundary points. Clearly in this case conjugate points are just the corresponding points in the two sheets of B' .

As a (not so trivial) example, let us prove that the double of a Möbius strip is a torus. However, it is not quite the same torus as in the case of the annulus, there are some subtle points that make all the difference in solving a harmonic problem on an annulus or on a Möbius strip. The discussion that follows is crucial to the understanding of these differences. I start by considering the projective plane, because the Möbius strip is obtained by making a hole in it ⁴⁾. A model of the projective plane is obtained by considering a disk - or a hemisphere - and closing the surface by identifying diametrically opposite points of the boundary. It is easy to see that, because of this identification, the orientation of the surface is reversed as one moves along a diameter. Just move along a diameter a small circle rotating

anticlockwise and check that when it reaches the boundary it emerges at the opposite end of the diameter rotating in a clockwise direction. The two-sheeted orientable covering of the projective plane is the sphere ⁷⁾. Indeed, cutting the sphere along the equator and identifying antipodal points one recovers the projective plane. Moreover, a path on the projective plane along which the orientation is reversed leads from one point to its antipodal, that is to say, from one sheet to the other.

Let us now cut a hole in the projective plane. In the sphere model, we must discard both the arctic and antarctic regions because antipodal points are identified (see Fig. 1). What is left is a zone of the sphere bounded by two parallels, i.e., an annulus. Because of the identification of antipodes only half of the zone is needed, say, the "visible" half. The reader will immediately recognize that this half zone, with its ends AB identified as indicated in Fig. 1, is the Möbius strip. The whole annulus is the two-sheeted orientable covering of the Möbius strip. We see that it has two boundary curves, as expected, since the Möbius strip has only one boundary.

Finally, the double of the Möbius strip is obtained by identifying the two boundary curves, and then it is easily seen that one obtains again a torus. However, the two boundaries must be identified by identifying corresponding boundary points, and it is clear from Fig. 1 that one of the circles must be rotated (or "twisted") by π before the surface is closed and the torus is obtained. This fact must be kept in mind because it turns to be the only difference in the solution of harmonic problems on an annulus or on a Möbius strip.

Finally, we finish this discussion of the double D by remarking that, since D is always closed and orientable, a theorem in topology of surfaces guarantees that the most general double will be a sphere with a number of handles. However, the previous examples show that one must be very careful in each particular case with the identification of conjugate points.

B) Abelian Integrals

Having defined the double D of B , one considers next the differentials $dZ(\xi)$ defined on it; called Abelian differentials. An Abelian differential is said to be of the first kind if it is regular at every point of D ; of the second kind if it is regular except for a double pole, of the third kind if it is regular except for two single poles with opposite residues. From now on, we shall refer to them as first, second and third Abelian differentials. Its integrals will be called Abelian integrals, and in general they depend on the limits of the integral and the paths connecting them. Because the surface is multiply connected, they will be many-valued functions on R .

The Abelian integrals are uniquely defined by their periodicity properties along certain closed paths - or "cycles" - on the surface D . Two cycles that can be made to coincide by a continuous deformation of the path on the surface are said to be homologous. If a cycle can be shrunk to a point, it is said to be homologous to zero. Clearly, on the sphere, all cycles are homologous to zero. However, that is not the case on a sphere with M handles. In this case, there are $2M$ non-homologous cycles such that any other cycle non-homologous to zero is homologous to a linear combination of them. In a more sophisticated language, we can say that the homology group of the sphere with M handles has $2M$ generators. We refer the reader to Fig. 2, where the case $M = 2$ is illustrated.

Let K, L be two cycles in D . Then their Kronecker index is defined to be the sum of intersections, counting $(+1)$ for every point where L crosses K from right to left, and (-1) for every point where it crosses K from left to right. Clearly $I(K, L) = -I(L, K)$. A "canonical basis" on the sphere with M handles is a set of $2M$ cycles $K_{2\mu-1}, K_{2\mu}$ ($\mu = 1, \dots, M$) satisfying

$$I(K_{2\mu-1}, K_{2\mu}) = 1 \quad (2.1)$$

with all the other pairs non-intersecting. Figure 2 also exhibits the canonical basis for the case $M = 2$. The M cycles $K_{2\mu}$ are said to be "dual" to the M cycles $K_{2\mu-1}$ - although this is mathematical jargon, and has nothing to do with Veneziano duality (more like dual diagrams).

Now we are ready to consider the uniqueness of a first Abelian differential $d\omega$. Let us define its periods by

$$P_{\mu}(d\omega) = \oint_{K_{2\mu-1}} d\omega \quad (2.2)$$

$$Q_{\mu}(d\omega) = \oint_{K_{2\mu}} d\omega \quad \mu=1, \dots, M \quad (2.3)$$

Then it can be shown that $d\omega$ is determined uniquely either by specifying P_{μ} ($\mu = 1, \dots, M$) or by specifying $\text{Re}P_{\mu}$, $\text{Re}Q_{\mu}$, $\mu = 1, \dots, M$. This theorem means that there are only M independent first Abelian differentials on the sphere with M handles. Let us introduce M first Abelian differentials $d\omega_{\nu}$, ($\nu = 1, \dots, M$) defined by

$$P_{\mu}(d\omega_{\nu}) = \oint_{K_{2\mu-1}} d\omega_{\nu} = \delta_{\mu\nu} \quad \mu, \nu = 1, \dots, M \quad (2.4)$$

Then it is clear that any first Abelian differential $d\omega$ with arbitrary periods $P_{\mu}(d\omega)$ can be expressed as linear combinations of them. The set $d\omega_{\nu}$ is said to form a "complex basis" of first Abelian differentials. In general, they will also have periods around the remaining dual cycles $K_{2\mu}$. Let us define the period matrix

$$B_{\mu\nu} = 2\pi i \oint_{K_{2\mu}} d\omega_\nu = 2\pi i Q_\mu(d\omega_\nu) \quad (2.5)$$

We shall need later on the fact that $\text{Re} B_{\mu\nu}$ is a non-singular matrix ⁴⁾.

The basic functional that we shall need in what follows - besides the first Abelian differentials - is a third Abelian differential $d\Omega_{ab}(\xi)$ regular on D except for poles of residue ± 1 at $\xi = a$ or b , respectively. It is defined uniquely by requiring that the real part of its Abelian integral, $\Omega_{ab}(\xi)$, is single-valued in D . In other words, $d\Omega_{ab}(\xi)$ has pure imaginary periods when one circles around any of the $2M$ canonical cycles

$$\text{Re} \oint_{K_{2\mu-1}} d\Omega_{ab}(\xi) = 0 \quad (2.6)$$

$$\text{Re} \oint_{K_{2\mu}} d\Omega_{ab}(\xi) = 0 \quad (2.7)$$

We emphasize that the uniqueness criteria for third Abelian differentials are different from the ones corresponding to first Abelian differentials that we discussed before. When the third Abelian differential is defined in the way we have indicated, one speaks of the "real normalization".

There exists another third Abelian differential $d\omega_{ab}(\xi)$ with poles of residue ± 1 at $\xi = a, b$, respectively, defined by the requirement that its periods along the canonical cycles

$K_{2\mu-1}$ vanish

$$\oint_{K_{2\mu-1}} d\omega_{ab}(\xi) = 0 \quad (2.8)$$

This is the so-called "complex normalization". The two third Abelian integrals $\Omega_{ab}(\xi)$ and $\omega_{ab}(\xi)$ differ by a linear combination of first Abelian integrals. We shall discuss this point again later on. Either $\Omega_{ab}(\xi)$ or $\omega_{ab}(\xi)$ is unique up to a complex constant, for any closed orientable Riemann surface D .

C) Green and Neumann Functions

The Green and Neumann functions of the original duality diagram B are defined in terms of the third Abelian integral $\Omega_{ab}(\xi)$ of its double D . By Green function $G(\xi, \eta)$ of B we understand the harmonic function on B which vanishes on the boundary and has a logarithmic singularity at $\xi = \eta$, where it behaves like

$$G(\xi, \eta) = \log \left| \frac{1}{\xi - \eta} \right| + \text{regular terms} \quad (2.9)$$

Schiffer and Spencer show that

$$G(\xi, \eta) = \frac{1}{2} [\Omega_{\eta\bar{\eta}}(\xi) - \Omega_{\eta\bar{\eta}}(\bar{\xi})] \quad (2.10)$$

The similarity with the method of images is now apparent. One duplicates the surface and then introduces a point charge of opposite strength at $\bar{\eta}$, the conjugate point of η . The Neumann function is defined by

$$\begin{aligned} N(\xi, \eta, \eta_0) &= N(\xi, \eta) - N(\xi, \eta_0) = \\ &= \frac{1}{2} [\Omega_{\eta\eta_0}(\xi) + \Omega_{\bar{\eta}\bar{\eta}_0}(\bar{\xi}) + \Omega_{\eta\eta_0}(\bar{\xi}) + \Omega_{\bar{\eta}\bar{\eta}_0}(\xi)] \end{aligned} \quad (2.11)$$

Only the difference of Neumann functions is defined because $N(\xi, \eta)$ itself is not a conformal invariant. These definitions are valid both for orientable and non-orientable surfaces. For non-orientable surfaces, the only difficulty is that $G(\xi, \eta)$ is not single-valued (one must construct the "quadruple" of B if one wants a single-valued

Green function). However, this need not concern us because we only need the Neumann function, which is single-valued in any case.

Notice that in Eq. (1.1) we can replace $N_B(z_i, z_j)$ by $[N_B(z_i, z_j) - N_B(z_i, z_0)]$, z_0 being an arbitrary fixed point in the interior of the surface. Momentum conservation guarantees that the extra term contributes only an over-all constant to the Veneziano integrand. Then, recalling the fact that on the boundary conjugate points coincide, we get

$$N_B(z_i, z_j) - N_B(z_i, z_0) = \Omega_{z, z_0}(z_j) + \Omega_{z, \bar{z}_0}(z_j) = R \Omega_{z, z_0}(z_j) \quad (2.12)$$

which gives then the logarithm of the Veneziano integrand in terms of the third Abelian integral with real normalization of the double D .

3. GENERAL SOLUTION OF NIELSEN'S ANALOGUE MODEL

We have seen in the previous section that the main problem in solving a harmonic problem on a Riemann surface is to construct the Abelian integral $\Omega_{ab}(\mathfrak{z})$ of a sphere with M handles. Consider, for example, the case $M=1$, in which case we have seen that D is a single torus. In this case it is possible to establish a one-to-one correspondence between the universal covering surface of the torus and the complex plane, in such a way that single-valued functions on the torus become doubly periodic functions in the complex plane. This is easily understood if one notices that single-valued functions on a torus have to reproduce themselves as one goes along the two canonical cycles. This is the reason why elliptic functions, i.e., meromorphic doubly periodic functions, occur in one-loop calculations, as we shall see later on.

In the same way as meromorphic single-valued functions on a torus are elliptic functions on the complex plane, meromorphic single-valued functions on an arbitrary closed Riemann surface are automorphic functions on the complex plane. Consider an infinite discontinuous group G of projective transformations T_n

$$z \rightarrow T_n(z) = \frac{\alpha_n z + \beta_n}{\gamma_n z + \delta_n} \quad (3.1)$$

in the sense that no other element of the group is infinitesimally close to I . This means that if we consider a point z of the complex plane, it belongs to a domain of points that are not the transforms of z under any element of the group. This domain is, roughly speaking, a so-called "fundamental region", and any element of the group will map it into another domain of the complex plane. Then the automorphic functions with respect to the group G are meromorphic functions with the property that

$$\phi(T_n(z)) = \phi(z) \quad ; \quad (3.2)$$

any $T_n \in G$.

For example, if the group consists of the translations $T_n(z) = z + n\omega$, then the automorphic functions are just the periodic functions with period ω and the fundamental region any strip of width ω in the complex plane. Similarly, the double translations $T_{n,m}(z) = z + n\omega_1 + m\omega_2$, $\text{Im}(\omega_2/\omega_1) > 0$, give rise to elliptic functions and the fundamental region is the parallelogram with sides ω_1, ω_2 .

The automorphic functions which are of relevance for the general solution of Nielsen's analogue model were considered by Burnside in 1891, inspired precisely by harmonic problems on surfaces with circular holes. I shall give here a short account of his results, and refer the interested reader to his very lucid original paper for more details. Consider a set of $2M$ circles in the complex plane, to be denoted by C_μ, C'_μ ($\mu = 1, \dots, M$) and introduce the M projective transformations T_μ that map C_μ into C'_μ , (see Fig. 3). The relation between T_μ and the circles is as follows. Given T_μ and T_μ^{-1}

$$T_\mu(z) = \frac{a_\mu z + b_\mu}{c_\mu z + d_\mu} \quad ; \quad T_\mu^{-1}(z) = \frac{-d_\mu z + b_\mu}{c_\mu z - a_\mu} \quad (3.3)$$

then the circles $|c_\mu z + d_\mu| = 1$, $|c_\mu z - a_\mu| = 1$ are called the isometric circles of T_μ and T_μ^{-1} , respectively; and they are precisely the two circles we are considering, C_μ and C'_μ . The region of the complex plane outside C_μ is mapped by T_μ into the inside of C'_μ . Now consider the infinite group of projective transformations G that has the T_μ ($\mu = 1, \dots, M$) as generators, that is to say, the infinite group of projective transformations obtained by taking any combination of products of positive and negative powers of the M basic transformations T_μ . Then, Burnside's automorphic functions are automorphic functions with respect to the group G so constructed.

It can be shown that a fundamental region of the group is the part of the complex plane outside all the $2M$ circles. It is well known that the complex plane with the point at infinity included is equivalent - by means of a stereographic projection - to a sphere, the so-called Riemann sphere. If, by bending and stretching the plane, the M pair of circles C_μ and C'_μ are made to coincide by identifying homologous points under T_μ , the fundamental region becomes a sphere with M handles. Burnside automorphic functions are therefore single-valued meromorphic functions on a sphere with M handles.

Let us consider a point ξ_μ on the circle C_μ and its homologous $T_\mu(\xi_\mu)$ on the circle C'_μ , and let us draw a path from ξ_μ to $T_\mu(\xi_\mu)$, (see Fig. 3). The set of M paths so constructed is what Burnside calls "barriers". When the sphere with M handles is constructed by bending and stretching, the circles $C_\mu = C'_\mu$ become the $K_{2\mu-1}$ cycles of a canonical basis, and the "barriers" the dual $K_{2\mu}$ cycles of the same canonical basis.

The theory of Burnside automorphic functions is based on the introduction of the Poincaré ϑ series

$$\vartheta(z, a) = \sum_{\alpha} \frac{1}{(c_{\alpha} z + d_{\alpha})^2} \frac{1}{T_{\alpha}(z) - a} \quad (3.4)$$

where

$$T_{\alpha}(z) = \frac{a_{\alpha} z + b_{\alpha}}{c_{\alpha} z + d_{\alpha}} \quad ; \quad a_{\alpha} d_{\alpha} - b_{\alpha} c_{\alpha} = 1 \quad (3.5)$$

The summation in Eq. (3.4) extends to all the elements of the group. From now on we shall use subindices α , β when we mean arbitrary elements of the group, and μ , ν when we restrict ourselves to the generators. If it were not for the convergence factor $(c_\alpha z + d_\alpha)^{-2}$, ϑ would be an automorphic function. As defined, it is an "automorphic form of dimension -2" with the property that

$$\vartheta(T_\beta(z), a) = (c_\beta z + d_\beta)^2 \vartheta(z, a) \quad (3.6)$$

Moreover, a crucial property is that the difference

$$\vartheta(z, a) - \vartheta(z, T_\beta(a)) = \vartheta(z, -\frac{d_\beta}{c_\beta}) \quad (3.7)$$

is independent of a . Notice that $(-d_\beta/c_\beta)$ is the centre of the isometric circle of the transformation T_β . Moreover, Burnside proved that there are only M independent $\vartheta(z, -d_\beta/c_\beta)$ functions, which can be taken to be $\vartheta(z, J_\nu)$, $\nu = 1, \dots, M$; $J_\nu = (a_\nu/c_\nu)$ being the centre of the circle C'_ν .

These functions $\vartheta(z, J_\nu)$ are important because their integrals $\phi_\nu(z)$

$$\phi_\nu(z) - \phi_\nu(z_0) = \int_{z_0}^z \vartheta(z, J_\nu) dz = \sum_{\alpha} \log \left[\frac{T_\alpha(z) - J_\nu}{T_\alpha(z_0) - J_\nu} \right] \quad (3.8)$$

are the M independent first Abelian integrals that exist on our Riemann surface. Not only that; they are proportional to the elements of a complex basis of first Abelian integrals, because Burnside proved that

$$\oint_{K_{2\mu-1}} d\phi_\nu(z) = 2\pi i \delta_{\mu\nu} \quad (3.9)$$

Moreover, the period matrix corresponding to the dual $K_{2\mu}$ cycles is given by

$$B_{\mu\nu} = \phi_{\mu}(T_{\nu}(z)) - \phi_{\nu}(z) = \int_z^{T_{\nu}(z)} \vartheta(z, J_{\mu}) dz \quad (3.10)$$

It can be shown that $B_{\mu\nu}$ is a symmetric matrix, independent of z . Finally, Burnside also constructed the third Abelian differential in the "complex normalization"

$$d\omega_{ab}(z) = \sum_{\alpha} \frac{1}{(c_{\alpha}z + d_{\alpha})^2} \left[\frac{1}{T_{\alpha}(z) - a} - \frac{1}{T_{\alpha}(z) - b} \right] \quad (3.11)$$

Indeed, the third Abelian integral

$$\omega_{ab}(z) = - \sum_{\alpha} \log \left[\left(\frac{T_{\alpha}(z) - a}{T_{\alpha}(z) - b} \right) \cdot \left(\frac{T_{\alpha}(z_0) - b}{T_{\alpha}(z_0) - a} \right) \right] \quad (3.12)$$

where the branch of the logarithm is such that it makes $\omega_{ab}(z_0) = 0$, does not change when we go along the $K_{2\mu-1}$ cycles [so $d\omega_{ab}(z)$ has zero periods along these cycles].

However, we need $\Omega_{ab}(z)$, the third Abelian integral in the real normalization. In order to construct it, we use the fact that, when the normalization is not specified, a third Abelian integral is defined up to a linear combination of first Abelian integrals, because its singularities are not modified by them. Then we write

$$\Omega_{ab}(z) = \omega_{ab}(z) + \sum_{\mu=1}^M \alpha_{\mu} \phi_{\mu}(z) \quad (3.13)$$

and use Eqs. (2.6) and (2.7). Equation (2.6) gives the M equations

$$\begin{aligned} \operatorname{Re} \oint_{K_{2\mu-1}} d\Omega_{ab}(z) &= \operatorname{Re} \oint_{K_{2\mu-1}} d\omega_{ab}(z) + \sum_{\nu=1}^M \operatorname{Re} [\alpha_{\nu} P_{\mu}(\phi_{\nu}(z))] \\ &= 2\pi i \operatorname{Im} \alpha_{\mu} \cdot \delta_{\mu\nu} = 0 \quad \mu, \nu = 1, \dots, M \end{aligned} \quad (3.14)$$

so we obtain the result that α_μ is real. Moreover, Eq. (2.7) gives another set of M equations that determine α_μ . Using the fact that

$$\omega_{ab}(T_\mu(z)) - \omega_{ab}(z) = \phi_\mu(b) - \phi_\mu(a) \quad (3.15)$$

then Eqs. (2.7), (2.5) and (3.13) give the set of equations

$$\operatorname{Re} [\phi_\mu(b) - \phi_\mu(a)] + \sum_{\nu=1}^M A_{\mu\nu} \alpha_\nu = 0 \quad (3.16)$$

where

$$A_{\mu\nu} = \operatorname{Re} B_{\mu\nu} \quad \mu, \nu = 1, \dots, M \quad (3.17)$$

Since $A_{\mu\nu}$ is non-singular (see previous section), the set of Eqs. (3.16) can be solved and we get, for the third Abelian integral

$$\Omega_{ab}(z) = \omega_{ab}(z) + \sum_{\mu, \nu=1}^M \phi_\mu(z) (\bar{A}^{-1})_{\mu\nu} \operatorname{Re} [\phi_\nu(a) - \phi_\nu(b)] \quad (3.18)$$

so that

$$\operatorname{Re} \Omega_{ab}(z) = \operatorname{Re} \omega_{ab}(z) + \sum_{\mu, \nu=1}^M \operatorname{Re} \phi_\mu(z) (\bar{A}^{-1})_{\mu\nu} \operatorname{Re} [\phi_\nu(a) - \phi_\nu(b)] \quad (3.19)$$

and $\operatorname{Re} \omega_{ab}(z)$ and $\operatorname{Re} \phi_\nu(z)$ are given by the theta series of Eqs. (3.8) and (3.12) by taking the modulus of the logarithm.

Finally, for further comparisons with the operator formalism, we need a formal expression for the real period matrix $A_{\mu\nu}$. Let us first consider $B_{\mu\nu}$. Starting from Eq. (3.10) and using Eq. (3.7)

$$B_{\mu\nu} = \int_b^{T_\nu(b)} \vartheta(z, T_\mu) dz = \int_b^{T_\nu(b)} [\vartheta(z, a) - \vartheta(z, T_\mu(a))] dz \quad (3.20)$$

The independence of $B_{\mu\nu}$ from the points a, b is guaranteed by Eq. (3.7) and the remark following Eq. (3.10). By formally integrating the θ series we get:

(3.21)

$$A_{\mu\nu} = \text{Re } B_{\mu\nu} = \sum_{\alpha} \log \left| \frac{T_{\alpha}(a) - T_{\nu}(b)}{T_{\alpha}(a) - b} \frac{T_{\alpha}(T_{\mu}(a)) - b}{T_{\alpha}(T_{\mu}(a)) - T_{\nu}(b)} \right|$$

4. ONE-LOOP ORIENTABLE AND NON-ORIENTABLE DIAGRAMS

We shall discuss here the case $M=1$, that is to say, the one-loop problem. The group of automorphisms G has only one generator, T , and it therefore consists of all the positive and negative powers of T , to be denoted by T_n , and $T_0 \equiv I$. The Möbius (homographic) transformation T can always be written as

$$\frac{T(v) - \xi_1}{T(v) - \xi_2} = K \frac{v - \xi_1}{v - \xi_2} ; \quad K = \rho e^{i\sigma} \quad (4.1)$$

where ξ_1, ξ_2 are the fixed points and K the multiplier. In this particular case it is much better to make a change of variables and work in the plane :

$$z = \frac{v - \xi_1}{v - \xi_2} \left(\frac{v_0 - \xi_1}{v_0 - \xi_2} \right) \quad (4.2)$$

We shall see at the end of next section, when we discuss the operator formalism, that in the configuration in which the Koba-Nielsen variables are on the unit circle, the limit points of the group are also on the unit circle. If ξ_1, ξ_2, v_0 are all on the unit circle, Eq. (4.2) maps the unit circle to the real axis. Therefore, working with the variable z corresponds, in the operator formalism, to real Koba-Nielsen variables. In this case the fixed points ξ_1, ξ_2 are mapped to $0, \infty$ respectively. Moreover, no generality is lost by making this transformation, and the group elements become simpler

$$z \rightarrow T_n(z) = K^n z \quad ; \quad n=0, \pm 1, \pm 2, \dots \quad (4.3)$$

In this configuration the two circles C_1 and C_1^* are concentric, $|z|=1$ and $|z|=p$, respectively, and the fundamental region is the annulus between them (not to be confused with the original annulus).

When we sew the two circles together, we get a torus, with $C_1 = C_1^*$ becoming the K_1 cycle. The duality diagram, in the orientable case, is the other annulus got by cutting this torus along the cycle K_2 , which corresponds to the real axis in the z plane.

Introduce now the variables

$$\begin{aligned} \omega &= \log z \\ \omega_a &= \log a \end{aligned} \quad (4.4)$$

Then, it is very simple to check that :

$$\begin{aligned} d \log \prod_n [T_n(z) - a] &= d \log \prod_{n=-\infty}^{+\infty} [1 - K^n e^{(\omega - \omega_a)}] \\ &= d \log \vartheta_1 \left(\frac{\omega - \omega_a}{2\pi i} \mid \tau \right) \end{aligned} \quad (4.5)$$

where

$$\tau = \frac{\log K}{2\pi i} = \frac{\log p}{2\pi i} + \frac{\sigma}{2\pi} \quad (4.6)$$

$\vartheta_1(z/\tau)$ being the Jacobi ϑ_1 functions. We use here the notation and conventions of Bateman. Equations (4.5) and (3.12) give the third Abelian integral in the complex normalization as

$$\omega_{ab} = \log \left[\frac{\vartheta_1 \left(\frac{\omega - \omega_a}{2\pi i} \mid \tau \right)}{\vartheta_1 \left(\frac{\omega - \omega_b}{2\pi i} \mid \tau \right)} \right] \quad (4.7)$$

up to constant terms that do not contribute because of momentum conservation. Moreover, in this case there is only one first Abelian integral $\phi(z)$. In this case we have that $J=0$ in Eq. (3.8) (the centre of the isometric circle C_1''). Then the first Abelian differential is ^{*} :

$$d\phi(z) = d \log \prod_n T_n(z) = d \log \prod_{n=-\infty}^{+\infty} K^n e^\omega = d \log e^\omega \quad (4.8)$$

so we obtain the result

$$\phi(z) = \log z \quad (4.9)$$

and the real part of its period B is $A = \log \rho$. Finally, Eq. (3.19) gives, using $z_i = r_i e^{i\theta_i}$, $z_j = r_j e^{i\theta_j}$

$$\operatorname{Re} \Omega_{z_i z_0}(z_j) = \log \left[e^{\frac{\log r_i \log r_j}{\log \rho}} \vartheta_1 \left(\frac{\log z_i - \log z_j}{2\pi i} \middle| \tau \right) \right] \quad (4.10)$$

where we have dropped the z_0 dependent terms that do not contribute because of momentum conservation.

Up to now, the discussion has been very general, and we have constructed the third Abelian integral on a torus, given by Eq. (4.10). Now we must identify our torus with the double of an annulus or of a Möbius strip. In particular, we must identify boundary points on the double because it is on these points that we need our Abelian integrals. The reader is, at this stage, urged to recall our discussion about the construction of D in the cases we are considering.

A) The Annulus

Consider the annulus $R \leq |z| \leq 1$ (see Fig. 4a). Its double is a torus, sewn along the K_2 cycle. Now let us cut this torus along the

^{*})

This argument is not very rigorous. Burnside's convergence proof assumes that ω is not a fixed point. However, (4.9) can be derived rigorously by calculating it before the transformation (4.2).

K_1 cycle, according to the discussion of the previous section. The resulting surface is a section of a pipe which, once straightened, looks again like an annulus A' (not to be confused with the original one) when seen from above. It is easy to see that, in this case, boundary points of B can be taken to be the segments of the real axis contained inside the double, as indicated in Fig. 4b; the segments $\overline{AA}$ corresponding to the external boundary $|z|=1$ and the segment $\overline{BB}$ to the internal one, $|z|=R$. The two bounding circles of A' are the ones that determine the generator of the groups of automorphisms. Clearly, in this case we have in (4.1)

$$\overline{\sigma} = 0 \quad (4.11)$$

and then

$$\tau = \frac{\ln \rho}{2\pi i} \quad (4.12)$$

Remember that, in Eq. (4.11), z_i, z_j are the boundary points on the double, so strictly speaking they must be taken as the points x_i, x_j indicated in Fig. 4b. Moreover, we have :

$$-\log |x_i| \log |x_j| \approx \frac{1}{2} \left[\log \left| \frac{x_i}{x_j} \right| \right]^2 \quad (4.13)$$

where we use the notation $\approx$ to indicate quantities that are equal up to functions of only one variable that do not contribute to the Veneziano integrand because of momentum conservation. Then, let us introduce the variable

$$x_{ij} = \frac{x_i}{x_j} \quad (4.14)$$

At this stage, two possibilities must be distinguished :

- i) the points p_i, p_j are on the same boundary (the line connecting them is "untwisted"). Equation (4.10) gives

$$e^{Re \Omega_{z_i z_o}(z_j)} = e^{\frac{\ln^2 |x_{ij}|}{2 \ln \rho}} \mathcal{D}_1 \left(\frac{\ln x_{ij}}{2\pi i} \middle| \frac{\ln \rho}{2\pi i} \right) \quad (4.15)$$

- ii) the points p_i, p_j are on different boundaries (the line connecting them is twisted). In this case (x_i/x_j) has a phase factor $e^{i\pi}$, and we get

$$e^{\frac{R}{2} \Omega_{z, z_0}(z)} = e^{\frac{\ln^2 |x_{ij}|}{2 \ln \rho}} \vartheta_1 \left(\frac{\ln x_{ij}}{2\pi i} + \frac{1}{2} \middle| \frac{\ln \rho}{2\pi i} \right) \quad (4.16)$$

Equations (4.15) and (4.16) are precisely the functions $\psi(x_{ij})$ and $\psi_T(x_{ij})$ introduced by Gross et al.⁸⁾ because in this case

$$\log \rho = \log \omega = \frac{2\pi^2}{\log R} \quad (4.17)$$

ω being the product of all the Chan variables of the propagators. It is extremely instructive to see what happens if the double of the annulus is cut along the K_2 cycle instead of the K_1 cycle as we did before. In this case the double R and the corresponding boundary points are exhibited in Fig. 4c. Clearly, in this case we must choose $\rho = R^2$. Then the modulus of the torus is given by

$$z = \frac{\log R}{i\pi} \quad (4.18)$$

Several different particular cases must be distinguished.

- i) If the momenta p_i, p_j are inserted in the external boundary of the annulus, $r_i = r_j = 1$ and the contribution of (4.11) to the Veneziano integrand is

$$\left[\vartheta_1 \left(\frac{\phi_i - \phi_j}{2\pi} | z \right) \right]^{-\alpha' p_i \cdot p_j} \quad (4.19)$$

- ii) If the momenta p_i, p_j are inserted in the interval boundary, $r_i = r_j = R$ and the contribution of (4.10) is

$$\left[R^{-1/2} \vartheta_1 \left(\frac{\phi_i - \phi_j}{2\pi} | z \right) \right]^{-\alpha' p_i \cdot p_j} \quad (4.20)$$

The factor $R^{-(\alpha'/2)p_i \cdot p_j}$ is the origin of the so-called Pommeranchuk-like singularity previously discussed by several authors.

- iii) If p_i is inserted in the internal boundary and p_j in the external boundary, $p_i = R$, $p_j = 1$, and the contribution of (4.11) is

$$\left[\vartheta_1 \left(\frac{\phi_i - \phi_j}{2\pi} + \frac{\tau}{2} \mid \tau \right) \right]^{-\alpha' p_i \cdot p_j} = \left[i \bar{R}^{1/4} e^{-\frac{i(\phi_i - \phi_j)}{2}} \vartheta_4 \left(\frac{\phi_i - \phi_j}{2\pi} \mid \tau \right) \right]^{-\alpha' p_i \cdot p_j} \quad (4.21)$$

However, the contribution of the phase factor $e^{-i(\phi_i - \phi_j)/2}$ will be cancelled by another one coming from the configuration $p_j p_i$, where both particles have been interchanged [see Eq. (1.1)]. Equations (4.18)-(4.21) completely agree with the results obtained by previous authors on one loop orientable diagrams. They also agree with the results obtained by cutting the double along the K_1 cycle, indeed, they can be obtained from them by making the Jacobi imaginary transformation - used by Neveu and Scherk to investigate loop divergences - on the ϑ_1 functions of Eqs. (4.15)-(4.16). Using the relation

$$\ln \omega = \frac{2\pi^2}{\ln R} \quad (4.22)$$

Equation (4.18) becomes

$$\tau = - \frac{2\pi i}{\ln \omega} \quad (4.23)$$

In complete agreement with the results of Gross et al. 8).

B) The Möbius Strip

At this stage the reader is urged to recall the construction of the double of the Möbius strip discussed in Section 2. Figure 5a shows a Möbius strip, and Fig. 5b its double cut along the K_2 cycle, which gives then back Fig. 1. In this case there is only one boundary, which corresponds to the circles $|z| = 1$ and $|z| = \rho$, and the identification

of the two circles as shown in Fig. 1b (and Fig.1) means that in this case the argument of the multiplier K of the generator T is [see Eq. (4.1)]

$$\sigma = \pi \quad (4.24)$$

Then the modulus of the torus is

$$\tau' = \frac{\ln \rho}{2\pi i} + \frac{1}{2} \quad (4.25)$$

In this case all the boundary points can be taken on the unit circle, and therefore we get

$$\operatorname{Re} \Omega_{z_i z_0}(z_j) = \log \left[\vartheta_1 \left(\frac{\phi_i - \phi_j}{2\pi} | \tau' \right) \right] \quad (4.26)$$

However, it must be kept in mind that ϕ_i and ϕ_j are the phases of the boundary points in the complex plane where we have constructed the double of the Möbius strip. One could imagine introducing angular co-ordinates on the Möbius strip itself, as indicated in Fig. 5a, by a relation of the form $\theta_i = 2\phi_i$. Therefore, in this case, two possibilities must be distinguished :

- i) contributions of terms like $p_j p_k$ not separated by the "twist".
In this case

$$\phi_k - \phi_j = \frac{\theta_k - \theta_j}{2} \quad (4.27)$$

and we have a contribution

$$\left[\vartheta_1 \left(\frac{\theta_j - \theta_k}{4\pi} | \tau' \right) \right]^{-\alpha' p_j p_k} \quad (4.28)$$

- ii) contributions of terms like $p_i p_j$, separated by the "twist". In this case one has

$$\phi_i - \phi_j = \frac{\theta_i - \theta_j}{2} + \pi \quad (4.29)$$

and we have a contribution

$$\left[\vartheta_1 \left(\frac{\theta_i - \theta_j}{4\pi} + \frac{1}{2} \mid \tau' \right) \right]^{-\alpha' p_i \cdot p_j} \quad (4.30)$$

By making the identification $\ln \rho = \pi^2 (\ln \omega)^{-1}$, Eqs. (4.25), (4.28) and (4.30) also agree with the previous results of Gross et al., obtained in the operator formalism, after transforming them by means of Jacobi's transformation⁸⁾. The reader can easily compare the quantities raised to the power $p_i \cdot p_j$ in Eqs. (4.28) and (4.30) with their ψ_N , ψ_{NT} functions and check that they are the same.

Incidentally, the derivation given here for the non-orientable diagrams also allows to obtain its duality properties in a trivial way; because the method of calculation shows that the location of the twist is topologically irrelevant. It is only when we introduce the angular co-ordinates on the Möbius strip that we must keep track of its location, but the result of the calculation is clearly independent of it.

We close this section with a few comments on the convergence of Poincaré ϑ series. We have seen that in the one-loop orientable case the phase of the multiplier is zero, and in the non-orientable case it is different from zero. In the first case, the transformation T is called "hyperbolic", and in the second case "loxodromic". If all the elements of a group are hyperbolic transformations, and if they have a common invariant circle, it is said to be a Fuchsian group. Otherwise, if it contains some loxodromic transformations, it is said to be a Kleinian group. Then in the one-loop orientable case we are dealing with the Fuchsian group and in the non-orientable case with a Kleinian group.

Burnside's proof of convergence of the series given by Eq. (3.4) originally applies to Fuchsian groups, and use is made of the fact that

all elements of a Fuchsian group leave a circle - the so-called principal circle - invariant (example : the real axis for the above annulus construction) and map interior points of the principal circle into interior points and exterior points into exterior points. Moreover, the isometric circles of the transformations of the group are orthogonal to the principal circle and so intersect it. This is no longer true if the group contains loxodromic transformations, except in the case $\sigma = \pi$ (the only of interest to us) in which there is a principal circle (again the real axis in the above Möbius strip construction) and the isometric circles are orthogonal to it, but the interior of the principal circle is mapped into the exterior, and vice versa. However, a close examination of Burnside's proof reveals that it is independent of this last fact, and therefore it also applies to the non-Fuchsian groups of interest in the dual resonance model. This is true for any number of loops, as we shall discuss in the next section.

5. N LOOP DIAGRAMS IN THE OPERATOR FORMALISM

The example of the one-loop diagram considered in the previous section shows that the construction of the double D of the original Riemann surface dictates the choice of the automorphism group G . In this section, I shall consider the general case of orientable and non-orientable N loop diagrams in the operator formalism, expanding the work done by Lovelace on the subject ³⁾. In particular, I shall present here a more complete discussion of some delicate points associated with the treatment of zero-mode oscillators.

The advantages of the operator formalism are evident : one does not need to stretch one's imagination to construct the double of a complicated non-orientable diagram because explicit formulae are given for the generators S of the group G ; and the measure of integration - missing in the analogue model - is also obtained.

The starting point in the derivation of the "sewing theorem" is Lovelace's N Reggeon vertex ⁹⁾, whose integrand can be written as

$$I = \pi \prod_{1 \leq i < j \leq N} e^{\langle a^{(i)} | U^{(i)} V^{(j)} | a^{(j)} \rangle} \quad (5.1)$$

Strictly speaking, Eq. (5.1) gives the coherent state matrix elements of the integrand of the N Reggeon vertex. Then $|a^{(i)}\rangle$ is an infinite component vector in the Hilbert space of the harmonic oscillators (internal excitations of the Reggeon) plus the Fubini-Veneziano "zero-mode" (centre-of-mass motion of the Reggeon)¹⁰⁾. Its harmonic oscillator components ${}_n|a^{(i)}\rangle$ are, for the time being, real coherent state parameters of the i^{th} Reggeon, and ${}_0|a^{(i)}\rangle = (2\alpha')^{\frac{1}{2}} k_i |0\rangle$, where $|0\rangle$ is a unit vector in the zero-mode subspace. Moreover, $V^{(i)}$ and $U^{(i)}$ are infinite matrices whose matrix elements are obtained by expanding in powers of z a given Möbius transformation

$$\frac{\omega^n}{\sqrt{n}} = \frac{1}{\sqrt{n}} \left(\frac{az+b}{cz+d} \right)^n = \sum_{m=0}^{\infty} U_{nm} \frac{z^m}{\sqrt{m}} \quad (5.2)$$

with the convention $1/\sqrt{0} = 1$. The infinite matrix is completely determined by the Möbius transformation, which, in turn, is determined by specifying how three given points in the complex plane transform under it. Therefore, in Lovelace's notation⁹⁾

$$U^{(i)} = \begin{bmatrix} z_{i-1} & z_i & z_{i+1} \\ 0 & \infty & 1 \end{bmatrix} \quad (5.3)$$

$$V^{(i)} = \begin{bmatrix} \infty & 0 & 1 \\ z_{i-1} & z_i & z_{i+1} \end{bmatrix} \quad (5.4)$$

where the upper line transforms into the lower. Using Eq. (5.2), it can be shown that $U_{no}^{(i)} = V_{on}^{(i)} = 0$. Therefore, there are no zero-mode intermediate state contributions in the matrix multiplication $U^{(i)} V^{(j)}$ ($U_{oo} V_{oo}$ contribute an over-all constant to the Veneziano integrand due to momentum conservation).

It will be very useful in what follows to introduce a symbol defined by

$$\Gamma = \begin{bmatrix} a & b & c \\ \bar{a}^{-1} & \bar{b}^{-1} & \bar{c}^{-1} \end{bmatrix} \quad (5.5)$$

corresponding to the Möbius transformation $\omega = 1/z$. Though Γ is not a Möbius matrix, it acts consistently on Möbius matrices. There are several properties of Γ that are very useful in what follows. Besides the obvious fact that $\Gamma^2 = 1$, we notice that

$$U^{(i)} V^{(i)} = \begin{bmatrix} \infty & 0 & 1 \\ 0 & \infty & 1 \end{bmatrix} = \Gamma \quad (5.6)$$

The twisted and untwisted propagators are defined by

$$D(x) = x^{L_0} = \begin{bmatrix} 0 & \infty & 1 \\ 0 & \infty & x \end{bmatrix} \quad (5.7)$$

$$P(x) = x^{L_0} \Omega (1-x)^W = \begin{bmatrix} 0 & \infty & 1 \\ x & 1 & 0 \end{bmatrix} \quad (5.8)$$

We shall use the symbol $\Delta(x)$ to represent either the twisted or the untwisted propagators. Using Eq. (5.2) we obtain in both cases $\Delta_{00} = 1$. We shall redefine $\Delta_{00}(x) = \exp(-k^2 \ln x / 2)$. Another important property of the symbol Γ we shall need later on is that it inverts both Δ_μ

$$\Gamma \Delta_\mu \Gamma = \Delta_\mu^{-1} \quad (5.9)$$

In order to construct an M loop diagram, we have to sew M pairs of Reggeons in the N Reggeon vertex, and then perform the loop momentum integrations. We shall discuss these two steps separately.

A) Sewing Reggeons

The N Reggeon vertex is invariant under an over-all projective transformation of the Koba-Nielsen variables z_i . This projective transformation only changes the z_i , the coherent state variables $|a^{(i)}\rangle$ are not affected by it⁹⁾. We are therefore free to do a projective transformation, sew the Reggeons by integrating over coherent state parameters and, at the end, undo the initial projective transformation if we wish.

Suppose we start with Koba-Nielsen variables on the unit circle²⁾. We then map the unit circle to the real axis by means of a projective transformation. The matrices $U^{(i)}$ and $V^{(i)}$ are now real, and following Lovelace's rules for matrix multiplication, it is easy to prove that, in this configuration

$$U^{(i)} = V^{(i)\dagger} \Gamma \quad (5.10)$$

and therefore, adopting the convention

$$U^{(i)} V^{(i)} = 0 \quad (5.11)$$

to cancel diagonal terms, the integrand of the N Reggeon vertex can be written in the more symmetric form

$$I = e^{\frac{1}{2} \sum_{ij} \langle a^{(i)} | V^{(i)\dagger} \Gamma V^{(j)} | a^{(j)} \rangle} \quad (5.12)$$

At this stage, the N Reggeon vertex can be formally factorized by means of a real functional integral

$$I = [det \Gamma]^2 \int \frac{\delta f}{\sqrt{\pi}} e^{-\frac{1}{2} \langle f | \Gamma | f \rangle + \sum_{i=1}^N \langle f | V^{(i)} | a^{(i)} \rangle} \quad (5.13)$$

This is admittedly only a formal step because Γ does not exist as a matrix in the Hilbert space of harmonic oscillators and $(\det \Gamma)^2$ has no meaning. However, we shall undo the δf functional integral at the end and cancel the factor $(\det \Gamma)^2$. The final result is correct and can be justified by more rigorous - and less transparent - methods ^{*}). Moreover, here by δf we mean

$$\frac{\delta f}{\sqrt{\pi}} = \prod_{n=1}^{\infty} \frac{\delta f_n}{\sqrt{\pi}} \quad (5.14)$$

Notice that there is no integration over a zero-mode component because of our previous remark that zero-mode intermediate states do not contribute to the matrix product $U^{(i)} V^{(j)}$.

At this stage we are ready to start sewing Reggeons. If the μ^{th} sewn pairs are $\langle a^{(i)} |$ and $| a^{(j)} \rangle$ we relabel

$$U_{\mu} = U^{(i)} = V_{\mu}^+ \Gamma \quad ; \quad \tilde{U}_{\mu} = U^{(j)} = \tilde{V}_{\mu}^+ \Gamma \quad (5.15)$$

$$\langle a^{(i)} | = \langle a_{\mu} | \quad ; \quad | a^{(j)} \rangle = | a_{\mu} \rangle \quad \mu=1, \dots, M \quad (5.16)$$

The matrices U_{μ} and $\tilde{U}_{\mu}$ must be distinguished because they depend on different Koba-Nielsen variables $(z_{\mu-1}, z_{\mu}, z_{\mu+1})$ and $(\tilde{z}_{\mu-1}, \tilde{z}_{\mu}, \tilde{z}_{\mu+1})$, respectively. Moreover, we shall keep the index i to label unsewn Reggeons. Finally, the integrand of the N Reggeon vertex with the propagators $\Delta_{\mu}(x_{\mu})$ included - x_{μ} being the Chan variable of line μ - and the unsewn Reggeons in the ground state is given by :

^{*}) This step is done in this way in order to emphasize the fact that we only need a real functional integral to linearize the dependence of the N Reggeon vertex on the coherent state variables $| a^{(i)} \rangle$. In fact, the Hermitian adjoint operation is not invariant under Möbius transformations. So if we define $\langle f |$ to be Möbius invariant and equal to the Hermitian adjoint $\langle f |$ in the unit circle configuration, then $\langle \tilde{f} | = \langle f | \Gamma$ after transformation to the real axis.

$$I' = [\det \Gamma]^2 \int \frac{\delta f}{\sqrt{\pi}} \exp \left\{ -\frac{1}{2} \langle f | \Gamma | f \rangle + \sum_i (2\alpha')^{1/2} p_i \langle z_i | f \rangle + \sum_{\mu=1}^M [\langle a_{\mu} | \Delta_{\mu} \tilde{V}_{\mu} \Gamma | f \rangle + \langle f | \tilde{V}_{\mu} | a_{\mu} \rangle] \right\} \quad (5.17)$$

where we have used $\langle 0 | V^{(i)} = \langle z_i |$, $\langle z_i |$ being a vector with components $z_i^{-n}/\sqrt{n}$. Let us now split $|a_{\mu}\rangle$ into its zero-mode components $|a_{\mu}^0\rangle$ and the internal excitation components $|\underline{a}_{\mu}\rangle$. The trace of I' over the oscillator modes is taken by making $|\underline{a}_{\mu}\rangle$ complex, identifying $\langle \underline{a}_{\mu}|$ with its complex conjugate and integrating over $|\underline{a}_{\mu}\rangle$ with the measure

$$\int \frac{1}{\pi} \delta \text{Re } \underline{a}_{\mu} \delta \text{Im } \underline{a}_{\mu} e^{-\langle a_{\mu} | a_{\mu} \rangle} \quad (5.18)$$

The procedure gives the result

$$\text{Tr } I' = [\det \Gamma]^2 \int \frac{\delta f}{\sqrt{\pi}} \exp \left\{ -\frac{1}{2} \langle f | \left\{ 1 - \sum_{\mu=1}^M (S_{\mu} + S_{\mu}^+) \right\} \Gamma | f \rangle + \sum_i (2\alpha')^{1/2} p_i \langle z_i | f \rangle + \sum_{\mu=1}^M \langle a_{\mu}^0 | X_{\mu} | f \rangle \right\} \quad (5.19)$$

where

$$S_{\mu} = \tilde{V}_{\mu} \Delta_{\mu} V_{\mu} \quad (5.20)$$

$$X_{\mu} = \Delta_{\mu} V_{\mu}^+ + \tilde{V}_{\mu}^+ \quad (5.21)$$

Finally, the integration over δf is done, with the result :

$$\begin{aligned} \text{Tr } I' = & \frac{1}{\|1 - \sum_{\mu=1}^M (S_{\mu} + S_{\mu}^{\dagger})\|^2} e^{-\frac{1}{2} \sum_{i,j} p_i p_j \langle z_i | \Gamma \frac{1}{1-z} | z_j \rangle} \\ & e^{-\frac{1}{2} \sum_{i,\mu} p_i \langle z_i | \Gamma \frac{1}{1-z} X_{\mu}^{\dagger} | a_{\mu}^0 \rangle} e^{-\frac{1}{2} \sum_{j,\nu} \langle a_j^0 | X_{\nu} \Gamma \frac{1}{1-z} | z_j \rangle p_j} \\ & e^{-\frac{1}{2} \sum_{\mu,\nu} \langle a_{\nu}^0 | X_{\nu} \Gamma \frac{1}{1-z} X_{\mu}^{\dagger} | a_{\mu}^0 \rangle} \end{aligned} \quad (5.22)$$

where

$$\Sigma = \sum_{\mu=1}^M (S_{\mu} + S_{\mu}^{\dagger})$$

At this stage we can undo our original projective transformation and go back to Koba-Nielsen variables on the unit circle. From now on, we shall work in this configuration. In this case, Eq. (5.12) no longer holds. However, an inspection of the matrices $U^{(i)}$, $V^{(i)}$ shows that, when the z variables are on the unit circle

$$U^{(i)} = V^{(i)\dagger} \quad (5.23)$$

This equation replaces therefore Eq. (5.11). It also follows from Eq. (5.23) that the continuation of Eq. (5.22) for $\text{Tr } I'$ from real z 's to z 's on the unit circle is done by suppressing the Γ factor that occurs in all matrix elements. Therefore, the final result is

$$\begin{aligned} \text{Tr } I' = & \Delta^{-2} e^{-\frac{1}{2} \sum_{i,j} p_i p_j \langle z_i | \frac{1}{1-z} | z_j \rangle} e^{-\frac{1}{2} \sum_{i,\mu} p_i \langle z_i | \frac{1}{1-z} X_{\mu}^{\dagger} | a_{\mu}^0 \rangle} \\ & e^{-\frac{1}{2} \sum_{j,\nu} \langle a_j^0 | X_{\nu} \frac{1}{1-z} | z_j \rangle p_j} e^{-\frac{1}{2} \sum_{\mu,\nu} \langle a_{\nu}^0 | X_{\nu} \frac{1}{1-z} X_{\mu}^{\dagger} | a_{\mu}^0 \rangle} \end{aligned} \quad (5.24)$$

where we have denoted by Δ the determinant that occurs in Eq. (5.22).

The first important point to notice here is that the S_μ are unitary operators when restricted to the subspace of harmonic oscillators such as we have them in Eq. (5.16). A heuristic proof (due to Lovelace) can be given, using the symbol Γ previously introduced

$$S_\mu^+ = U_\mu^+ \Delta_\mu \tilde{V}_\mu^+ = V_\mu \Delta_\mu \tilde{U}_\mu = \tilde{U}_\mu^{-1} \Gamma \Delta_\mu \Gamma \tilde{V}_\mu^{-1} = \tilde{U}_\mu^{-1} \Delta_\mu^{-1} \tilde{V}_\mu^{-1} \quad (5.25)$$

by (5.9). More rigorously (also Lovelace) S_μ leaves the unit circle invariant [see Eq. (5.44) below] and therefore belongs to the unitary D_1^+ representation of $O(2,1)$ as can be checked from comparison of Gliozzi's generators ¹¹⁾ with Bargmann's ¹²⁾. S_μ acts only on the $n \geq 1$ part of the Reggeon. Therefore

$$\Sigma = \sum_{\mu=1}^M (S_\mu + S_\mu^{-1}) \quad (5.26)$$

We must keep in mind that we have not yet applied the constraints given by Eq. (5.4), arising from the fact that there are no diagonal elements in the N Reggeon vertex. For the sewn Reggeons, such a constraint means that $S_\mu S_\mu^{-1} = 0$, so that all terms containing adjacent factors S_μ and S_μ^{-1} are to be zeroed in the expansion of $(1 - \Sigma)^{-1}$. What is left are all the elements of a group G of projective transformations generated by the M operators S_μ . We shall therefore write

$$\frac{1}{1 - \Sigma} = \sum_{\alpha} T_{\alpha} \quad (5.27)$$

where the sum runs over all the elements of the group of automorphisms generated by the S_μ . In Eq. (5.27), T_{α} is an infinite matrix, not to be confused with the projective transformation $T_{\alpha}(z)$ defined by Eq. (3.5). Consider the matrix elements of $(1 - \Sigma)^{-1}$. Using the fact that ¹¹⁾

$${}_n(T_{\alpha} | z_i \rangle) = \sum_m (T_{\alpha})_{nm} \frac{z_i^m}{\sqrt{m}} = \frac{(T_{\alpha}(z_i))^m}{\sqrt{m}} \quad (5.28)$$

we get

$$\langle z_i | T_\alpha | z_j \rangle \approx \sum_\alpha \log |z_i - T_\alpha(z_j)| \quad (5.29)$$

where we use the symbol $\approx$ to indicate equal up to a quantity that does not contribute to the Veneziano integrand because of momentum conservation. Using Eq. (3.12), it is quite straightforward to check that this is part of the contribution to the Neumann function $N_B(z_i, z_j)$ of a Riemann surface whose double is defined by the set of generators S_μ . It is precisely the part of $N_B(z_i, z_j)$ coming from $\text{Re } \omega_{z_i z_0}(z_j)$, the third Abelian integral of the Riemann surface in the complex normalization. The matrix element given by Eq. (5.29) are the only ones in Eq. (5.24) that are independent of the loop momenta. We now proceed to show that the loop momentum integration contributes precisely the terms needed to change the third Abelian integral from the complex to the real normalization.

B) Loop Momentum Integrations

The loop momentum integration in this case just amounts to an integration over the zero-mode components $|a_\mu^0\rangle$, which are proportional to the loop momenta k_μ . However, this step requires a very careful discussion. Let us first notice that $|a_\mu^0\rangle$ is always multiplied by X_μ^+ . Using the fact that the symbol Γ can be written as

$$\Gamma = U_\mu V_\mu \quad (5.30)$$

and that it inverts both the twisted and the untwisted propagators, we have

$$\begin{aligned} X_\mu^+ |a_\mu^{(0)}\rangle &= (V_\mu \Delta_\mu + \tilde{V}_\mu) |a_\mu^0\rangle = \\ &= (U_\mu^{-1} \Delta_\mu^{-1} \Gamma + \tilde{V}_\mu) |a_\mu^0\rangle \end{aligned} \quad (5.31)$$

In order to decide what is $\Gamma |a_\mu^0\rangle$, we must remember how the zero-mode is introduced. Any operator valued function of z , $F(z)$, can be expanded in terms of irreducible representations of $SU(1,1)$ - isomorphic to

the group of projective transformations. One possible way of doing it is to expand in terms of the representations $D_{\epsilon}^{(+)}$ of the $SU(1,1)$ algebra ^{10),13)}

$$F(z) = \sum_{m=0}^{\infty} a^{(m)} \left[\frac{(m-1+2\epsilon)!}{m!} \right]^{1/2} z^{\epsilon+m} \quad (5.32)$$

Let ϵ be a small positive real number. Expanding to first order in ϵ one gets ^{10),13)}

$$F(z) = \frac{1}{2} (1 + \epsilon \ln z) \left(\frac{1}{\epsilon} \sqrt{2\alpha'} p - \frac{i q}{\sqrt{2\alpha'}} \right) + \sum_{m=1}^{\infty} a^{(m)} \frac{z^m}{\sqrt{m}} \quad (5.33)$$

In the limit $\epsilon \rightarrow 0$, Eq. (5.33) is not well defined by itself. However, we know that singular terms will cancel anyway in the Veneziano integrand ¹⁰⁾. If we forget about terms that either vanish or diverge when $\epsilon \rightarrow 0$, because total momentum conservation eventually cancels divergent terms, the "zero-mode" component is

$$\frac{1}{2} \left[\sqrt{2\alpha'} p \ln z - \frac{i q}{\sqrt{2\alpha'}} \right] \quad (5.34)$$

If we consider now $\Gamma F(z) \Gamma$, the net effect is to replace z by $1/z$ in Eq. (5.32). Taking the limit $\epsilon \rightarrow 0$ just changes the sign of $\ln z$ in Eq. (5.34). Since q_0 does not contribute to the Veneziano amplitude ¹⁰⁾, we conclude that the zero component - the momentum - changes sign under Γ (in fact this is just momentum conservation for the sewn line). Then Eq. (5.31) can be written

$$X_{\mu}^{\dagger} |a_{\mu}^{\circ}\rangle = (S_{\mu}^{-1} - 1) |\tilde{z}_{\mu}\rangle (2\alpha')^{1/2} k_{\mu} \quad (5.35)$$

where

$$_n |\tilde{z}_{\mu}\rangle = \left(\tilde{V}_{\mu} |0\rangle \right) = \frac{\tilde{z}_{\mu}^n}{\sqrt{n}} \quad (5.36)$$

Therefore, the integration of $\text{Tr } I'$ over the loop momenta reads

$$\begin{aligned} & \Delta^{-2} e^{-\frac{1}{2} \sum_{ij} p_i p_j \langle z_i | \frac{1}{1-\Sigma} | z_j \rangle} \int \left(\prod_{\mu=1}^n d^4 k_\mu \right) e^{-\frac{1}{2} \sum_{i,\mu} p_i k_\mu \langle z_i | \frac{1}{1-\Sigma} (S_\mu^{-1}) | \tilde{z}_\mu \rangle} \\ & e^{-\frac{1}{2} \sum_{j,\nu} p_j k_\nu \langle \tilde{z}_\nu | (S_\nu^{-1}) \frac{1}{1-\Sigma} | z_j \rangle} e^{-\frac{1}{2} \sum_{\mu,\nu} k_\mu k_\nu \langle \tilde{z}_\mu | (S_\mu^{-1}) \frac{1}{1-\Sigma} (S_\nu^{-1}) | \tilde{z}_\nu \rangle} \end{aligned} \quad (5.37)$$

These Gaussian integrations can be easily done. They involve an inversion of the $M \times M$ matrix that multiplies k_μ, k_ν and that can be done because the matrix is positive definite. The final result is (after elimination of diagonal terms)

$$\frac{1}{\Delta^2 \det A} e^{-\frac{1}{2} \sum_{ij} p_i p_j N_B(z_i, z_j)} \quad (5.38)$$

where $N_B(z_i, z_j)$ is given by

$$N_B(z_i, z_j) = \sum_{\alpha} \langle z_i | T_{\alpha} | z_j \rangle + \sum_{\mu,\nu=1}^M \langle z_i | \frac{1}{1-\Sigma} (S_\mu^{-1}) | \tilde{z}_\mu \rangle (A^{-1})_{\mu\nu} \cdot \langle \tilde{z}_\nu | (S_\nu^{-1}) \frac{1}{1-\Sigma} | z_j \rangle \quad (5.39)$$

and

$$A_{\mu\nu} = \langle \tilde{z}_\nu | (S_\nu^{-1}) \frac{1}{1-\Sigma} (S_\mu^{-1}) | \tilde{z}_\mu \rangle \quad (5.40)$$

We have used the notation $N_B(z_i, z_j)$ anticipating the fact that the right-hand side of Eq. (5.39) is the Neumann function of the Riemann surface whose double is defined by the generators S_μ . We have already seen that the first term of the right-hand side can be written as

$$\sum_{\alpha} \langle z_i | T_{\alpha} | z_j \rangle \approx R_{\omega_{z_i z_j}}(z_j) \quad (5.41)$$

Moreover, using Eqs. (3.7) and (3.8) we immediately get that

$$\begin{aligned} \sum_{\alpha} \langle z_i | T_{\alpha} (S_{\mu}^{-1} - 1) | \tilde{z}_{\mu} \rangle &= \sum_{\alpha} \log |z_i - T_{\alpha}(\tilde{z}_{\mu})| - \sum_{\alpha} \log |z_i - T_{\alpha}(S_{\mu}^{-1}(\tilde{z}_{\mu}))| \\ &= \text{Re } \phi_{\mu}(z_i) \end{aligned} \quad (5.42)$$

where $\phi_{\mu}(z_i)$ is a first Abelian integral, independent of $\tilde{z}_{\mu}$. Moreover, we can also see that $A_{\mu\nu}$ given by Eq. (5.40) is precisely the real period matrix given by Eq. (3.21); and is independent of $\tilde{z}_{\mu}$, $\tilde{z}_{\nu}$ for the same reasons that the period matrix of Section 3 is independent of a , b . Therefore we conclude that the right-hand side of Eq. (5.39) is precisely the Neumann function. The calculation done in this section neatly shows that the loop momentum integration contributes to the Neumann function the piece containing the first Abelian integrals, that is to say, the piece associated with the existence of harmonic functions on the surface which are the real parts not of single-valued, but rather of multiple-valued analytic functions.

C) The Analogue Riemann Surface

We have succeeded in proving that the operator formalism gives the M loop Veneziano integrand in terms of the Neumann function of a certain Riemann surface B whose double D is a sphere with M handles defined by the M generators S_{μ} . More explicitly, the generators are given by the projective transformations

$$S_{\mu} = \begin{bmatrix} \infty & 0 & 1 \\ \tilde{z}_{\mu-1} & \tilde{z}_{\mu} & \tilde{z}_{\mu+1} \end{bmatrix} \begin{bmatrix} 0 & \infty & 1 \\ 0 & \infty & x_{\mu} \end{bmatrix} \begin{bmatrix} z_{\mu-1} & z_{\mu} & z_{\mu+1} \\ 0 & \infty & 1 \end{bmatrix} \quad (5.43)$$

if the μ^{th} propagator is untwisted, and by

$$S_{\nu} = \begin{bmatrix} \infty & 0 & 1 \\ \tilde{z}_{\nu+1} & \tilde{z}_{\nu} & \tilde{z}_{\nu+1} \end{bmatrix} \begin{bmatrix} 0 & \infty & 1 \\ x_{\mu} & 1 & 0 \end{bmatrix} \begin{bmatrix} z_{\nu-1} & z_{\nu} & z_{\nu+1} \\ 0 & \infty & 1 \end{bmatrix} \quad (5.44)$$

if the $\vee$ th propagator is twisted. It is now easy to see that $S_{\vee}$ given by Eq. (5.44) leaves the unit circle invariant. Indeed, reading from right to left we have : unit circle anticlockwise $\rightarrow$ real axis leftwards $\rightarrow$ real axis rightwards $\rightarrow$ unit circle anticlockwise. Moreover $S_{\vee}$ transforms the interior (or exterior) of the unit circle into itself. Therefore, $S_{\vee}$ is what we have called in Section 4 a hyperbolic transformation, of the type found in the one-loop planar calculation. Remember that we have sewn the symmetric vertex, and therefore a twisted propagator gives an untwisted line. Therefore, for planar diagrams, all the generators are hyperbolic transformations, the group is Fuchsian, and Burnside's proof of convergence of the ϑ series holds at once.

It is also straightforward to check that in the case of the untwisted propagator (twisted line) the generator S_{μ} given by Eq. (5.43) transforms the unit circle anticlockwise $\rightarrow$ unit circle clockwise and the interior of the unit circle into its exterior and vice versa. Therefore, the group is not Fuchsian but the fact that the unit circle is invariant is sufficient to generalize Burnside's proof of the ϑ series, as discussed in Section 4. In the original N Reggeon vertex, one has to integrate over all the Koba-Nielsen variables. This means that the M loop Veneziano formula is obtained by integrating Eq. (5.38) (some trivial factors involving α_0 must still be added to the integrand) over the Koba-Nielsen variables of the unsewn Reggeons, z_i ; and the Koba-Nielsen variables of the sewn Reggeons, $z_{\mu}, \tilde{z}_{\mu}$. This last integration corresponds to variations of the Riemann surface B, whereas the former one corresponds to variations of the boundary points where the currents are attached, in the language of the analogue model.

This interpretation of the integrand of the M loop Veneziano formula is particularly useful to analyze the singularities of the M loop amplitude. Domain variational theory can be used to analyze how the Neumann function varies when the boundaries of the Riemann surface are varied. It can be shown in general that variations of the boundary give rise to singularities in the amplitude only when they change the topological structure of the Riemann surface, i.e., when a hole shrinks to zero or when two holes collide. A first step in this direction has been given by Lovelace³⁾, who analyzed the so-called Pomeranchuk-like singularity to all orders in the number of loops.

D) The Measure of Integration

The measure of integration is explicitly exhibited in Eq. (5.38) besides some trivial factors containing α_0 that I have not bothered to include. It must be emphasized that $\det A$ is a finite determinant, taken in the M dimensional space where $A_{\mu\nu}$ is defined. On the contrary, Δ is an infinite determinant, and care must be exercised in its definition. We have

$$\frac{1}{\Delta} = \frac{1}{\det(1-Z)} = e^{\text{Tr} \ln(1-Z)} = e^{\sum_{N=1}^{\infty} \frac{\text{Tr}(Z^N)}{N}} \quad (5.45)$$

However, in computing $\sum^N$ we must remember our "rule of thumb" $S_{\mu} S_{\mu}^{-1} = 0$ arising from the absence of diagonal terms in the original N Reggeon vertex. Then, Δ^{-1} can be written as

$$\frac{1}{\Delta} = e^{\sum_{\alpha} \frac{\text{Tr}(T_{\alpha})}{n_{\alpha}}} \quad (5.46)$$

where the summation extends to all the elements of the group of automorphisms and n_{α} is an integer defined by

$$n_{\alpha} = \sum_{\mu=1}^M n_{\alpha\mu} \quad (5.47)$$

$n_{\alpha\mu}$ being the number of times that the generator S_{μ} is contained in T_{α} , counting (+1) for either S_{μ} or S_{μ}^{-1} . I have already discussed in Section 4 the fact that any arbitrary projective transformation $T_{\alpha}(z)$ can be written as

$$\frac{T_{\alpha}(z) - \xi_{1\alpha}}{T_{\alpha}(z) - \xi_{2\alpha}} = K_{\alpha} \frac{z - \xi_{1\alpha}}{z - \xi_{2\alpha}} \quad (5.48)$$

in terms of its fixed points ξ_1, ξ_2 (assuming ξ_1, ξ_2 finite and $\xi_1 \neq \xi_2$) and the multiplier K_{α} . It is easy to show that K_{α} is the ratio of the two eigenvalues of the 2×2 matrix

$$\begin{pmatrix} a_{\alpha} & b_{\alpha} \\ c_{\alpha} & d_{\alpha} \end{pmatrix} \quad (5.49)$$

and that K_α is as good a multiplier as K_α^{-1} (the order in which one must take the two fixed points is not defined). Therefore, multipliers can always be taken with $|K_\alpha| \leq 1$. This is known to be the correct choice in the one-loop case ¹¹⁾. Thus

$$K(\bar{T}_\alpha) = K(T_\alpha) \quad (5.50)$$

It has been shown in Ref. 11) that the eigenvalues of the infinite matrix T_α are given by

$$(\lambda_\alpha)_n = K_\alpha^n \quad (5.51)$$

Then, in this case, it is simple to compute $\text{Tr}(T_\alpha)$ and write Δ^{-2} in the form

$$\frac{1}{\Delta^2} = \prod_{n=1}^{\infty} e^{4 \sum_{\alpha} \frac{K_\alpha^n}{n_\alpha}} \quad (5.52)$$

where the extra power of 2 in the exponential is due to the fact that T_α and T_α^{-1} have the same multiplier. We are therefore summing over half of the elements of the group in Eq. (5.52); inverses have been eliminated. Then we split this half of the group in two classes of elements: a first class of elements T_α that can be written in the form $T_\alpha = (T_\beta)^R$ ($R=2,3,\dots$) and a second class of elements that cannot be written in such a form. For example, $T_1 T_4^2 T_3$ belongs to the second class, whereas $T_1 T_2 T_1 T_2$ belongs to the first class.

Using the fact that

$$n(T_\beta^N) = N n_\beta ; \quad K(T_\beta^N) = K_\beta^N \quad (5.53)$$

the first class of elements can be summed explicitly and we get the final result

$$\Delta^2 = \prod_{\alpha}' \prod_{n=1}^{\infty} (1 - K_\alpha^n)^{4/n_\alpha} \quad (5.54)$$

where $\prod'$ means an infinite product restricted to the second class of group elements. It is very simple to identify in Eq. (5.54) the M partition functions associated with each individual loop. All of them have $n_\alpha = 1$, and K_α are just the multipliers of the generators of the group G . However, we see that there are now infinitely many partition functions which could in principle produce new exponential divergences. The renormalization programme must therefore be re-examined. A first step in this direction has been given by Kaku and Scherk¹⁴⁾, who analyzed in detail the two-loop case with a formula analogous to Eq. (5.54). They concluded that, besides the exponential divergences associated with each individual loop, there is only one extra partition function that gives an exponential divergence, and that is the one in which K_α is the multiplier of $S_1 S_2$. This is true only when spurious states have been eliminated from the two loops. The general problem of N loops is currently under investigation.

ACKNOWLEDGEMENT

I am grateful to Professor C. Lovelace for enlightening discussions and suggestions.

R E F E R E N C E S

- 1) H.B. Nielsen, NORDITA preprint (1969);
D. Fairlie and H.B. Nielsen, Nuclear Phys. B20, 637 (1970).
- 2) Z. Koba and H.B. Nielsen, Nuclear Phys. B10, 633 (1969).
- 3) C. Lovelace, CERN preprint TH.1203 (1970) - to be published in Phys.Letters.
- 4) M. Schiffer and D. Spencer, "Functionals of Finite Riemann Surfaces", Princeton (1954).
- 5) L. Ford, "Automorphic Functions", Chelsea (1951).
- 6) W. Burnside, Proc.London Math.Soc. 23, 49 (1891).
- 7) H.S.M. Coxeter, "Introduction to Geometry", Wiley (1961).
- 8) D.J. Gross, A. Neveu, J. Scherk and J. Schwarz, Princeton preprint, to be published.
- 9) C. Lovelace, CERN preprint TH.1184 (1970) - to be published in Phys.Letters.
- 10) S. Fubini and G. Veneziano, M.I.T. preprint, to be published.
- 11) V. Alessandrini, D. Amati, M. Le Bellac and D. Olive, CERN preprint TH.1160 (1970), to be published.
- 12) V. Bargmann, Ann.Math. 48, 568 (1947).
- 13) L. Clavelli and P. Ramond, N.A.L. preprint, to be published.
- 14) M. Kaku and J. Scherk, Berkeley preprint, to be published.

FIGURE CAPTIONS

- Figure_1 Construction of the two-sheeted orientable covering of the Möbius strip ; by cutting a hole in the two-sheeted covering of the projective plane.
- Figure_2 Canonical cycles $K_{2\mu-1}$, $K_{2\mu}$ on a sphere with two handles.
- Figure_3 Fundamental region for Burnside automorphic functions and canonical cycles on the sphere with M handles.
- Figure_4 a) The annulus considered in Section 5.
b) Its double cutted along the K_1 cycle.
c) Its double cutted along the K_2 cycle.
- Figure_5 a) The Möbius strip.
b) Construction of its double.

FIG. 1

FIG. 2

FIG. 3

FIG. 5a

FIG. 5b