CM-P00058648

Ref.TH.1217-CERN

LOWER BOUNDS TO THE $\pi^0 \pi^0$ SCATTERING LENGTH FROM
UNITARITY, ANALYTICITY AND CROSSING SYMMETRY

A.K. Common ^{*)} and R. Wit ^{**)}

CERN - Geneva

A B S T R A C T

It is shown that for $\pi^0 \pi^0 - \pi^0 \pi^0$ scattering a rigorous lower bound to the value of the scattering amplitude $F(s,t,u)$ at $s=4$ and $t=2$ may be obtained from unitarity, analyticity and crossing symmetry. For a sufficiently small ω wave scattering length the corresponding threshold value of the forward scattering amplitude $F(4,0,0)$ must be greater than -2.30 pion Compton wavelengths, which corresponds to a scattering length -1.15 .

^{*)} On leave of absence from the University of Kent at Canterbury, England until 1st October 1970.

^{**)} On leave of absence from the Jagellonian University, Cracow, Poland.

Ref.TH.1217-CERN

20 August 1970

1. INTRODUCTION

The number and variety of exact results on $\pi\pi$ scattering which have been obtained in the last few years are rather impressive ¹⁾. Analyticity, unitarity and crossing symmetry prove to be very powerful constraints on the possible behaviour of the scattering amplitudes. Many of the rigorous relations deal with the low energy structure of the partial wave amplitudes and the scattering amplitudes themselves. One of the well-known results of this type is the lower bound to the $\pi^0\pi^0$ S wave scattering length a_0 . The existence of such a bound in the axiomatic framework was proved by Łukaszyk and Martin ²⁾. Coming out as a by-product of a general proof of absolute bounds of the scattering amplitude inside the Mandelstam triangle, the first rigorous lower bound ^{*})

$$-19 \leq a_0 \quad (1.1)$$

was certainly overestimated and not very restrictive. In a subsequent paper by Bonnier and Vinh Mau ³⁾ this bound was substantially improved becoming

$$-4 \leq a_0. \quad (1.2)$$

From the technique developed by these two authors it also turned out that any improvement on the absolute upper bounds of the modulus of the scattering amplitude inside the Mandelstam triangle leads at the same time to an improvement on the lower bound to a_0 . Therefore, any effort to get a better lower bound to a_0 can go in two different ways: 1) a direct improvement on a_0 ; 2) an improvement on the upper bound of the scattering amplitudes in the unphysical region $0 < s < 4$, $0 < t < 4$, $0 < u < 4$. Both possibilities will be discussed in the next sections.

^{*}) Throughout this paper we use the natural system of units:

$$\mu = \hbar = c = 1.$$

2. NOTATIONS AND PRELIMINARIES

Let us consider the $\pi^0 \pi^0$ scattering amplitude $F(s, t, u)$ for which the partial wave decomposition reads

$$F(s, t) = \frac{\sqrt{s}}{k} \sum_{\substack{l=0 \\ \text{even}}}^{\infty} (2l+1) f_l(s) P_l \left(1 + \frac{2t}{s-4} \right), \quad (2.1)$$

where $s = 4(k^2 + 1)$ is the squared energy in the centre-of-mass system and k the relative c.m. momentum. Unitarity yields

$$|f_l(s)|^2 \leq \text{Im} f_l(s), \quad s \geq 4$$

and the scattering lengths are defined by

$$a_l = \lim_{s \rightarrow 4} f_l(s) / k^{2l+1}.$$

As already mentioned by one of us ⁴⁾ when evaluating the amplitude $F(s, t, u)$ at threshold for fixed, positive t we sum in fact over these scattering lengths ^{*)}. For example, when $s=4$ and $t=2$,

$$F(4, 2, -2) = 2a_0 + \frac{2}{\sqrt{\pi}} \sum_{l=2}^{\infty} (2l+1) a_l 2^l \frac{\Gamma(l+1/2)}{\Gamma(l+1)}. \quad (2.2)$$

Now ⁵⁾

$$a_{l+2} \leq \frac{1}{16} \frac{(l+2)(l+1)}{(l+5/2)(l+3/2)} a_l, \quad l \geq 2$$

^{*)} It will be taken as understood from now on that all sums are over even l .

so that

$$F(4,0,0)=2a_0 \leq F(4,2,-2) \leq 2a_0 + \frac{32a_2}{\sqrt{\pi}} \sum_{l=2}^{\infty} (2l+1) 2^{-l} R(l) \frac{\Gamma(l+1/2)}{\Gamma(l+1)} \quad (2.3)$$

where

$$R(l) = \begin{cases} 1 & , \quad l=2, \\ \prod_{k=4}^l \frac{k(k-1)}{k^2 - 1/4} & , \quad l \geq 4. \end{cases}$$

With the present estimates ⁶⁾ for $a_2 (=1.6 \times 10^{-5})$, it appears that if we can obtain a lower bound to $F(4,2,-2)$ it will be also substantially a lower bound to $2a_0$. To derive such a lower bound for $F(4,2,-2)$ we can make the assumption

$$F(4,2,-2) < 0 \quad (2.4)$$

since if the reverse were true we would have immediately a better lower bound, i.e., zero.

The inequality (2.4) is crucial for our method and as we see, there exists a large margin for further improvements on a_0 which still will not violate it. Therefore, from now on we concentrate our attention on $F(4,2,-2)$ and, instead of looking for a lower bound to a , we start to investigate the lower bound to $F(4,2,-2)$. Of course, the choice $t=2$ is arbitrary and, in fact, any positive value of t in the range $0 < t < 4$ can be introduced into play. The algebraic structure of our equations becomes, however, very simple for $t=2$ and this circumstance will make our considerations more transparent ^{*)}.

^{*)} On the other hand, crossing symmetry and the convexity property of the scattering amplitude in the unphysical region make the lower bound to $F(4,2,-2)$ most important in our discussion - cf., Ineqs (3.9), (3.10).

As seen later the numerical analysis has been performed for all interesting values of t .

We would like to start with the twice subtracted dispersion relation

$$F(s, 2, 2-s) - F(s_0, 2, 2-s_0) = \frac{2(s-2+s_0)(s-s_0)}{\pi} \int_{\frac{1}{2}}^{\infty} \frac{ds' (s'-1) A(s', 2)}{(s'-s_0)(s'-s)(s'-2+s)(s'-2+s_0)} \quad (2.5)$$

where $A(s', 2) \equiv \text{Im } F(s', 2)$ and $0 < s_0 < 4$. There are two immediate consequences which follow from Eq. (2.5) and the positivity of $A(s, 2)$:

- i) if $F(4, 2, -2) < 0$ then $F(s, 2)$ does not have any zero in the twice cut s plane;
- ii) for $1 \leq s \leq 4$ the function $F(s, 2)$ is an increasing function of s .

For arbitrary t in the interval $0 \leq t \leq 4$ this statement is true for all $4 \geq s \geq 2-t/2$. The statement i) can be generalized in a similar way: if $F(4, t, -t) < 0$ for $0 \leq t < 4$ then $F(s, t)$ is free of zeros. A direct application of this result is the Cauchy theorem for the function $\ln F(s, 0)$ which leads to

$$|F(2, 0, 2)| = \exp \left[\frac{1}{\pi} \int_{\frac{1}{2}}^{\infty} \frac{ds \ln |F(s, 0)|}{\sqrt{s(s-4)} (s-2)} \right]. \quad (2.6)$$

3. A SIMPLE TECHNIQUE FOR OBTAINING LOWER BOUNDS TO $F(s, t, u)$ IN MANDELSTAM'S TRIANGLE *)

Under the assumption $F(4, 2, -2) < 0$ we apply in this section a simple method for obtaining lower bounds to $F(s, t, u)$ inside the Mandelstam triangle. This method, introduced in its general form in

*) The use of this method was suggested to us by Dr. A. Martin.

physics by Meiman ⁷⁾, was further developed by Martin ⁸⁾, and it proves to be very useful for our particular purposes.

We start by applying the Schwartz inequality to the forward scattering amplitude so that

$$\begin{aligned}
 |F(s,0)|^2 &= \left| \frac{\sqrt{s}}{k} \sum_{l=0}^{\infty} (2l+1) f_l(s) \right|^2 \\
 &\leq \frac{s}{k^2} \sum_{l=0}^{\infty} (2l+1) |f_l(s)|^2 P_l(x) \sum_{l=0}^{\infty} (2l+1) / P_l(x) \\
 &\leq H(s,2) \frac{\sqrt{s}}{k} \sum_{l=0}^{\infty} (2l+1) / P_l(x)
 \end{aligned} \tag{3.1}$$

where we have used the unitarity condition for the partial waves; here $x = 1 + 4/(s-4)$. Now, we introduce Ineq. (2.4) into (2.5) and make use of (3.1) to get finally

$$|F(2,0,2)| \geq \frac{8}{\pi} \int_{\frac{1}{2}}^{\infty} \frac{ds (s-1) |F(s,0)|^2}{\sqrt{s(s-4)} (s-2) s (s+2) \sum_{l=0}^{\infty} (2l+1) / P_l(x)} . \tag{3.2}$$

It is convenient to map the twice-cut s plane onto a unit circle in the w plane by means of the two transformations

$$z = \left(\frac{s-2}{2} \right)^2, \quad w = \frac{i - \sqrt{z-1}}{i + \sqrt{z+1}} = r e^{i\alpha}.$$

Then, formula (3.2) can be written in the form

$$\frac{1}{|F(2,2,0)|} \geq \frac{1}{2\pi} \int_{-\pi}^{\pi} d\alpha W(\alpha) |F_R(e^{i\alpha})|^2, \quad (3.3)$$

where

$$F_R(e^{i\alpha(s)}) = F(s,0)/F(2,0,2)$$

and

$$W(\alpha) = \frac{1}{2} \frac{1+2V}{(1+v)(2+v) \sum_{l=0}^{\infty} (2l+1)/P_l(1+2(v-1)^{-1})},$$

$$V = \sqrt{1 + tg^2 \alpha/2}. \quad (3.4)$$

Using the inequality of arithmetic and geometric means in (3.3)

$$\frac{1}{|F(2,0,2)|} \geq \exp \left[\frac{1}{2\pi} \int_{-\pi}^{\pi} d\alpha W(\alpha) \right] \times \exp \left[\frac{1}{2\pi} \int_{-\pi}^{\pi} d\alpha \ln |F_R(e^{i\alpha})|^2 \right]. \quad (3.5)$$

Since, as stated before, due to Ineq. (2.4) the forward scattering amplitude does not have zeros in the s plane so that the function $\ln F(s(w),0)$ is analytic in the unit circle in the w plane. Applying Cauchy's theorem we get

$$\frac{1}{2\pi} \int_{-\pi}^{\pi} d\alpha \ln |F_R(e^{i\alpha})|^2 = 2 \ln F_R(\kappa=0); \quad (3.6)$$

note, however, that the point $w=0$ corresponds to $s=2$ and, therefore, $\ln F_R(w=0)=0$. In this way we finally obtain from (3.5)

$$|F(2,0,2)| \leq \exp\left(-\frac{1}{2\pi} \int_{-\pi}^{\pi} d\alpha \ln W(\alpha)\right) \quad (3.7)$$

and numerical integration yields

$$-8.90 \leq F(2,0,2) \leq 0. \quad (3.8)$$

It is important to remember that the bound (3.8) has been obtained under the assumption that $F(4,2,-2)$ is negative. Similar calculations have been made for other values of s , t and u (but under the same condition) and the results are as follows

$$\begin{aligned} -12.30 &\leq F(2/3, 4/3, 0) \leq 0, & -8.90 &\leq F(2,2,0) \leq 0, \\ -10.80 &\leq F(4/3, 4/3, 4/3) \leq 0, & -11.10 &\leq F(3,2,-1) \leq 0. \end{aligned}$$

Some of these bounds can be improved immediately. Convexity and crossing symmetry of the scattering amplitude $F(s,t,u)$ provide us with the following chains of inequalities

$$-8.90 \leq F(2,2,0) \leq F(3,2,-1) \leq F(4,2,-2) \leq 0, \quad (3.9)$$

$$-8.90 \leq F(2,0,2) \leq F(8/3, 0, 4/3) \leq F(4,0,0) \leq 0, \quad (3.10)$$

and thus, the inequality (3.8) appears to be the most restrictive one.

4. USE OF ŻUKASZUK AND MARTIN'S METHOD

In deriving the bound to $F(2,2,0)$ in the previous section some majorizations were made like, e.g., the Schwartz inequality. Therefore, we may expect to get better bounds by introducing more refined techniques, e.g., the method due to Żukaszuk and Martin²⁾. We will adopt the latter to get a direct lower bound on $F(4,2,-2)$ and hence effectively to a_0 .

By definition

$$|F(s,0)| \leq \frac{\sqrt{s}}{k} \sum_{\ell=0}^{\infty} (2\ell+1) |f_{\ell}(s)| \equiv \bar{F}(s), \quad (4.1)$$

and from the unitarity condition

$$\bar{H}(s,t) \equiv \frac{\sqrt{s}}{k} \sum_{\ell=0}^{\infty} (2\ell+1) |f_{\ell}(s)|^2 P_{\ell}(x) \leq H(s,t) \quad (4.2)$$

where $x = 1 + 2t(s-4)^{-1}$.

Let $\phi(\bar{F}, s, 2)$ be the minimum of $\bar{A}(s, 2)$ for fixed $\bar{F}(s)$. This minimum occurs for the following distribution of partial waves

$$|f_{\ell}(s)| = 1, \quad \ell = 0, 2, \dots, 2L \quad (4.3)$$

$$|f_{\ell}(s)| = c(s)/P_{\ell}(x), \quad \ell = 2L+2, \dots \quad (4.4)$$

$$P_{2L(s)}(x) < c(s) \leq P_{2L(s)+2}(x). \quad (4.5)$$

For given $\bar{F}(s)$ the relations (4.3), (4.4) and (4.5) define $c(s)$ and $L(s)$ uniquely. It is also very easy to prove that $\partial \phi / \partial \bar{F} > 0$ so that $\phi(|F|, s, t) \leq \phi(\bar{F}, s, t)$. More generally, all the functions

$$\begin{aligned} & \phi(\bar{F}, s, t), & \frac{\partial \phi}{\partial \bar{F}} \\ & \phi(\bar{F}, s, t) / \bar{F}, \end{aligned}$$

are increasing functions ²⁾ of $\bar{F}$.

Unitarity and the assumption (2.4) allow us to write (2.5) and (2.6) in the following form

$$-F(4, 2, -2) + \frac{16}{\pi} \int_1^{\infty} \frac{ds (s-1) \phi(\bar{F}, s, 2)}{(s-4)(s-2)s(s+2)} \leq |F(2, 2, 0)| \leq \exp \left[\frac{4}{\pi} \int_2^{\infty} \frac{ds \ln \bar{F}(s)}{\sqrt{s(s-4)}(s-2)} \right]. \quad (4.6)$$

Once again we map onto the unit circle in the w plane to get

$$|F(4, 2, -2)| + \frac{1}{\pi} \int_0^{\pi} d\alpha W_1(\alpha) \phi(\bar{F}, \alpha) \leq \exp \left[\frac{1}{\pi} \int_0^{\pi} d\alpha \ln \bar{F}(\alpha, 2) \right] \quad (4.7)$$

where

$$\phi(\bar{F}, s, 2) = \phi(\bar{F}, \alpha)$$

and

$$W_1(\alpha) = \frac{1 + 2\sqrt{1 + \tan^2 \alpha/2}}{\tan \alpha/2 (2 + \sqrt{1 + \tan^2 \alpha/2})}.$$

Since $\phi(\bar{F}, \alpha)/\bar{F}$ increases with $\bar{F}$ for any fixed α , Ineq. (4.7) does not allow for any unlimited growth in $\bar{F}$ and, therefore, we get an upper bound to the right-hand side of (4.7). The extremum occurs by the calculus of variations when

$$W_1(\alpha) \bar{F}(\alpha) \frac{\partial \phi}{\partial \bar{F}} = 1, \quad (4.8)$$

where λ is independent of α . Let us rewrite the Ineq. (4.7) in the following way

$$0 \leq |F(4,2,-2)| \leq \mathcal{D}(\lambda) \equiv E(\lambda) - I(\lambda) \quad (4.9)$$

where by definition

$$I(\lambda) = \frac{1}{\pi} \int_0^{\pi} d\alpha H_1(\alpha) \phi(\bar{F}, \alpha), \quad E(\lambda) = \exp \left[\frac{1}{\pi} \int_0^{\pi} d\alpha \ln \bar{F}(\alpha) \right].$$

It is obvious that not all values of λ are allowed: for some values of λ Ineq. (4.3) might be violated. Starting from any value of λ for which the Ineq. (4.9) holds in its strong form and increasing the argument of $I(\lambda)$ and $E(\lambda)$ we finally get, for a critical value of

$$I(\lambda_{cr}) = E(\lambda_{cr}).$$

This value of λ gives us, according to (4.6), a possible upper bound to $|F(2,2,0)|$. As will be proved in the Appendix it is at the same time a true upper bound to $|F(2,2,0)|$.

Let us consider the behaviour of $\mathcal{D}(\lambda)$. For $\lambda = 0$ we obtain $\mathcal{D}(0) = I(0) - E(0) = 0$ [cf. Eqs. (4.7) and (4.8)]. What we are now interested in is the behaviour of $\mathcal{D}(\lambda)$ for λ in the interval $0 \leq \lambda < \lambda_{cr}$. Obviously $\mathcal{D}(\lambda)$ has there an upper bound ($= E(\lambda_{cr})$) but its behaviour may not be in general very simple although $E(\lambda)$ and $I(\lambda)$ are both increasing functions of λ . More detailed information about $\mathcal{D}(\lambda)$ comes out from the computer. What we observe is that in an interval $0 < \lambda < \lambda_{max}$ the function $\mathcal{D}(\lambda)$ increases monotonically and for $\lambda_{max} < \lambda < \lambda_{cr}$ it decreases (also in a monotonic way). The unique maximum at λ_{max} provides us with an upper bound on $|F(4,2,-2)|$ as seen from Ineq. (4.3).

The results of our numerical analysis are

$$\lambda_{cr} = 6.00, \quad E(\lambda_{cr}) = 8.50,$$

$$\lambda_{max} = 1.40, \quad E(\lambda_{max}) = 4.50, \quad \mathcal{D}(\lambda_{max}) = 2.30.$$

In other words, starting from the assumption $F(4,2,-2) < 0$ we have proved that

$$-8.50 \leq F(2,2,0) \leq 0, \quad (4.10)$$

$$-2.30 \leq F(4,2,-2) \leq 0. \quad (4.11)$$

In comparison with (3.8) we have slightly improved the lower bound to $F(2,2,0)$ but the main result is contained in (4.11). Indeed, if we take into account the estimate of the higher partial waves contributions given by (2.3) (with a very small value of a_2), we get immediately from (4.11)

$$-1.15 < a_0. \quad (4.12)$$

Thus we have improved by a factor 3 on the lower bound to a_0 given in Ref. 3). Note, however, that for $|F(4,2,-2)| = 2.30$ the corresponding upper bound to $|F(2,2,0)|$ reads

$$|F(2,2,0)| \leq 4.50.$$

Similar upper bounds can be obtained for $|F(4/3,4/3,4/3)|$ and $|F(3,2,-1)|$. Introducing these improved bounds into the estimates obtained by Bonnier and Vinh Mau it may happen that we improve also significantly their older result. We postpone the discussion of this rather interesting question to the next section.

5. INVERSE AMPLITUDE APPROACH

As we have learned in Section 2, if $F(4,2,-2) < 0$ the function $F(s,2)$ does not have any zero in the whole s plane. Even if $F(4,2,-2)$ is positive the function $F(s,2) - F(4,2)$ is still different from zero everywhere in the s plane except at threshold ⁴⁾ [where it vanishes like $(s-4)^{1/2}$]. Therefore, the function

$[F(s,2)-F(4,2,-2)]^{-1}$ is analytic in the cut s plane but to write a Cauchy formula for it we need some knowledge about the asymptotic behaviour of $F(s,2)$. Suppose

$$\int_s^\infty \frac{ds}{\phi(s,2)} < \infty ; \quad (5.1)$$

then certainly

$$- \frac{1}{F(s,2)-F(4,2)} = \frac{2}{\pi} \int_4^\infty \frac{ds' (s'-1) \operatorname{Im} F(s',2)}{(s'-s)(s'-2+s) \left[(\operatorname{Re} F(s',2)-F(4,2))^2 + (\operatorname{Im} F(s',2))^2 \right]} . \quad (5.2)$$

In fact, the assumption (5.1) means

$$\lim_{s \rightarrow \infty} \phi(s,2)/\ln s = \infty . \quad (5.3)$$

Note that if

$$|F(s,2)| (\ln s)^{1+\epsilon} \xrightarrow{s \rightarrow \infty} 0 \quad (5.4)$$

the Cauchy formula for $F(s,2)$ can be written without subtractions and in this case we get immediately

$$0 < F(4,2,-2) \quad (5.5)$$

since $\operatorname{Im} F(s,2) > 0$ for $s \geq 4$.

Of course, if (5.4) holds the dispersion relation (2.5) remains true but (5.2) certainly does not. So, starting with the Cauchy formula (5.2) we lose some generality of our predictions for a_0 but this loss will be small since the difference between (5.3) and (5.4) consists only in slightly more than two powers of $(\ln s)$ in the asymptotic behaviour of $F(s,2)$.

What kind of useful information can be drawn from (5.2)? If we neglect the real part contribution in the denominator of the integrand, introduce the unitarity condition in its very weak form (4.2), put $s=2$ and make use of (2.4) we finally get

$$1 + |F(4,2,-2)| \times I \leq |F(2,2,0)| \times I \leq I \times E \quad (5.6)$$

where

$$I = \frac{2}{\pi} \int_1^{\infty} \frac{ds (s-1)}{(s-2)s \phi(\bar{F}, s, 2)} \quad , \quad E = \exp \left(\frac{4}{\pi} \int_1^{\infty} \frac{ds \ln \bar{F}(s)}{\sqrt{s(s-4)} (s-2)} \right).$$

Now, in close analogy to what has been proved in Section 4 we expect that Ineq. (5.6) also does not allow for an unlimited growth ^{*)} of $\bar{F}(s)$. Therefore, we ask for a minimum of I for fixed $|F(4,2,-2)|$ and given E (note that ϕ is an increasing function of $\bar{F}$). The extremum condition on $\bar{F}(s)$ yields in this case

$$\frac{\sqrt{s(s-4)} \frac{(s-1) \bar{F}}{s \phi^2} \frac{\partial \phi}{\partial \bar{F}}}{} = 2\beta \quad (5.7)$$

Given β we solve this equation for all physical values of s then calculate $I(\beta)$ and $E(\beta)$. For β between 0.05 and 400.0 we observe that $I(\beta)$ and $I(\beta)E(\beta)$ are increasing functions of β but $E(\beta)$ decreases with growing argument ^{**)}. For the critical

^{*)} To prove this assertion some additional assumptions must be introduced.

^{**)} Note that Eq. (5.7) is only an analogue of Eq. (4.8). The latter leads to an automatic increase of $\bar{F}$ and ϕ with increasing λ (for any fixed α) since $\partial \phi / \partial \bar{F}$ is an increasing function of $\bar{F}$. It is very hard to deduce a similar conclusion from the more complicated structure of (5.7) but just as in the case when the method of Zukaszuk and Martin is used we are only interested in the behaviour of integrals of ϕ and $\bar{F}$ over s as functions of the parameters λ and β .

value of β for which $E(\beta_{cr})I(\beta_{cr})=1$ the Ineq. (5.6) obviously gives $F(4,2,-2)=0$. Then, in the interval $\beta_{cr} < \beta < \beta_{max}$, the maximum values of $|F(4,2,-2)|$ allowed by Ineq. (5.6) are increasing but in the range $\beta_{max} < \beta < 400$ the function $|F_{max}(4,2,-2;\beta)|$ reveals a monotonic decrease. The numerical analysis gave

$$\beta_{cr} = 0.27, \quad E(\beta_{cr}) \leq 8.40$$

$$\beta_{max} = 1.0, \quad E(\beta_{max}) \leq 4.40, \quad |F_{max}(4,2,-2;\beta_{max})| \leq 2.30.$$

With obvious changes one gets in exactly the same way the relevant bounds to $F(3,2,-1)$ and $F(8/3,4/3,0)$. Before listing them we would like to show, however, how to get an upper bound on $|F(4/3,4/3,4/3)|$ since in this case the approach presented above does not apply directly.

From the Cauchy formula *)

$$\frac{1}{F(s, 4/3)} = \frac{2}{\pi} \int_{\frac{1}{2}}^{\infty} \frac{ds' (s'-2 + 2/3) \operatorname{Im} F(s', 4/3)}{|F(s', 4/3)|^2 (s'-s)(s'-4+s+4/3)}$$

we get in a straightforward way

$$|F(4/3, 4/3, 4/3)| \leq |F(8/3, 4/3, 0)| \sup_{s \geq \frac{1}{2}} \frac{(s-4/3)^2}{s(s-8/3)};$$

hence

$$|F(4/3, 4/3, 4/3)| \leq 1.34 |F(8/3, 4/3, 0)|. \quad (5.8)$$

Therefore, from an upper bound on $|F(8/3,4/3,0)|$ one gets immediately an upper bound $|F(4/3,4/3,4/3)|$. This is the way in which we obtain the estimates for $F(4/3,4/3,4/3)$.

*) This Cauchy formula holds since as previously $F(s,4/3)$ has no zeros when $F(4,4/3,-4/3) < 0$.

The qualitative results we have got for the corresponding critical values of β are as follows ^{*)}

$$\begin{aligned} -10.50 \leq F(8/3, 4/3, 0) \leq 0, & \quad -8.40 \leq F(2, 2, 0) \leq 0, \\ -14.10 \leq F(4/3, 4/3, 4/3) \leq 0, & \quad -9.20 \leq F(3, 2, -1) \leq 0. \end{aligned} \quad (5.9)$$

Similarly, for the relevant values of $\beta_{\max}$ we found

$$\begin{aligned} -3.00 \leq F(4, 4/3, -4/3) \leq 0, & \quad -2.30 \leq F(4, 2, -2) \leq 0; \\ & \quad -2.40 \leq F(4, 2, -2) \leq 0. \end{aligned} \quad (5.10) \quad (**)$$

If the lower bounds in (5.10) were saturated we would get the inequalities

$$\begin{aligned} -5.30 \leq F(8/3, 4/3, 0) \leq 0, & \quad -4.40 \leq F(2, 2, 0) \leq 0, \\ -7.10 \leq F(4/3, 4/3, 4/3) \leq 0, & \quad -4.80 \leq F(3, 2, -1) \leq 0. \end{aligned} \quad (5.11)$$

Finally, again as in Section 3, convexity and crossing symmetry yield

$$\begin{aligned} F(2, 2, 0) &\leq F(3, 2, -1), \\ F(2, 0, 2) &\leq F(8/3, 4/3, 0). \end{aligned}$$

Therefore, using again (5.8)

$$\begin{aligned} -8.40 \leq F(8/3, 4/3, 0) \leq 0, \\ -11.30 \leq F(4/3, 4/3, 4/3) \leq 0, & \quad -8.40 \leq F(3, 2, -1) \leq 0, \end{aligned} \quad (5.12)$$

^{*)} The first and third inequalities hold for $F(4, 4/3) < 0$ and second and fourth inequalities hold for $F(4, 2) < 0$.

^{**) This inequality comes out from the numerical analysis of this section applied to $F(3, 2, -1)$ in place of $F(2, 2, 0)$.}

and similar inequalities follow ^{*)} from (5.11). The last ones will be used explicitly in the next section.

6. CONCLUSIONS

Let us briefly summarize the results we have obtained in the last three sections (for simplicity we discuss only the case $t = 2$).

The extended Meiman's techniques give the following upper bound to $F(2,2,0)$

$$|F(2,2,0)| \leq 8.90. \quad (6.1)$$

The method of Łukaszuk and Martin provides us with

$$|F(2,2,0)| \leq 8.50, \quad (6.2)$$

$$|F(4,2,-2)| \leq 2.30. \quad (6.3)$$

If the last inequality were saturated we would get a better bound for $|F(2,2,0)|$

$$|F(2,2,0)| \leq 4.50. \quad (6.4)$$

From the inverse amplitude approach one gets

$$|F(2,2,0)| \leq 8.40 \quad (6.5)$$

^{*)} It should be noted that these bounds are derived under the assumption that $F(4,2,-2) \leq 0$ while bounds for the corresponding amplitudes in (5.3) are derived under the slightly weaker assumption $F(4,4/3,-4/3) \leq 0$.

$$|F(4,2,-2)| \leq 2.30 \quad (6.6)$$

and if the first inequality becomes an equality we have

$$|F(2,2,0)| \leq 4.40. \quad (6.7)$$

Despite of quite different ways in which the results were obtained they are fairly close to each other. This indicates the crucial role of the assumption (2.4) which was built in from the beginning in all our considerations. The inverse amplitude bounds, which to the extent indicated in the previous section are not completely rigorous, passed through several numerical tests for the stability of the results against approximations which have been made during the calculations. The cross-checks turned out to be satisfactory^{*)} and, therefore, we take the bounds from the last section as a starting point for the discussion given below.

What is the relation between our results and the lower bound to a_0 obtained by Bonnier and Vinh Mau? Introducing the constraints on $|F(3,2,-1)|$, $|F(2,2,0)|$ and $|F(4/3,4/3,4/3)|$ which follow from convexity, (5.11) and (5.12) into the upper bound on $|F(s,t)|$ given by Eq. (18) of Ref. 3), we found that

$$-2.30 \leq a_0. \quad (6.8)$$

This is an essential improvement in comparison with (1.2) but Ineq. (6.8) is still not compatible with our estimate

$$-1.15 \leq a_0 \quad (6.9)$$

*) Note, however, that the bounds published in Refs. 4) and 9) have been slightly relaxed.

(higher partial waves combination being neglected). Unfortunately, Ineq. (6.3) is also not very restrictive if practical applications are taken into account. The best solution obtained from a detailed analysis of the existing data suggests ¹⁰⁾

$$a_0 \simeq 0.02 \tag{6.10}$$

and therefore the Ineq. (6.3) should be considered rather as an example of how far we can proceed at the moment using only analyticity, unitarity, crossing symmetry and computer facilities. If further improvements could be made to the bound (6.3) to bring it much closer to the value given by (6.10) (if it does not change in the meantime) then the situation would be much more interesting.

ACKNOWLEDGEMENTS

We would like to thank Dr. D. Atkinson for a clarifying discussion. We are very grateful to Dr. A. Martin for suggesting to us the method developed in Section 3, for providing us with the proof contained in the Appendix and for helpful comments.

The warm hospitality extended to us at the Theoretical Study Division is acknowledged.

A P P E N D I X

We wish to present here the proof that the condition (4.3) $E(\lambda) - I(\lambda) \geq 0$ puts an upper bound on $E(\lambda)$ and that it is given by $E(\lambda_{cr})$. This result has been used in the original work by Zukaszuk and Martin ²⁾ [see in particular the discussion after Eq. (13)] but the proof was not given there ^{*)}.

Take any general value of $\lambda = \lambda_0$ for which

$$I(\lambda_0) < E(\lambda_0)$$

i.e.,

$$\frac{1}{\pi} \int_0^\pi d\alpha W_1(\alpha) \phi(\bar{F}_0, \alpha) \leq \exp \left[\frac{1}{\pi} \int_0^\pi d\alpha \ln \bar{F}_0(\alpha) \right] \quad (\text{A.1})$$

and

$$\bar{F}_0 \frac{\partial \phi}{\partial \bar{F}_0} = \lambda_0 . \quad (\text{A.2})$$

Also, by definition of λ_{cr}

$$\frac{1}{\pi} \int_0^\pi d\alpha W_1(\alpha) \phi(\bar{F}_{cr}, \alpha) = \exp \left[\frac{1}{\pi} \int_0^\pi d\alpha \ln \bar{F}_{cr}(\alpha) \right] \quad (\text{A.3})$$

where

$$\bar{F}_{cr} \frac{\partial \phi}{\partial \bar{F}_{cr}} = \lambda_{cr} . \quad (\text{A.4})$$

^{*)} We would like to thank Dr. A. Martin for providing this proof.

We wish to prove that

$$\exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \bar{F}_0(\alpha)\right] \leq \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \bar{F}_{cr}(\alpha)\right]. \quad (\text{A.5})$$

Since both $\partial\phi/\partial\bar{F}$ and $\partial^2\phi/\partial\bar{F}^2$ are positive ²⁾

$$\phi(\bar{F}_0) - \phi(\bar{F}_{cr}) \gg (\bar{F}_0 - \bar{F}_{cr}) \frac{\partial\phi}{\partial\bar{F}_{cr}}. \quad (\text{A.6})$$

Hence, from (A.6) and then (A.1) and (A.3)

$$\begin{aligned} \frac{1}{\pi} \int_0^{\bar{\pi}} d\alpha [\bar{F}_0(\alpha) - \bar{F}_{cr}(\alpha)] \frac{\partial\phi}{\partial\bar{F}_{cr}} &< \frac{1}{\pi} \int_0^{\bar{\pi}} d\alpha [\phi(\bar{F}_0, \alpha) - \phi(\bar{F}_{cr}, \alpha)] \\ &< \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \bar{F}_0(\alpha)\right] - \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \bar{F}_{cr}(\alpha)\right]. \end{aligned} \quad (\text{A.7})$$

Using (A.4) in the left-hand side of the above inequality

$$\begin{aligned} \frac{\lambda_{cr}}{\pi} \int_0^{\bar{\pi}} d\alpha \left(\frac{\bar{F}_0(\alpha)}{\bar{F}_{cr}(\alpha)} - 1 \right) &< \left\{ \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \bar{F}_{cr}(\alpha)\right] \right\} \times \\ &\times \left\{ \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \frac{\bar{F}_0(\alpha)}{\bar{F}_{cr}(\alpha)}\right] - 1 \right\} \end{aligned}$$

and applying the inequality of the arithmetic and geometric means

$$0 \leq \left\{ \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \bar{F}_{cr}(\alpha)\right] - \lambda_{cr} \right\} \times \left\{ \exp\left[\frac{1}{\pi}\int_0^{\bar{\pi}} d\alpha \ln \frac{\bar{F}_0(\alpha)}{\bar{F}_{cr}(\alpha)}\right] - 1 \right\}. \quad (\text{A.8})$$

From the convexity of ϕ we get

$$\phi(\bar{F}_{cr}, \alpha) < \bar{F}_{cr} \frac{\partial \phi}{\partial \bar{F}_{cr}} = \lambda_{cr} \quad (\text{A.9})$$

since $\phi = 0$ when $\bar{F} = 0$. Therefore, (A.3) and (A.9) lead to

$$\exp \left[\frac{1}{\pi} \int_0^{\pi} d\alpha \ln \bar{F}_{cr}(\alpha) \right] = \frac{1}{\pi} \int_0^{\pi} d\alpha \phi(\bar{F}_{cr}, \alpha) < \lambda_{cr} \quad (\text{A.10})$$

and introducing this result in (A.18) we obtain finally

$$\exp \left[\frac{1}{\pi} \int_0^{\pi} d\alpha \ln \bar{F}_0(\alpha) \right] - \exp \left[\frac{1}{\pi} \int_0^{\pi} d\alpha \ln \bar{F}_{cr}(\alpha) \right] \leq 0$$

which proves our initial statement.

It follows from above that there is only one value of $\lambda = \lambda_{cr}$ for which $E(\lambda_{cr}) = I(\lambda_{cr})$. Indeed, suppose there were two such values λ_1 and λ_2 . Then, from the preceding analysis we get $E(\lambda_1) = E(\lambda_2)$ and since $E(\lambda)$ increases monotonically with λ we must have $\lambda_1 = \lambda_2$. Hence λ_{cr} is unique.

R E F E R E N C E S

- 1) A. Martin, Lecture given at IX Internationale Universitätswochen für Kernphysik, Schladming (1970) (Lecture notes).
- 2) L. Łukaszyk and A. Martin, Nuovo Cimento 52, 122 (1967).
- 3) B. Bonnier and R. Vinh Mau, Phys.Rev. 165, 1323 (1967);
cf. Phys.Letters 24B, 477 (1967).
- 4) R. Wit, CERN preprint TH.1182 (to be published in Nuclear Phys.).
- 5) A. Martin, Nuovo Cimento 47, 265 (1967);
A.K. Common, Nuovo Cimento 63, 863 (1969);
F.J. Yndurain, Nuovo Cimento 64, 225 (1969).
- 6) J.C. Le Guillou, A. Morel and H. Navelet, private communication;
F.J. Yndurain has informed us that the Veneziano model gives a
very similar value for the 2 wave scattering length.
- 7) N.N. Meiman, Soviet Phys. JETP 17, 830 (1963).
- 8) A. Martin, in High Energy Physics, Vienna (p.155) (1965).
- 9) R. Wit, to be published in Nuovo Cimento.
- 10) D. Morgan and G. Shaw, Columbia University preprint, NYO-1932(2)-164
(1970).