CP VIOLATION AND THE $K \rightarrow \mu^+ \mu^-$ PUZZLE

M.K. Gaillard
CERN - Geneva
and
Laboratoire de Physique Théorique et
Particules Élémentaires
Orsay *)

A B S T R A C T

The unitarity relations for $K \rightarrow \mu^+ \mu^-$ are studied under the assumption of CPT alone, and including $\gamma\gamma$, $\pi\pi$, and the CP even $\pi\pi\gamma$ states in the absorptive parts for both K_S and K_L . A new bound for $K_S \rightarrow \mu\mu$ is obtained and it is shown explicitly that superweak theory contradicts present experimental data under these assumptions.

*) Laboratoire associé au C.N.R.S.

1. - INTRODUCTION

It is well known that unitarity together with CP and CPT invariance leads to a lower bound ¹⁾ on the $K_L \rightarrow \mu^+ \mu^-$ decay rate if only the two-photon intermediate state contributes to the absorptive part of the amplitude. This bound is in disagreement with the experimental upper bound ²⁾ for that process.

Assuming CPT invariance alone, Christ and Lee ³⁾ have recently shown that the present experimental information on $K_L \rightarrow \mu^+ \mu^-$ and $K_L \rightarrow \gamma\gamma$ implies a large branching ratio :

$$\Gamma(K_S \rightarrow \mu^+ \mu^-) / \Gamma_S \geq 5 \times 10^{-7} \quad (1)$$

if only the two photon state contributes to the absorptive part for $K_L \rightarrow \mu^+ \mu^-$.

In this note we generalize their result by allowing for an arbitrarily large $\pi\pi$ contribution and using experimental information on $K_S \rightarrow \mu^+ \mu^-$. We shall also show explicitly that if the super weak theory of CP violation is assumed, there remains a discrepancy with experiment under these assumptions.

2. - FORMALISM

Unitarity and CPT give the following relations for T matrix elements :

$$i(T_{i\hat{i}} - T_{i\hat{i}}^*) = \sum_n T_{i\hat{n}} T_{\hat{n}\hat{i}}^* = \sum_n T_{i\hat{n}}^* T_{n\hat{i}} \quad (2)$$

where $\hat{n}$ is the CPT conjugate of n . In the decay $K \rightarrow \mu^+ \mu^-$ there are two final states which have opposite CP and add incoherently in the rates, so that we may consider them separately. We denote these states by

$$\begin{aligned}
 (\mu\mu)_+ &= {}^3P_0 & C = P = CP = +1 \\
 (\mu\mu)_- &= {}^1S_0 & C = +1 \quad P = CP = -1
 \end{aligned}
 \tag{3}$$

For any $\pi\pi$ or $\gamma\gamma$ state $C = +1$, $CP = P$. Therefore parity conservation alone implies that CP is conserved in the transitions :

$$\pi\pi \rightarrow \mu\mu \quad \pi\pi \rightarrow \gamma\gamma \quad \gamma\gamma \rightarrow \mu\mu$$

Furthermore quantum electrodynamics for muons insures that CP is conserved in the process $\pi\pi\gamma \rightarrow \mu\mu$ (for total angular momentum zero) since in the order relevant to our discussion hadrons are involved only through the annihilation of two on-shell pions to a virtual photon. Current conservation and hermiticity imply that this process is CP conserving.

Therefore, if we neglect 3π intermediate states and $\pi\pi\gamma$ states ⁴⁾ except for the one arising from pion bremsstrahlung only the CP odd two-photon state $(\gamma\gamma)_-$ contributes to the absorptive part of $K \rightarrow (\mu^+\mu^-)_-$ and $K \rightarrow (\gamma\gamma)_-$ has no absorptive part.

Adopting the usual conventions we define even and odd eigenstates of the CP operator, K_1 and K_2 respectively. The phase convention is chosen so that

$$K_2 \rightarrow 2\pi \quad (I=0) \tag{4}$$

If we neglect the quantity ϵ' which measures the amplitude $K_2 \rightarrow 2\pi$ ($I=2$) and is known to be small ⁵⁾ then

$$K_2 \rightarrow 2\pi, \quad 2\pi\gamma \text{ (Bremsstrahlung)} \tag{5}$$

Then only the intermediate state $(\gamma\gamma)_+$ will contribute to the absorptive part of $K_2 \rightarrow (\mu\mu)_+$ and $K_2 \rightarrow (\gamma\gamma)_+$ has no absorptive part.

From Eq. (2) we thus obtain the following constraints :

$$\text{Im } T_{2\mu_-} = \frac{1}{2} \text{Re} (T_{2\gamma_-} T_{\gamma\mu_-}^*) = T_{2\gamma_-} \sqrt{\varphi_-} \quad (6a)$$

$$\text{Re } T_{1\mu_-} = -\frac{1}{2} \text{Im} (T_{1\gamma_-} T_{\gamma\mu_-}^*) = +i T_{1\gamma_-} \sqrt{\varphi_-} \quad (6b)$$

$$\text{Im } T_{2\mu_+} = -\frac{1}{2} \text{Im} (T_{2\gamma_+} T_{\gamma\mu_+}^*) = i T_{2\gamma_+} \sqrt{\varphi_+} \quad (6c)$$

where $T_{2\mu_-} = T_{K2} \rightarrow (\mu\mu)_-$, $T_{\gamma\mu_-} = T(\gamma\gamma)_- \rightarrow (\mu\mu)_-$, etc. The T matrix elements are understood to contain the relevant phase space factors so that their absolute squares give the partial widths and the $\varphi_{\pm}$ factors are obtained from explicit calculation ^{3),4)} of $\gamma\gamma \rightarrow \mu^+\mu^-$

$$\begin{aligned} \varphi_- &\approx 1.2 \times 10^{-5} \\ \varphi_+ &\approx 0.8 \varphi_- \end{aligned} \quad (7)$$

the remaining relation

$$\text{Re } T_{1\mu_+} = \frac{1}{2} \text{Re} (\sum_n T_{2n} T_{n\mu_+}^*) \quad (8)$$

is not useful since $\pi\pi$ and $\pi\pi\gamma$ states contribute here.

3. - CONSEQUENCES FOR $K \rightarrow \mu\mu$

To obtain constraints for K_L and K_S decays we use the relations :

$$K_S = K_1 + \varepsilon K_2 \quad (9a)$$

$$K_L = K_2 + \varepsilon K_1 \quad (9b)$$

$$\varepsilon \approx 2 \times 10^{-3} e^{i\pi/4}$$

Note that $\mathcal{E}$ cannot be neglected since replacing K_2 by K_L in Eqs. (6) gives

$$\Gamma_{K_L \rightarrow \mu\mu} \approx |T_{2\mu-}|^2 + |T_{2\mu+}|^2 \geq |\text{Im} T_{2\mu-}|^2 + |\text{Re} T_{2\mu+}|^2 = \varphi_- |T_{2\gamma-}|^2 + \varphi_+ |T_{2\gamma+}|^2 \quad (10)$$

Then since $\varphi_+ < \varphi_-$ [cf. Eq. (7)]

$$\Gamma_{K_L \rightarrow \mu\mu} \geq \varphi_+ \Gamma_{K_L \rightarrow \gamma\gamma} \approx 4.8 \times 10^{-9} \Gamma_L \quad (11)$$

in contradiction with the experimental bound ²⁾

$$\Gamma_{K_L \rightarrow \mu\mu} \leq 1.8 \times 10^{-9} \Gamma_L \quad (12)$$

For this reason the contribution of the $\pi\pi$ state to $K_L \rightarrow \mu\mu$ through the admixture of K_1 in K_L is not a priori negligible.

Next we note that with our assumptions (neglect of the 3π and direct emission $\pi\pi\gamma$ states) super weak theory is in contradiction with experiment : this theory predicts

$$K_2 \nrightarrow (\mu\mu)_+, (\gamma\gamma)_+ \quad (13)$$

$$K_1 \nrightarrow (\mu\mu)_-, (\gamma\gamma)_-$$

Then there is no interference between K_1 and K_2 in K_L decay, and we have

$$\Gamma_{K_L \rightarrow \mu\mu} = |T_{2\mu-}|^2 + |\mathcal{E} T_{1\mu+}|^2 \leq 1.8 \times 10^{-9} \Gamma_L \quad (14)$$

Equation (6a) together with Eq. (13) gives

$$\varphi_- |T_{2\gamma-}|^2 \leq |T_{2\mu-}|^2 \leq 1.8 \times 10^{-9} \Gamma_L \quad (15)$$

Taking out the factor φ_- in Eq. (15) and separating $T_{K_L \rightarrow \gamma\gamma}$ into contributions from K_1 and K_2 we obtain the inequality

$$|T_{2\gamma}|^2 = \Gamma_{K_L \rightarrow \gamma\gamma} - |\varepsilon T_{1\gamma}|^2 \leq 1.5 \times 10^{-4} \Gamma_L \quad (16)$$

From the rate for $K_L \rightarrow \gamma\gamma$:

$$\Gamma_{K_L \rightarrow \gamma\gamma} = 5 \times 10^{-4} \Gamma_L \quad (17)$$

we then obtain a lower bound for $K_S \rightarrow \gamma\gamma$:

$$\Gamma_{K_S \rightarrow \gamma\gamma} = |T_{1\gamma}|^2 + |\varepsilon T_{2\gamma}|^2 \geq 3.5 \times 10^{-4} \Gamma_L / |\varepsilon|^2 \approx 0.14 \Gamma_S \quad (18)$$

whereas the experimental limit is much lower ⁶⁾

$$\Gamma_{K_S \rightarrow \gamma\gamma} / \Gamma_S \leq 1.6 \times 10^{-3} \quad (19)$$

4. - GENERAL CASE

Using Eq. (9b) for K_L we obtain from Eqs. (6) :

$$\begin{aligned} \text{Im } T_{L\mu} = T_{2\gamma} \sqrt{q_-} + \text{Im}(\varepsilon T_{1\mu}) = T_{2\gamma} \sqrt{q_-} + \text{Im} \varepsilon i T_{1\gamma} \sqrt{q_-} + \\ + \text{Re} \varepsilon \text{Im } T_{1\mu} \end{aligned} \quad (20)$$

For $K_L \rightarrow (\gamma\gamma)_-$ we have the relation

$$\begin{aligned} T_{L\gamma} = T_{2\gamma} + \varepsilon T_{1\gamma} = T_{2\gamma} + i \text{Im} \varepsilon T_{1\gamma} + \text{Re} \varepsilon T_{1\gamma} \\ = T_{2\gamma} + i \text{Im} T_{1\gamma} - i \text{Re} \varepsilon \text{Re} T_{1\gamma} / \sqrt{q_-} \end{aligned} \quad (21)$$

so that Eq. (19) may be written

$$\sqrt{Q_-} T_{L\gamma_-} = \text{Im} T_{L\mu_-} - \text{Re} \epsilon (\text{Im} T_{1\mu_-} + i \text{Re} T_{1\mu_-}) \quad (22)$$

This is identical to the result of Christ and Lee for the CP odd state since only the two-photon state can contribute to the absorptive part.

The relation for the CP odd state is less restrictive ; from Eqs. (6) we have :

$$iT_{L\gamma_+} \sqrt{Q_+} = \text{Re} T_{L\mu_+} - \text{Re} (\epsilon T_{1\mu_+}) = i (T_{L\gamma_+} - \epsilon T_{1\gamma_+}) \sqrt{Q_+} \quad (23)$$

Following Christ and Lee, we consider Eqs. (21) and (23) as a relation between complex vectors :

$$x - y = z \quad (24)$$

with

$$\begin{aligned} x &= (\text{Im} T_{L\mu_-}, \text{Re} T_{L\mu_+}) \quad |x|^2 \leq K_{L \rightarrow \mu\mu} \\ y &= (i \text{Re} \epsilon T_{1\mu_-}, \text{Re} (\epsilon T_{1\mu_+})) \quad |y|^2 \leq \Gamma_{K_1 \rightarrow \mu\mu} |\epsilon|^2 \\ z &= (\sqrt{Q_-} T_{L\gamma_-}, i \sqrt{Q_+} (T_{L\gamma_+} - \epsilon T_{1\gamma_+})) \end{aligned} \quad (25)$$

The triangular inequality $|z| \leq |x| + |y|$ gives the inequality for $K \rightarrow \mu\mu$ rates :

$$\{Q_- |T_{L\gamma_-}|^2 + Q_+ |T_{L\gamma_+} - \epsilon T_{1\gamma_+}|^2\}^{1/2} \leq |\epsilon| \sqrt{\Gamma_{K_2 \rightarrow \mu\mu}} + \sqrt{\Gamma_{K_L \rightarrow \mu\mu}}$$

where we have replaced ϵT_1 by ϵT_S since $\epsilon^2 T_L$ is negligible compared with T_L .

Since [cf. Eqs. (17) and (19)]

$$|\varepsilon T_{S\delta+}|^2 \leq 6.4 \times 10^{-9} \Gamma_S \approx 4.1 \times 10^{-6} \Gamma_L \ll \Gamma_{K_L \rightarrow \gamma\gamma} \quad (26)$$

the worst limit on $\Gamma_{K_S \rightarrow \mu\mu}$ occurs if both K_S and K_L decay only into the $(\gamma\gamma)_+$ state. In that case we obtain :

$$|\varepsilon|^2 \Gamma_{K_S \rightarrow \mu\mu} \geq (\sqrt{39} \Gamma_L \times 10^{-5} - \sqrt{\Gamma_{K_L \rightarrow \mu\mu}})^2 \geq 4 \times 10^{-10} \Gamma_L \quad (27)$$

or

$$\Gamma_{K_S \rightarrow \mu\mu} \geq 10^{-4} \Gamma_L \approx 1.6 \times 10^{-7} \Gamma_S \quad (28)$$

This corresponds to maximal K_S, K_L interference. If there is no interference in the CP even $\gamma\gamma$ state, the contribution of $\Gamma_{K_S \rightarrow \gamma\gamma}$ to the inequality (24) is negligible and we obtain

$$\Gamma_{K_S \rightarrow \mu\mu} \geq \frac{1}{|\varepsilon|^2} (\sqrt{48} \Gamma_L \times 10^{-5} - \sqrt{\Gamma_{K_L \rightarrow \mu\mu}})^2 \geq 2.8 \times 10^{-7} \Gamma_S \quad (29)$$

Finally, if the $\pi\pi$ and $\pi\pi\gamma$ contributions are ignored altogether, Eq. (22) takes the form of Eq. (21) and the bound (28) is improved by a factor of two (independently of whether or not interference occurs in the $\gamma\gamma$ decay) because it is $(\text{Re } \varepsilon)^2 \Gamma_{K_S \rightarrow \mu\mu}$ rather than $|\varepsilon|^2 \Gamma_{K_S \rightarrow \mu\mu}$ which is limited. This is just the result of Christ and Lee.

5. - CONCLUSIONS

The question studied here is whether the discrepancy between the theoretical lower bound and the experimental upper bound for $K_L \rightarrow \mu\mu$ can be resolved without introducing an unexpectedly large contribution of the 3π state to the absorptive part of the amplitude (the $\pi\pi\gamma$ contribution can be calculated with some confidence and is found to be small ⁴⁾).

The answer implied by the above discussion is yes, provided that :

- (a) super weak theory is wrong and there is a significant direct CP violation in $K^0 \rightarrow (\mu^+\mu^-)$ and/or $(\gamma\gamma)$, as suggested in Ref. 3), and,
- (b) the rate for $K_S \rightarrow (\mu\mu)$ is much larger than the rate for $K_L \rightarrow (\mu\mu)$. We must have either

$$9 \times 10^4 \gtrsim \Gamma_{K_S \rightarrow \mu\mu} / \Gamma_{K_L \rightarrow \mu\mu} \gtrsim 5.5 \times 10^4 \quad (30)$$

and a large K_S, K_L interference in the two photon decay, or

$$\Gamma_{K_S \rightarrow \mu\mu} / \Gamma_{K_L \rightarrow \mu\mu} \gtrsim 9 \times 10^4 \quad (31)$$

An experimental limit on the $K_L K_S$ interference term would push the bound for $K_S \rightarrow \mu\mu$ towards the higher value.

We take pleasure in thanking Professors J.S. Bell and J. Prentki for lively discussions.

REFERENCES

- 1) G. Quigg and J.D. Jackson - UCRL Report 18487 (unpublished) ;
L.M. Sehgal - Nuovo Cimento 45, 785 (1966) ; Phys.Rev. 183,
1511 (1969).
- 2) A.R. Clark, T. Elliott, R.C. Field, H.R. Frisch, R.R. Johnson,
L.T. Kirth and W.A. Wenzel - Phys.Rev.Letters (to be
published).
- 3) M. Christ and T.D. Lee - Columbia University Preprint NYO-1932(2)-
195 (1971).
- 4) For estimates of these contributions, see :
B.R. Martin, E. de Rafael and J. Smith - Phys.Rev. D2, 179 (1970) ;
G.R. Farrar and S.B. Treiman - Phys.Rev.Letters (to be published) ;
M.K. Gaillard - CERN Preprint TH. 1327 (1971) ;
Ref. 3) above.
- 5) Experiments show $\epsilon' < 0.2 \epsilon$. See, e.g., :
J.M. Gaillard - Proceedings Daresbury Study Week-End on K Decay (1971).
- 6) B. Wolff, J.C. Chollet, J.M. Gaillard, M. Jane, J.P. Repellin and
K. Schubert - Private communication and reported by
P. Darriulat, Conférence sur la Physique des Hautes
Energies, Aix en Provence, Supplément au Journal de
Physique, Fasc. 11-12, C6 (1970).