***D*-branes, string cosmology, and large extra dimensions**

Antonio Riotto

Theory Division, CERN, CH-1211, Geneva 23, Switzerland

(Received 30 April 1999; published 10 May 2000)

D-branes are fundamental in all scenarios where there are large extra dimensions and the string scale is much smaller than the four-dimensional Planck mass. We show that this current picture predicts a period of inflation on our brane which is driven by the large extra dimensions and the issue of the graceful exit becomes inextricably linked to the problem of the stabilization of the extra dimensions, suggesting the possibility of a common solution. We also show that branes may violently fluctuate along their transverse directions in curved spacetime, possibly leading to a period of brane-driven inflation. This phenomenon plays also a crucial role in many other cosmological issues, such as the smoothing out of the cosmological singularities.

PACS number(s): 98.80.Cq, 04.50.+h, 11.10.Kk, 11.25.Mj

D-branes are classical solutions which are known to exist in many string theories [1]. A *Dp*-brane denotes a configuration which extends along *p* spatial directions and is localized in all other spatial transverse directions. From the string point of view, a *D*-brane is a soliton on which open string endpoints may live and whose mass (or tension) is proportional to the inverse of the string coupling g_s . This means that *D*-branes become light in those regions of the moduli space where the string coupling is large. This simple property has led to a new interpretation for singularities of various fixed spacetime backgrounds in which strings propagate and provided a generic and physically intuitive mechanism for smoothing away cosmological singularities in the strong coupling or large curvature regime of the evolution of the early Universe [2,3].

The presence of branes in string theory has also given rise to the possibility that the standard model (SM) gauge fields live on branes rather than in the bulk of spacetime. This scenario corresponds to a novel compactification in which gravity lives in the bulk of the ten dimensional spacetime, but the other observed fields are confined to a brane of lower dimension [4]. In particular, one might imagine the situation where the SM particles live on a three brane and the directions transverse to the three brane being compact. There might be also other branes separated from ours in the extra directions. Since the usual gauge and matter fields live on the brane, the only long-range interaction which sees the extra dimensions is gravity.

In the weakly coupled heterotic string there is a fundamental relation between the string scale M_s , the four-dimensional Planck mass M_p , the value of the dilaton field ϕ and the gauge coupling constant α_g , $(M_s/M_p)^2 \sim \alpha_g \sim e^\phi$ and the string scale is fixed to be about 10^{18} GeV. However, in other regions of the moduli space the string scale can be much smaller [4] and it is tempting to imagine [5] that the string scale is not far from the TeV region. If this is the case, the small ratio M_s/M_p could be a consequence of large compactified dimensions [6,7] through a scaling relation like $M_p^2 = M_s^{n+2} V_n$, where V_n is the volume of the *n* large extra dimensions. For $M_s \sim \text{TeV}$ and $n \geq 2$, the radius of the large extra dimensions may be as large as the millimeter. Large extra dimensions are only allowed if the SM

fields are confined to live on a membrane orthogonal to the large extra dimensions and this hypothesis fits rather nicely with the current picture of string theory and *D*-branes.

Not surprisingly, one expects many interesting cosmological implications of this scenario (some of them have already been discussed in [2,8]) and the possibility of large extra dimensions is potentially exciting for several reasons.

In the presence of some matter with energy density ρ , the evolution of the Universe is controlled by the equation $H^2 \equiv (\dot{a}/a)^2 = (8\pi\rho/3M_p^2)$, where a is the scale factor and H is the Hubble rate. This implies that for an expanding Universe, ρ and H^2 are proportional, they increase or decrease together in time. This is true only if M_p is constant. Now, in the weakly coupled string theory where all compactified dimensions have size $\sim M_s^{-1}$, M_p is controlled by the dilaton and it is perfectly possible to have growing H while ρ decreases, provided the dilaton ϕ is also growing. On this peculiar feature is based the whole idea of pre-big-bang (PBB) cosmology where there exist superinflationary solutions for the scale factor a having a growing—rather than a constant—Hubble parameter and a growing dilaton field [9]. The suggestion from weakly coupled string theory is therefore rather compelling, since it is possible to generate a scenario containing a dilaton-driven PBB inflation.

These simple considerations suggest a simple way of getting inflation on our brane: since M_p is now governed by the size b of the large extra dimensions, it is possible to obtain superinflationary solutions provided that the extra dimensions change in time [10,11], that is a radion-driven PBB cosmology. The Einstein action in $4+n$ dimensions and metric $g_{MN} = (1, -a^2\delta_{ij}, -b^2\delta_{mn})$ reads

$$S \sim \int d^4x a^3 b^n \left[R - 2n \frac{\dot{b}}{b} - 6n \frac{\dot{a}}{a} \frac{\dot{b}}{b} - n(n-1) \frac{\dot{b}^2}{b^2} \right], \quad (1)$$

where R is the usual four-dimensional Ricci scalar. Setting $b^n = e^{c\sigma}$, with $c = (n/n-1)^{1/2}$, and integrating by parts the term $\dot{b}$ in Eq. (1), the action for the radion field σ becomes

$$S \sim \int d^4x a^3 e^{c\sigma} [R + \partial_\mu \sigma \partial^\mu \sigma]. \quad (2)$$

Were $c=1$, string cosmology experts would recognize the action of the dilaton field coupled to gravity (with $\phi = -\sigma$). For us $c=1$ in the limit of $n \rightarrow \infty$ and in this regime we can recover the more familiar results of dilaton-driven PBB cosmology. The evolution of the radion is governed by the equation obtained by varying the lapse function

$$\dot{\sigma} = \epsilon_{\pm} H, \quad \epsilon_{\pm} = -3c \pm \sqrt{9c^2 - 6}, \quad (3)$$

while the equation for the Hubble parameter becomes

$$\dot{H} = -\alpha_{\pm} H^2, \quad \alpha_{\pm} = 3 + c\epsilon_{\pm}. \quad (4)$$

For each value of n , α_{-} is negative and it generates a superinflationary solution, that is H grows until a singularity is reached, say at $t=0$. In terms of the scale factors a and b , the solutions are given by $a(t) = a_0(t/t_0)^{-\gamma_1}$ and $b(t) = b_0(t/t_0)^{\gamma_2}$ where t and t_0 are negative and $\gamma_1 = |1/\alpha_{-}|$, $\gamma_2 = |c\epsilon_{-}/(n\alpha_{-})|$. Since $\epsilon_{-} < 0$, the volume of the internal dimensions is contracting. Suppose that a and b start with comparable values at low curvature at $t \sim t_0$, say $a \sim b \sim M_s^{-1}$; the subsequent evolution is such that the spatial dimensions of our three brane superinflate, whereas the extra dimensions evolve to values smaller than M_s^{-1} . In string theory, due to the presence of winding states in the closed string (gravitational) sector, the low-energy physics is equivalent as if there was a radius $b_I = (bM_s^2)^{-1}$ which is much larger than the string length and the roles of winding and Kaluza-Klein momenta are interchanged [by performing a T-duality transformation $\lambda \rightarrow \lambda/(bM_s)$, the dimensionless coupling $\lambda = 8/\alpha_g(M_s/M_p)^2$ goes to $\lambda = 4\alpha_g$, which is still in the perturbative regime]. On the other hand, in this T -dual description, the open string states—which give rise to the ordinary non gravitational matter on the three brane—have only heavy winding modes. As a result, in the dual theory the extra dimensions appear to be large. The dynamics of the radion field b coupled to the four-dimensional gravity leads automatically to an inflationary stage on our three brane and explains the huge hierarchy between the sizes of our observed Universe and of the extra dimensions if $|\gamma_1| \gg |\gamma_2|$. Inflating the observable dimensions by the required amount depends upon the parameters, in particular the number of extra dimensions. Similarly, the amount of density perturbations generated during the inflationary stage will depend crucially on the stringy scale M_s and the number of extra dimensions. This is a crucial issue deserving a careful analysis which is beyond the scope of this paper.

The cosmological solution stops being valid when the Hubble rate becomes of the order of M_s . However, differently from the dilaton-driven PBB cosmology where one has to face a regime of strong coupling when approaching the big-bang singularity at $t \rightarrow 0$, in the radion-driven PBB scenario the singularity may take place well inside the perturbative regime and be understood as the process of decompactifying the extra dimensions. The issue of the graceful exit in string cosmology becomes therefore inextricably linked to the problem of the stabilization of the extra dimensions, suggesting the possibility of a common solution.

There is—however—another feature of D -branes that may help in solving the radion-driven PBB singularity problem as well as give new insight into other fundamental cosmological issues: this is the peculiar behavior of fluctuations transverse to the brane in curved spacetime. The action governing the dynamics of a generic Dp -brane (neglecting the gauge fields) is [1] $S = -T_p \int d^{p+1}\xi e^{-\phi} \sqrt{-\det G_{\alpha\beta}}$, where T_p^{-1} is the brane tension, ξ^{α} ($\alpha = 0, \dots, p$) parametrize the D -brane world-volume and $G_{\alpha\beta} = g_{\mu\nu}(X) \partial_{\alpha} X^{\mu} \partial_{\beta} X^{\nu}$ is the induced metric. Given a brane configuration $\bar{X}^{\mu}(\xi)$ in a spacetime of D -dimensions, arbitrary fluctuations transverse to this configuration (the would-be Goldstone bosons of the broken translational invariance along the directions perpendicular to the brane) are parametrized by $(D-p)$ world-volume scalars $y^a(\xi)$ as $X^{\mu}(\xi) = \bar{X}^{\mu}(\xi) + \bar{n}_a^{\mu}(\xi) y^a(\xi)$, where $\bar{n}_a^{\mu}(\xi)$ [$a = 1, \dots, (D-p)$] are the $(D-p)$ vectors normal to the brane at the point $\bar{X}(\xi)$. The expansion of the metric in Riemann coordinates is [12] $g_{\mu\nu}(X) = \bar{g}_{\mu\nu} - \frac{1}{3} y^a y^b \bar{R}_{\mu\rho\nu\sigma} \bar{n}_a^{\rho} \bar{n}_b^{\sigma} + \mathcal{O}(y^3)$, where the overbar indicates that the quantity is evaluated at $\bar{X}$. If the extrinsic curvature of the membrane is small, the normal vectors are $\bar{n}_a^{\rho} \approx \delta_a^{\rho}$, and therefore $g_{\mu\nu}(X) = \bar{g}_{\mu\nu} - \frac{1}{3} y^a y^b \bar{R}_{\mu a \nu b} + \mathcal{O}(y^3)$ [2].

Let us begin, for simplicity, by considering a single Dp -brane in an expanding accelerating ($\ddot{a} > 0$) background where the dimensions orthogonal to the brane are static. We shall consider for definiteness a background where the common scale factor a of the spatial coordinates $\vec{\xi}$ is inflating, $a(\eta) \sim (-\eta)^{-q}$, where $0 < q \leq 1$ and $d\eta = (dt/a)$ is the conformal time, $-\infty < \eta < 0$. For $q \neq 1$ the Hubble rate H grows with time till a singularity is reached and the De Sitter background with constant H is recovered for $q = 1$. The equation of motion for small transverse oscillations $y_a(\xi)$ (y_a is the comoving displacement) is

$$\ddot{y}_a + (p-1)(\dot{a}/a)\dot{y}_a - \nabla^2 y_a = 0, \quad (5)$$

where $\dot{y}_a \equiv \partial y_a / \partial \eta$. The canonical conjugate momentum is $\pi_a = \tau_p a^{p-1} \dot{y}_a$, where $\tau_p = T_p / g_s$; imposing the canonical commutation relation $[y_a(\eta, \vec{\xi}), \pi_b(\eta, \vec{\xi}')] = i \delta^{(p)}(\vec{\xi} - \vec{\xi}')$, we can expand the solution as

$$y^a(\eta, \vec{\xi}) = \frac{1}{\tau^{1/(p+1)}} \int \frac{d^p k}{(2\pi)^{p/2}} [a_k h_k^a(\eta) e^{-i\vec{k} \cdot \vec{\xi}} + a_k^{\dagger} h_k^{a\dagger}(\eta) e^{i\vec{k} \cdot \vec{\xi}}], \quad (6)$$

where a_k and $a_k^{\dagger}$ are the annihilation and creation operators, respectively. The solution is very simple, $h_k^a(\eta) = (-\eta)^{\beta} [c_1 H_{\beta}^1(-k\eta) + c_2 H_{\beta}^{1*}(-k\eta)]$, where $\beta = (q(p-1)+1)/2$. If we quantize the modes and define the vacuum state by $a_k|0\rangle = 0$, then different choices of vacuum correspond to different choices of c_1 and c_2 . The adiabatic vacuum corresponds to $c_2 = 0$ and this (Heisenberg) state is the state we will assume the brane is in.

The behavior of a mode $h_k^a(\eta)$ is very simple. Suppose from now on to consider the case $p=3$, i.e., we consider a three brane. As long as the physical wavelength ak^{-1} of a mode is inside the Hubble radius, $h_k^a(\eta)$ oscillates with constant *physical* amplitude $ah_k^a(\eta)$. As the physical wavelength grows it crosses the Hubble radius and the *comoving* amplitude $h_k^a(\eta)$ becomes frozen, so that the physical amplitude grows like a . In this state one can calculate the mean square transverse displacement

$$\langle \mathbf{y}^2 \rangle \equiv \int \frac{d^3 \xi}{V_3} \langle 0 | \mathbf{y}^2(\eta, \xi) | 0 \rangle = \frac{D-4}{\tau^{1/2}} \int \frac{d^3 k}{(2\pi)^3} |h_k^a|^2, \quad (7)$$

where $\mathbf{y}^2 \equiv \sum_a y_a^2$ and $V_3 = \int d^3 \xi$. The physical displacement $a\mathbf{y}$ has the usual ultraviolet divergence as in flat space: physical wavelengths with $k \gg aH$ are always well inside the horizon at a given time and their amplitude is unaffected by the expansion. Their contribution leads to the usual $\int d^3 k k^{-1}$ divergence. We can subtract this divergence. Modes with $k \ll a_i H_i$, where a_i and H_i are the values of the scalar factor and Hubble rate at the beginning of the inflationary stage, are always well outside the horizon and simply match on adiabatically to the modes before and after inflation. One therefore finds at $\eta \approx \eta_f \approx 0$ (for $q \neq -1$)

$$a^2 \langle \mathbf{y}^2 \rangle \approx a^2 \frac{D-4}{\tau^{1/2}} \int_{a_i H_i}^{a_f H_f} H \frac{d^3 k}{(2\pi)^3} |h_k^a|^2 \sim \frac{D-4}{M_s^4 \eta_f^2}, \quad (8)$$

where we have set $\tau \sim M_s^4$. One can picture this result by saying that the modes with wavelength of the order of the Hubble radius have a physical width $\langle \mathbf{y}_p^2 \rangle \approx M_s^{-2}$ which gets amplified by the expansion after they cross the Hubble radius. Transverse fluctuations grow in time. In the special case of de Sitter with constant rate of expansion H , one finds that $\langle \mathbf{y}^2 \rangle \approx M_s^{-4} H^2 \ln(EH) \sim M_s^{-4} H^3 t$, where E is the total number of e -foldings.

More interestingly, one can calculate the energy acquired by each mode in this process. The energy is given by $E \sim M_s^4 \int d^3 \xi a^3 [1 + 1/2 \dot{\mathbf{y}}^2/a^2 + 1/2 (\nabla \mathbf{y})^2/a^2]$, where the first term is just the classical stretching. Just as in (8), we find

$$\frac{\langle (\nabla \mathbf{y})^2 \rangle}{a^2} \approx \left(\frac{H}{M_s} \right)^4. \quad (9)$$

The term $\langle \dot{\mathbf{y}}^2 \rangle$ gives no contribution after the flat spacetime subtraction. Thus we deduce that the fractional energy in the transverse perturbations grows with time as H increases with time and their contribution to the energy density becomes sizeable when $H \sim M_s$. Physical transverse fluctuations of the three brane—even though initially oscillating—will eventually grow and give rise to an instability. Therefore, as the three brane is stretched by the expansion of the Universe, transverse fluctuations grow, the thickness of the brane gets larger and larger and the brane cannot be considered as a static object any longer. This effect resembles the instability and growth of the string modes in accelerated backgrounds [11,13].

This growth is accompanied by a huge production of light pseudo-Goldstone bosons of the broken translation invariance. Let us try to feed back the effect of this production into the expansion of the Universe. As the system evolves towards the singularity, more and more energy is transferred into transverse fluctuations of the brane till—eventually—they dominate the energy density of the Universe. Since $\langle (\nabla \mathbf{y})^2 \rangle \gg \langle \dot{\mathbf{y}}^2 \rangle$, the gas of pseudo-Goldstone bosons is characterized by an energy density $\rho_y \approx -3p_y$, where p_y is the pressure of the gas. We thus obtain an effective negative pressure. Assuming a flat Universe, the equation for the acceleration $\ddot{a}$ alone becomes $\ddot{a}/a \propto (\rho_y + 3p_y) = 0$ and we discover that the back reaction of the gas of pseudo-Goldstone bosons onto the evolution of the Universe is to halt the period of acceleration. This effect again reminds the amplification of string fluctuations in accelerated backgrounds [11,13].

It should be stressed that this result is only valid when the stress tensor is of the perfect fluid type, i.e. when it becomes possible to neglect the viscosity terms due to mutual and self-interactions. These terms are not negligible when the approximation of small transverse fluctuations breaks down. The role played by the transverse fluctuations may be even more dramatic, though. Remember that in the considerations made so far we have supposed that the directions transverse to our three brane were static. Assume now that these directions are suffering a period of decelerating contraction, while the spatial dimensions of the brane are accelerating. This is—for instance—what happens during the stage of radion-driven inflation previously described. The fluctuations y_a transverse to the brane get a tachyonic mass $m_y^2 = 1/3(\dot{b}/b + 3\dot{b}/b\dot{a}/a)$, where now the dots stand for derivatives with respect to the cosmic time t . The situation is very similar to the thermal tachyon which occurs at the Hagedorn temperature in string theory [14,15]. In that case a condensate of winding states forms when the periodic imaginary times becomes too small. In the present case the tachyon forms because of the nontrivial geometry of the spacetime. As a given physical wavelength ak^{-1} of a transverse fluctuation y_a crosses the horizon, the comoving amplitude h_k^a does *not* freeze out; instead the transverse physical amplitude grows *faster* than the scale factor a and therefore faster than the physical coordinates defining the brane. In fact the brane tends to lose its own identity as the thickness of the brane grows faster than the physical coordinates defining the three brane itself; the brane violently fluctuates along the transverse directions. This phenomenon holds for branes of any dimensionality. If one goes beyond the approximation of small transverse fluctuations and computes the higher order terms in the tachyon effective potential, one can show that—along with these huge fluctuations—branes become almost tensionless [2]. This is hardly surprising since the brane can fluctuate along the transverse directions paying no price in energy. When the tension becomes negative, branes become unstable and—at least in the regime of strong coupling—it becomes easier and easier for the gravitational background to produce further massless branes. On the other hand, when the transverse sizes of the branes become larger than the horizon, the

branes presumably break down and decay, but branes are continuously created. The rate of brane creation due to these quantum effects may be so fast to balance the dilution of the brane density due to the expansion. One might be therefore led into a phase of *constant* brane density and exponentially expanding Universe, phase of *brane-driven* inflation [16] reminiscent of the old idea of string-driven inflation [17]. The huge transverse fluctuations of branes in curved space-time may be also relevant to the issue of baryon asymmetry production on our three brane if we accept the idea that there might be other branes separated from ours in the extra directions. Even though on our brane global charges, such as the baryon and the lepton number, are conserved with a great accuracy, this is not necessarily the case on other branes. This breaking might be communicated to our brane by messenger fields that leave in the bulk [18]. However, this communication is suppressed by the enormous distance separating the branes in the bulk and this is the reason why today the baryon number is conserved in SM interactions. Nevertheless, at early epochs in the evolution of the Universe the thickness of a brane is highly fluctuating. This makes it possible for two branes separated in the bulk to overlap. One can also envisage various other dynamical phenomena like the melting of two or more branes [2]. If our brane was considerably overlapping with another brane where the breaking on the baryon number is $\mathcal{O}(1)$, a significant amount of baryon number might have been deposited on our three brane, thus

explaining the observed baryon asymmetry of our Universe [19].

The considerations reported here are very preliminary and certainly leave many questions unanswered. For instance, what are the initial conditions for the Universe (or perhaps just for our region of the Universe) right before the onset of inflationary PBB stage driven by the radion field? What is the subsequent dynamics of the gas of pseudo Goldstone bosons which is generated at $H \sim M_s$ and whose presence may help in avoiding the singularity? How does the system get into the familiar radiation-dominated stage? At this point open strings attached to the branes have to play a role since the expansion time H^{-1} is of the order of the typical scale of the stringy network M_s^{-1} . It seems safe therefore to assume that this gas of strings attached to the branes and pseudo Goldstone bosons reaches thermal equilibrium. The fact that fundamental strings have a limiting temperature $\sim M_s$ may turn out to be crucial since it implies that if radiation is created and is in thermal contact with the strings, it cannot attain a density higher than $\sim M_s^4$, and the Universe necessarily is dominated by strings.

We are grateful to T. Banks, R. Brustein, E. Copeland, K. Dienes, M. Dine, S. Dimopoulos, T. Gherghetta, G. F. Giudice, D. Lyth, R. Rattazzi, G. Veneziano, and especially to M. Maggiore for many useful discussions.

-
- [1] See, for instance, J. Polchinski, “TASI Lectures on D-branes,” hep-th/9611050.
 - [2] M. Maggiore and A. Riotto, Nucl. Phys. **B548**, 427 (1999).
 - [3] T. Banks, W. Fischler, and L. Motl, J. High Energy Phys. **01**, 019 (1999).
 - [4] E. Witten, Nucl. Phys. **B471**, 135 (1996).
 - [5] J. Lykken, Phys. Rev. D **54**, 135 (1996).
 - [6] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Lett. B **429**, 263 (1998).
 - [7] I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, Phys. Lett. B **436**, 257 (1998).
 - [8] N. Arkani-Hamed, S. Dimopoulos, and G. Dvali, Phys. Rev. D **59**, 086004 (1999); K.R. Dienes, E. Dudas, T. Gherghetta, and A. Riotto, Nucl. Phys. **B543**, 387 (1999); N. Arkani-Hamed, S. Dimopoulos, and J. March-Russell, Phys. Rev. D (to be published), hep-th/9809124; K. Benakli and S. Davidson, *ibid.* **60**, 025004 (1999); D.H. Lyth, Phys. Lett. B **448**, 191 (1999); N. Kaloper and A. Linde, Phys. Rev. D **59**, 101303 (1999); G. Dvali and S.-H. H. Tye, Phys. Lett. B **450**, 72 (1999); T. Banks, M. Dine, and A. Nelson, J. High Energy Phys. **06**, 014 (1999); N. Arkani-Hamed *et al.*, hep-ph/9903224; G. Dvali and G. Gabadadze, Phys. Lett. B **460**, 47 (1999); G. Dvali and M. Shifman, Phys. Rep. **320**, 107 (1999).
 - [9] G. Veneziano, Phys. Lett. B **265**, 287 (1991); M. Gasperini and G. Veneziano, Astropart. Phys. **1**, 317 (1993).
 - [10] D. Sahdev, Phys. Lett. **137B**, 155 (1984).
 - [11] M. Gasperini, N. Sanchez, and G. Veneziano, Nucl. Phys. **B364**, 365 (1991).
 - [12] See, for instance, A.Z. Petrov, *Einstein Spaces* (Pergamon, Oxford, 1969).
 - [13] M. Gasperini, N. Sanchez, and G. Veneziano, Int. J. Mod. Phys. A **6**, 3853 (1991).
 - [14] B. Sathiapalan, Phys. Rev. D **35**, 3277 (1987).
 - [15] Ya.I. Kogan, Pis'ma Zh. Éksp. Teor. Fiz. **72**, 1349 (1977) [JETP Lett. **45**, 709 (1987)].
 - [16] A. Riotto (unpublished).
 - [17] N. Turok, Phys. Rev. Lett. **60**, 549 (1988).
 - [18] N. Arkani-Hamed and S. Dimopoulos, hep-ph/9811353.
 - [19] See also Dvali and Gabadadze [8].