CERN LIBRARIES, GENEVA

SCAN-0010082

ОБЪЕДИНЕННЫЙ
ИНСТИТУТ
ЯДЕРНЫХ
ИССЛЕДОВАНИЙ

Дубна

E1-99-332

Z.U.Usubov*, O.B.Abdinov*

A SEARCH FOR QUARK COMPOSITENESS
AT THE LHC.
DIJET ANGULAR DISTRIBUTIONS

1. 4. 99
Submitted to «Ядерная физика»

Yadernaya Fizika
pod redaktsionnoy
Uchebnoye Fizika
*Institute of Physics, Baku, Azerbaijan

1999

The existence of an underlying substructure will be good key for understanding of the unresolved problems of Standard Model such as the presence of few generations of quarks and leptons, the fundamental properties of particles.

The quark substructure would appear as an excess of the high P_T jets compared to the level predicted by QCD or with dijet angular distributions more isotropic than what is expected from a point-like quark theory. Dijet angular distributions have been studied by the CDF [1] and $D0$ [2] experiments at a center-of-mass(CM) energy of $1.8 TeV$. The highest E_T reached so far at the *Tevatron*, $440 GeV$, corresponds to a distance scale $10^{-17} cm$. The experimental data have been compared with QCD predictions including compositeness. No evidence of quark substructure was found. Previous studies of dijet invariant mass spectrum reported by $UA1$ [3], $UA2$ [4] at $\sqrt{s} = 630 GeV$ and by CDF [5] have also shown the data that were consistent with QCD predictions. The CDF [6] and $D0$ [7] results on high-mass Drell-Yan cross-section measurement sets lower limit on the quark-electron compositeness scale about $5.5 TeV$. From the ratio of charged current to neutral current cross-section measurement in the CCFR fixed target neutrino experiment[8] at *Tevatron* limit on $\Lambda \sim 8 TeV$ was achieved.

In future hadron colliders, the search for quark substructure will continue. Here we investigate the effect of quark compositeness in dijet angular distribution as would be seen by $ATLAS$ [14] at the *LHC*. The same problem for high E_T jet spectrum was pointed out earlier[9]. To simulate a scenario with quark substructure the event generator $PYTHIA-5.7$ [10] has been used. This has allowed to use a simple phenomenological approach of contact interactions between quark constituents with a compositeness scale Λ [11], where the sign of the *effective Lagrangian* for a flavor diagonal definite chirality current is positive (destructive interference) or negative (constructive interference). The data simulated in the framework of *Standard Model (SM)* are compared with those obtained assuming quark compositeness. The analysis is based on a samples of ~ 280800 pp -interactions at $\sqrt{s} = 14 TeV$ which corresponds to the sample of dijet events expected after one month of *LHC* operation at a luminosity $\mathcal{L} = 10^{33} cm^{-2}s^{-1}$. The simulated event sample included the following hard-scattering subprocesses: $qq, qg, gg, g\gamma, q\gamma, \gamma\gamma$. The $\gamma^*/Z, W, tt$ production subprocesses also enabled. To get a sufficiently large

number of events with high P_T jets in a reasonable *CPU* time. a cut on the transverse momentum of the hard scattering subprocess was set to 600 GeV . Under these conditions, the contributions from the qq , qg and gg process represent 97.5% of the total cross section $3.370 \cdot 10^{-7}\text{ mb}$. Initial and final state QCD and QED radiation, fragmentation and decay of partons and particles, multiple interactions were enabled. First-order running α_S calculations were applied. The Λ_{QCD} value is chosen according to the parton distribution functions (pdf) parametrizations, used in *PYTHIA*. For the Q^2 scale in the hard scattering $2 \rightarrow 2$ process, $Q^2 = (m_{T1}^2 + m_{T2}^2)/2$ was used. The detector performance was simulated using the *ATLFAST*[12] package which provides a reliable estimate of the detector response to hadronic jets. Jets were reconstructed with *ATLFAST* using the standard procedure of summing the energy deposited in a cone of radius $\Delta R = \sqrt{\Delta\eta^2 + \Delta\phi^2} = 0.7$. All calorimeter cells with $E_T > 1.5\text{ GeV}$ are taken as possible initiators of clusters. The total E_T summed over all cells in a cone ΔR should be larger than 15 GeV . Jets are reconstructed down to $|\eta| \leq 5.0$.

The analysis was made in terms of an angular variable $\chi \equiv e^{(|\eta_1 - \eta_2|)}$, where $\eta_{1,2}$ are the pseudorapidities of the two leading jets. For the case of $2 \rightarrow 2$ parton scattering, it is related to the CM scattering angle Θ^* as follows:

$$\chi = \frac{1 + |\cos\Theta^*|}{1 - |\cos\Theta^*|} \quad (1)$$

This definition makes the comparison with theory more straightforward[13]. The investigation of the dijet angular distribution, $(1/N)(dN/d\chi)$, was made in four dijet invariant mass bins. The dijet invariant mass is defined as

$$M_{jj} = \sqrt{(E_1 + E_2)^2 - (P_1 + P_2)^2}, \quad (2)$$

where $(E_{1,2}, P_{1,2})$ are the 4-vector of the two leading jets. For all dijet invariant mass bins the E_T -threshold for the highest E_T jet was 400 GeV . *Table1* shows the selection cuts for the highest E_T jet for the various invariant dijet mass bins, together with the average M_{jj} and the number of events per bin.

The dijet angular distributions for these dijet mass bins are shown in *Figure 1*, and *Figure 2* for destructive and constructive interference, respectively. From the figures, one can see that quark compositeness leads to an enhancement in the distribution at low values of χ in comparison to

<i>Mass bins</i> (<i>GeV</i>)	2000-2300	2300-2800	2800-3400	>3400
<i>E_T thresholds</i> (<i>GeV</i>)	400	400	400	400
<i>N_{ev}</i>	18562	15781	7722	5228
<i>Average M_{jj}</i> (<i>GeV</i>)	2136	2512	3050	4048

Table 1: Characteristics of the invariant mass bins for the high E_T jets.

the SM prediction. The dijet mass range above 3400 GeV is good for the determination of isotropic contributions to the dijet angular distribution in pp -interactions at $\sqrt{s} = 14\text{ TeV}$ for Λ up to 8 TeV . One can see as well that the sensitivity is greater for a constructive interference than for destructive one.

To estimate the limits on the quark compositeness scale the CDF -group used the variable R_χ which is the ratio of the number of events with $\chi < \chi_0$ to the number with $\chi > \chi_0$. In our case a value of $\chi_0 = 5$ was used. *Figure 3* shows the R_χ as a function of the dijet mass for different values of the compositeness scale Λ , in the constructive case, when two, and all quarks are composite. The data are plotted at the average mass for each mass bin. In the case that all quarks are composite and for a compositeness scale $\Lambda = 16\text{ TeV}$ the data differ from the SM predictions by 1.5σ and 3.5σ , for destructive and constructive interference, respectively.

In *Figure 4* the dependence of R_χ on the scale Λ , for the constructive and destructive cases, and when two and all quarks are composite are compared. It is clear, that there is not enough sensitivity to distinguish the cases of two or all quarks compositeness.

R_χ is not very sensitive to the pdf as illustrated in *Figure 5*, where the values of R_χ are shown for the mass bin of M_{jj} above 3400 GeV . These predictions have been obtained for $\Lambda_{ud,all}^- = 8000\text{ GeV}$ for the cases when two or all quarks are composite. Note that in the rest of the analysis, *PYTHIA* was used with the default structure function *CTEQ2L*.

R_χ is also insensitive to the jet cone radius ΔR .

The sensitivity to the calorimeter resolution has been studied[15]. In *ATLFAST*, the jet energy is smeared according to $\sigma_E/E \sim 50\%/\sqrt{E} + 2\%$ in the central region ($|\eta| < 3$) and $\sigma_E/E \sim 100\%/\sqrt{(E)} + 7\%$ in the forward calorimeters ($3 < |\eta| < 5$).

In order to investigate the influence of a change of the constant term

on $(1/N)(dN/d\chi)$, we simply multiplied and divided the constant term by two. Changes in the stochastic were also considered. There is no significant impact of those changes on the dijet angular distribution.

The non-linear response of the hadron calorimeter can affect the observed difference between the SM and compositeness scenario, or fake a compositeness signal. To study this effect, we considered a non-linearity of the jet E_T scale according to the relation[16]:

$$E_T(meas) = E_T \cdot \frac{1}{c(1. + (e/h - 1.) \cdot 0.11 \cdot \ln E_T)} \quad (3)$$

where $E_T(meas)$ and E_T are measured and true jet transverse energy; $e/h = 1.36$ and c is adjusted such that at $50 GeV$ the scale is unchanged. Such a dependence on E_T leads to a deviation from linearity of 6.5, 9.3 and 12.3% for 400, 1000 and 3000 GeV , respectively. In *Figure 6* we compare the SM and composite quarks predictions with and without non-linearity effects. For this choice of dijet mass bin intervals and jet E_T , we see that no fake signal is created and that the angular distribution is quite insensitive to effects of non-linearity.

To study the data sensitivity to the quark compositeness signal for higher scale Λ_{all}^- , a analysis was performed for integrated luminosity of $30 fb^{-1}$ and $300 fb^{-1}$. *Figure 7* and *Figure 8* shows the difference between compositeness quark and SM predictions, divided by the SM predictions for dijet angular distributions.

In conclusion, the study based on *PYTHIA-5.7* generated and processed through *ATLFAST* shows that high mass dijet angular distribution has a good discovery capability for compositeness. Small sensitivity to the pdf choice, calorimeter energy resolution and non-linearity effects makes the dijet angular distribution a powerful tool for future high statistics data analyses. One month of LHC operation at $10^{33} cm^{-2} s^{-1}$ and $\sqrt{s} = 14 TeV$ allows the discovery of quark substructure if the constituent interaction constant is of the order $14 TeV$. To reach a lower limit of $25(40) TeV$ for interaction constant a data taking at integrated luminosity of $30(300) fb^{-1}$ is needed.

We would like to thank D. Froidevaux, F. Gianotti, I. Hinchliffe and M. Bosman for help and valuable comments.

Figure 1: Dijet angular distributions for various mass bins in case of destructive interference.

Figure 2: Dijet angular distributions for various mass bins in case of constructive interference.

Figure 3: R_χ as a function of the dijet mass M for different values of the compositeness scale Λ when two (a) and all quarks (b) are composite. The data are plotted at the average mass for each mass bin.

Figure 4: The dependence of R_χ on the scale Λ , for the constructive and destructive cases, when two and all quarks are composite.

Figure 5: The dependence of R_x on the parton distribution function.

Figure 6: Standard Model and composite quark predictions with and without non-linearity effects.

Figure 7: The difference between composite quark and Standard Model predictions, divided by the Standard Model for dijet angular distribution at integrated luminosity $L = 30 \text{ fb}^{-1}$.

Figure 8: The same as in **Figure 7**, but for an $L = 300 \text{ fb}^{-1}$.

References

- [1] CDF Collaboration, F. Abe et al., Phys.Rev.Lett. 77, 5336 (1996).
- [2] D0 Collaboration, B. Abbott et al., Phys.Rev.Lett. 80, 666 (1998).
- [3] UA1 Collaboration, C. Albajar et al., Phys. Lett. B209, 127 (1988).
- [4] UA2 Collaboration, J. Alitti et al., Z. Phys. C49, 17 (1991).
- [5] CDF Collaboration, F. Abe et al., Phys. Rev. D41, 1722 (1990),
Phys. Rev. Lett. 71, 2542 (1993).
- [6] CDF Collaboration, F.Abe et al., Phys.Rev.Lett. 79, 2198 (1997).
- [7] D0 Collaboration, B. Abbott et al., Fermilab-Pub-98/391-E.
- [8] CCFR Collaboration, K.S.McFarland et al., Fermilab-Pub-97/001-E.
- [9] Z.U. Usubov, Preprint JINR E1-99-298, Dubna, Russia, 1999.
- [10] T. Sjostrand, Preprint LU tp 95-20, Computer Physics Commun. 82,
74 (1994).
- [11] E. Eichten, K. Lane and M. Peskin, Phys. Rev. Lett. 50, 811 (1983).
E. Eichten et al., Rev. Mod. Phys. 56, 579 (1984).
- [12] E. Richter-Was, D. Froidevaux and L. Poggioli, ATLAS Internal Note,
PHYS-No-079 (1996).
- [13] V. Barger and R.J.N. Phillips, Collider Physics, (Addison-Wesley, Red-
wood City, CA, 1987)
- [14] ATLAS Collaboration, Technical Proposal, CERN/LHCC/94-43.
- [15] F. Ariztizabal et al., Nucl. Instr. and Meth. A349, (1994), 384.
- [16] M. Bosman, VI Int.Conf. on Calorimetry in HEP - Frascati, June 8-14,
1996 Frascati Physics series Vol.VI,(pp.299-310)

Received by Publishing Department
on December 17, 1999.

**The Publishing Department
of the Joint Institute for Nuclear Research
offers you to acquire the following books:**

Index	Title
E1,2-97-79	Proceedings of the XIII International Seminar on High Energy Physics Problems. Relativistic Nuclear Physics and Quantum Chromodynamics. Dubna, 1996 (2 volumes, 364 p. and 370 p., in English)
D5,11-97-112	Proceedings of the 9th International Conference «Computational Modelling and Computing in Physics». Dubna, 1996 (378 p., in English)
E2-97-213	Proceedings of the V International Seminar on Interaction of Neutron with Nuclei «Neutron Spectroscopy, Nuclear Structure, Related Topics». Dubna, 1997 (446 p., in English)
E2,4-97-263	Proceedings of the Third International Conference «Renormalization Group'96». Dubna, 1996 (436 p., in English)
E10-97-272	Proceedings of the Data Acquisition Systems of Neutron Experimental Facilities (DANEF'97). Dubna, 1997 (325 p., in English)
D19-97-284	Proceedings of the International Symposium «Problems of Biochemistry. Radiation and Space Biology». Dubna, 1997 (2 volumes 284 p. and 405 p., in Russian and English)
E2-97-413	Proceedings of the VII Workshop on High Energy Spin Physics (SPIN'97). Dubna, 1997 (398 p., in English)
E15-98-57	Proceedings of the III International Workshop «Hyperfine Structure and Nuclear Moments of Exotic Nuclei by Laser Spectroscopy». Poznan, 1997 (200 p., in English)
D13-98-66	Proceedings of the XVII International Symposium on Nuclear Electronics. Varna, 1997 (242 p., in English and Russian)
E1,2-98-154	Proceedings of the XIII International Seminar on High Energy Physics Problems «Relativistic Nuclear Physics and Quantum Chromodynamics». Dubna, 1996 (2 volumes 300 p. and 282 p., in English)
E3,14-98-168	Proceedings of the German-Russian User Meeting «Condensed Matter Physics with Neutrons at IBR-2». Dubna, 1998 (126 p., in English)
E3-98-202	Proceedings of the VI International Seminar on Interaction of Neutrons with Nuclei Neutron Spectroscopy, Nuclear Structure, Related Topics. Dubna, 1998 (352 p., in English)
D1,2-98-215	Proceedings of the International Workshop «Relativistic Nuclear Physics: from MeV to TeV». Dubna, 1998 (384 p., in English and Russian)
E2-98-254	Proceedings of the International Workshop Hadronic Atoms and Positronium in the Standard Model. Dubna, 1998 (260 p., in English)

Index	Title
D9,11-98-273	Proceedings of the 4th International Workshop «Beam Dynamics and Optimization». Dubna, 1997 (162 p., in English and Russian)
E17,19-98-305	Proceedings of the International Workshop on Deuteration of Biological Molecules for Structural and Dynamic Studies. Applications to Neutron Scattering and NMR. Dubna, 1998 (100 p., in English)
E1,2-98-307	Proceedings of the International School–Seminar «Actual Problems of Particle Physics». Gomel, 1997, Belarus (2 volumes 304 p. and 220 p., in English)
E2-98-372	Proceedings of the III International Workshop «Classical and Quantum Integrable Systems». Erevan, 1998 (200 p., in English)
E9-99-26	Proceedings of the International Conference on High Energy Accelerators. Dubna, 1998 (432 p., in English)
E2-99-35	Proceedings of the XI International Conference on Problems of Quantum Field Theory. Dubna, 1998 (508 p., in English)
E5-99-38	Proceedings of the International Workshop «Self-Similar Systems». Dubna, 1998 (404 p., in English)
E9-99-92	Proceedings of the XVII International Conference on High Energy Accelerators HEACC'98. Dubna, 1998 (435 p., in English)

Please apply to the Publishing Department of the Joint Institute for Nuclear Research for extra information. Our address is:

Publishing Department
Joint Institute for Nuclear Research
Dubna, Moscow Region
141980 Russia
E-mail: publish@pds.jinr.dubna.su.

Усубов З.У., Абдинов О.Б.

E1-99-332

Поиск структуры кварков на LHC.

Анализ угловых распределений адронных струй

Рассмотрены эффекты в угловых распределениях адронных струй, по которым может быть обнаружена структура кварков на LHC. Изучено влияние функции распределения кварков, нелинейности отклика и энергетического разрешения калориметра на эффекты, связанные со структурой кварков. Приведены оценки значений константы связи кварков, которые могут быть достигнуты в эксперименте при разных светимостях LHC.

Работа выполнена в Лаборатории ядерных проблем ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 1999

Usubov Z.U., Abdinov O.B.

E1-99-332

A Search for Quark Compositeness at the LHC.

Dijet Angular Distributions

This paper considers the effects of dijet angular distributions coming from a model of quark compositeness. The influence of the parton distribution function, calorimeter non-linearity and energy resolution is investigated. The data sensitivity to the quark compositeness scale for low and high LHC luminosity is studied.

The investigation has been performed at the Laboratory of Nuclear Problems, JINR.

Preprint of the Joint Institute for Nuclear Research. Dubna, 1999