United Nations Educational Scientific and Cultural Organization
and
International Atomic Energy Agency
THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**PROJECTIVE NORMALITY OF THE WONDERFUL COMPACTIFICATION
OF SEMISIMPLE ADJOINT GROUPS**

S. Senthamarai Kannan¹
Chennai Mathematical Institute, 92, GN Chetty Road, Chennai-17, India
and
The Abdus Salam International Centre for Theoretical Physics, Trieste, Italy.

Abstract

We prove that if G is a semisimple adjoint group over an algebraically closed field of arbitrary characteristic, X is the wonderful compactification of G and $\widehat{G}$ is a simply connected covering of G , then for any ample line bundle L over X , the cone over X given by L is normal.

MIRAMARE – TRIESTE

October 2000

¹E-mail: kannan@ictp.trieste.it; kannan@smi.ernet.in

1 Introduction

In [4], what are known as “wonderful compactifications” of symmetric varieties were constructed and studied by C. De Concini and C. Procesi. More precisely, if G is a semisimple adjoint group over the field of complex numbers, H is the subgroup of all fixed points of an involution σ of G that is induced by an involution $\hat{\sigma}$ of the simply connected covering $\hat{G}$ of G , then, they have constructed a complete embedding $\overline{G/H}$ of the homogeneous space G/H , with boundary being a union of normal crossing divisors. In particular, one gets such a compactification $\overline{G}$ for the group G (G being considered as $(G \times G)/\Delta(G)$). In [10], E. Strickland has proved that the wonderful compactifications for the group exists over algebraically closed fields of positive characteristics. In [5], C. De Concini and T.A. Springer have proved that these compactifications for arbitrary symmetric space G/H exists when the base field is of characteristic $p \neq 2$. In [6], G. Faltings raised the question: For what $\hat{G}$ -linearised line bundles L , the cone over $\overline{G/H}$ given by L is normal?

The aim of this paper is to provide an affirmative answer to his question for the case of the wonderful compactification of a semisimple adjoint group G over an algebraically closed field of arbitrary characteristic. To be more precise, we prove that if G is a semisimple adjoint group over an algebraically closed field of arbitrary characteristic, $\hat{G}$ is a simply connected covering of G , then for any $\hat{G} \times \hat{G}$ -linearised very ample line bundle L over the wonderful compactification $\overline{G}$, the cone over $\overline{G}$ given by L is normal. In section 3, we prove this result when the base field is the field of complex numbers. For a more precise statement, see Theorem 3.4. In section 4, we prove the result for algebraically closed fields of positive characteristic by using the properties of good filtrations.

2 Notations and basic Theorems

Throughout sections 2 and 3, we fix the following notations: Let G be a semisimple adjoint group over the field of complex numbers. Let $\pi : \hat{G} \longrightarrow G$ be a simply connected covering of G . Let $k[\hat{G}]$ denote the co-ordinate ring of $\hat{G}$. Let T be a maximal torus of G and B be a Borel subgroup of G containing T . Let B^- be the opposite Borel subgroup of G determined by T and B . Let $\hat{T}$ denote the pull back of T in $\hat{G}$. Let $\hat{B}$ (resp. $\hat{B}^-$) denote the pull back of B (resp. B^-) in $\hat{G}$. Let W denote the Weyl group of G with respect to T .

Let Φ denote the set of roots with respect to T as above and $\Phi^+ \subset \Phi$ denote the set of positive roots with respect to B as above.

Let $\Delta \subset \Phi^+$ denote the set of simple roots. We label the elements of Δ by :

$$\Delta := \{\alpha_1, \alpha_2, \dots, \alpha_l\}, \quad l = \text{rank}(G).$$

Let χ denote the weight lattice of G with respect to T . Let $E := \chi \otimes \mathbb{R}$ denote the $\mathbb{R}$ span of χ . Let $(\cdot, \cdot)$ denote the positive definite bilinear form on E induced by the Killing form of the Lie algebra of G and define $\langle \alpha, \beta \rangle = \frac{2(\alpha, \beta)}{(\beta, \beta)}$. Now, consider the fundamental weights. Since, they form a dual basis of the simple coroots, we label them:

$$\{\varpi_1, \dots, \varpi_l\},$$

where ϖ_i is dual to α_i . Let w_0 denote the longest element of W with respect to Δ .

In this paragraph, we recall some basic properties of the wonderful compactification of the symmetric spaces. Let G be a semisimple adjoint group, let σ be an involution of G that is induced by an involution $\hat{\sigma}$ of $\hat{G}$. Let H denote the subgroup of all fixed points of σ in G . Let H' denote the fixed points of $\hat{\sigma}$ in $\hat{G}$. Then, there exist a σ -stable maximal torus T of G and a Borel subgroup B of G containing T with the following property :

1. The dimension of $\{t \in T : \sigma(t) = t^{-1}\}$ is maximal.
2. For any positive root α (with respect to B), if $\sigma(\alpha)$ is a positive root, then $\sigma(\alpha) = \alpha$.

Let $\{\gamma_1, \gamma_2, \dots, \gamma_j\}$ be the set of all simple roots that are not fixed by σ . Then, σ induces a permutation $\tilde{\sigma}$ on the set $\{1, 2, \dots, j\}$.

Also, there is only one closed G -orbit in the wonderful compactification $\overline{G/H}$, say G/P and the restriction map $\text{Pic}(\overline{G/H}) \rightarrow \text{Pic}(G/P)$ is injective. For a dominant weight λ , let V_λ denote the irreducible representation of $\hat{G}$ with highest weight λ . Let λ be a dominant weight such that $V_\lambda^{H'}$ is non zero. Then, [by Theorem 8.3, pp 30-31 [4]], the global sections of the line bundle L_λ on $\overline{G/H}$ is given by:

$$2.1 \quad H^0(\overline{G/H}, L_\lambda) = \bigoplus_{\nu \in \Gamma_\lambda} V_\nu^*, \quad \text{where } \Gamma_\lambda \text{ denote the set of all dominant weights } \nu \text{ that are of the form } \nu = \lambda - \left(\sum_{i \leq \tilde{\sigma}(j)} m_i (\gamma_i - \sigma(\gamma_i))\right), \text{ with } m_i \in \mathbb{Z}_{\geq 0}.$$

Now, we apply these to our special involution $\sigma : G \times G \rightarrow G \times G$ sending $(x, y) \mapsto (y, x)$.

Here, the symmetric space is the group $G = (G \times G)/\Delta(G)$. Let X denote the wonderful compactification of G .

Here, we should take $T \times T$ as a σ -stable maximal torus with the above maximal property. We should take $B \times B^-$ ($T \subset B$) as a Borel subgroup of $G \times G$ containing $T \times T$ with the above property. Here, the unique closed $G \times G$ -orbit is $G/B \times G/B^-$, and the image of $\text{Pic}(X) \rightarrow \text{Pic}(G/B \times G/B^-)$ consists of line bundles associated to weights of the form $(\lambda, -\lambda)$. For any dominant weight λ , let L_λ denote the line bundle over X associated to λ and let

$H^0(X, L_\lambda)$ denote the global sections of L_λ over X . Let $\leq$ denote the dominant ordering on χ . That is $\nu \leq \lambda$ if and only if $\lambda - \nu = \sum_{i=1} m_i \alpha_i$ for some $m_i \in \mathbb{Z}_{\geq 0}$.

Now, applying the above statement 2.1 to our special involution of $G \times G$, we have

2.2 $H^0(X, L_\lambda) = \bigoplus_{\nu \leq \lambda} \text{End}(V_\nu^*)$, where the sum is taken over all dominant weights ν such that $\nu \leq \lambda$.

3 Projective normality

We now prove that for any line bundle L_λ on X , there is a unique (up to multiplication by a constant) injective homomorphism

$$\phi_\lambda : H^0(X, L_\lambda) \longrightarrow k[\widehat{G}] \quad (3.0.1)$$

of $\widehat{G} \times \widehat{G}$ modules.

For a proof of this fact: Since $G = G \times G / \Delta(G)$ is a $G \times G$ stable open subset of X , and the natural map $\pi : \widehat{G} = \widehat{G} \times \widehat{G} / \Delta(\widehat{G}) \longrightarrow G = G \times G / \Delta(G)$ is a surjection, the natural map $\pi : \widehat{G} = \widehat{G} \times \widehat{G} / \Delta(\widehat{G}) \longrightarrow X$ is a dominant map and hence for any line bundle L over X , $H^0(X, L)$ is a $\widehat{G} \times \widehat{G}$ -sub module of the $\widehat{G} \times \widehat{G}$ module $H^0(\widehat{G}, L)$. Since $k[\widehat{G}]$ is a unique factorisation domain, the pull back $\pi^*(L)$ on $\widehat{G}$ is trivial and hence we have

$$H^0(\widehat{G}, L) = k[\widehat{G}].$$

Moreover, this identification is unique up to constants since the units of the k - algebra $k[\widehat{G}]$ are only nonzero constants. Thus, (3.0.1) holds.

We now prove that for any two line bundles L_λ and L_μ , the following diagram is commutative:

$$\begin{array}{ccc} H^0(X, L_\lambda) \otimes H^0(X, L_\mu) & \longrightarrow & H^0(X, L_\lambda \otimes L_\mu) \\ \downarrow & & \downarrow \\ k[\widehat{G}] \otimes k[\widehat{G}] & \longrightarrow & k[\widehat{G}] \end{array} \quad (3.0.2)$$

Here the horizontal arrows are the natural maps, the vertical map on the left is $\phi_\lambda \otimes \phi_\mu$ and the vertical map on the right is $\phi_{\lambda+\mu}$.

For a proof of this observation:

By the Peter-Weyl Theorem, the $\widehat{G} \times \widehat{G}$ - module $k[\widehat{G}] = \bigoplus_\nu \text{End}(V_\nu^*)$, where the sum is taken over all dominant weights ν . Using this and the fact that $H^0(X, L_\lambda) = \bigoplus_{\nu \leq \lambda} \text{End}(V_\nu^*)$, it is easy to see that (3.0.2) holds.

Now, we prove the following Lemma.

Lemma 3.1. *Let λ and μ be two dominant weights. Let λ_1 be a dominant weight such that $\lambda_1 \leq \lambda$ and μ_1 be a dominant weight such that $\mu_1 \leq \mu$. Let ν be a dominant weight such that V_ν is a $\widehat{G}$ -submodule of the tensor product $V_{\lambda_1} \otimes V_{\mu_1}$. Then, the image of $\text{End}(V_{\lambda_1}^*) \otimes \text{End}(V_{\mu_1}^*)$ under the multiplication map $H^0(X, L_\lambda) \otimes H^0(X, L_\mu) \longrightarrow H^0(X, L_{\lambda+\mu})$ contains $\text{End}(V_\nu^*)$.*

Proof: Since $\widehat{G}$ is linearly reductive and V_ν^* is a quotient of $V_{\lambda_1}^* \otimes V_{\mu_1}^*$, we can assume that V_ν^* is a submodule of $V_{\lambda_1}^* \otimes V_{\mu_1}^*$.

Now, consider the $\widehat{B}$ action on $\text{End}(V_{\lambda_1}^*)$ on the right and consider the $\widehat{B}^-$ action on $\text{End}(V_{\mu_1}^*)$ on the right. By the observation 2.2 and the hypothesis $\lambda_1 \leq \lambda$ and $\mu_1 \leq \mu$, we have $\text{End}(V_{\lambda_1}^*) \subset H^0(X, L_\lambda)$ and $\text{End}(V_{\mu_1}^*) \subset H^0(X, L_\mu)$. So, by our identification (3.0.1), we have $\text{End}(V_{\lambda_1}^*) \subset H^0(X, L_\lambda) \subset k[\widehat{G}]$ and $\text{End}(V_{\mu_1}^*) \subset H^0(X, L_\mu) \subset k[\widehat{G}]$. Therefore, by the Borel-Weil Theorem [cf [1]], we have $V_{\lambda_1}^* = \{f \in \text{End}(V_{\lambda_1}^*) : f(xb) = \lambda_1(b).f(x) \text{ for } x \in \widehat{G}, b \in \widehat{B}\} = \{f \in k[\widehat{G}] : f(xb) = \lambda_1(b).f(x) \text{ for } x \in \widehat{G}, b \in \widehat{B}\}$ and $V_{\mu_1}^* = \{f \in \text{End}(V_{\mu_1}^*) : f(xb^-) = w_0(\mu_1)(b^-).f(x) \text{ for } x \in \widehat{G}, b^- \in \widehat{B}^-\} = \{f \in k[\widehat{G}] : f(xb^-) = w_0(\mu_1)(b^-).f(x) \text{ for } x \in \widehat{G}, b^- \in \widehat{B}^-\}$.

Fix our choice of $V_{\lambda_1}^*$ and $V_{\mu_1}^*$ as above and let $f = \sum_i f_i \otimes g_i \in V_{\lambda_1}^* \otimes V_{\mu_1}^*$, $f_i \in V_{\lambda_1}^*$, $g_i \in V_{\mu_1}^*$ be such that $(\sum_i f_i \otimes g_i)(x, x)$ is zero, for every $x \in \widehat{G}$. Then, we have $f((xb, xb^-)) = (\lambda_1(b))(w_0(\mu_1)(b^-))f(x, x) = 0$ for every $x \in \widehat{G}$, $b \in B$ and $b^- \in B^-$. Since the elements (xb, xb^-) form a dense open set in $\widehat{G} \times \widehat{G}$, f must vanish identically on $\widehat{G} \times \widehat{G}$. Thus, $V_{\lambda_1}^* \otimes V_{\mu_1}^*$ is mapped injectively via the multiplication map $\text{End}(V_{\lambda_1}^*) \otimes \text{End}(V_{\mu_1}^*) \subset k[\widehat{G}] \otimes k[\widehat{G}] \longrightarrow k[\widehat{G}]$. By the hypothesis $\lambda_1 \leq \lambda$, $\mu_1 \leq \mu$ and by Theorem(2.1), we have $\text{End}(V_{\lambda_1}^*) \subset H^0(X, L_\lambda)$ and $\text{End}(V_{\mu_1}^*) \subset H^0(X, L_\mu)$. Therefore, by the observation (3.0.2), the $\widehat{G}$ module $V_{\lambda_1}^* \otimes V_{\mu_1}^*$ is mapped injectively into $H^0(X, L_{\lambda+\mu})$ via the multiplication map $H^0(X, L_\lambda) \otimes H^0(X, L_\mu) \longrightarrow H^0(X, L_{\lambda+\mu})$. Since V_ν^* is a $\widehat{G}$ -submodule of $V_{\lambda_1}^* \otimes V_{\mu_1}^*$, V_ν^* must be isomorphic to a $\widehat{G}$ submodule of the image of $\text{End}(V_{\lambda_1}^*) \otimes \text{End}(V_{\mu_1}^*)$. Since the image of $\text{End}(V_{\lambda_1}^*) \otimes \text{End}(V_{\mu_1}^*)$ is a $\widehat{G} \times \widehat{G}$ submodule of $k[\widehat{G}]$, it must contain $\text{End}(V_\nu^*)$.

Therefore, using the diagram (3.0.2), it is easy to see that $\text{End}(V_\nu^*)$ is a $\widehat{G} \times \widehat{G}$ -submodule of the image of $H^0(X, L_\lambda) \otimes H^0(X, L_\mu) \longrightarrow H^0(X, L_{\lambda+\mu})$ under the multiplication map.

This completes the proof of the Lemma.

We now prove the following Lemma.

Lemma 3.2. *Let λ and μ be two dominant weights. Let ν be a dominant weight satisfying $\nu \leq \lambda + \mu$. Then, there is a dominant weight $\lambda_1 \leq \lambda$ and there is a dominant weight $\mu_1 \leq \mu$ such that V_ν is $\widehat{G}$ -submodule of the tensor product $V_{\lambda_1} \otimes V_{\mu_1}$.*

Proof: Let λ and μ be two dominant weights. Let ν be a dominant weight such that $\nu \leq \lambda + \mu$. We wish to prove that there are dominant weights λ_1, μ_1 such that $\lambda_1 \leq \lambda$, $\mu_1 \leq \mu$

and an element w of the Weyl group W such that $\nu = \lambda_1 + w(\mu_1)$, and then using PRV conjecture [cf [9]], we prove that V_ν is a $\widehat{G}$ -submodule of the tensor product $V_{\lambda_1} \otimes V_{\mu_1}$.

By [sections (13.4) and (21.3)] of [8], ν is a weight of $V_{\lambda+\mu}$. Therefore, we have $\nu = \nu_1 + \nu_2$, where ν_1 is a weight of V_λ and ν_2 is a weight of V_μ . We have $\lambda - \nu_1 = (\sum_i^l m_i \alpha_i)$ for some $m_i \in \mathbb{Z}_{\geq 0}$.

We first prove that there is a dominant weight λ_1 such that $\lambda_1 \leq \lambda$ and there is a weight ν_2' of V_μ such that $\nu = \lambda_1 + \nu_2'$ by induction on the sum $\sum_i^l m_i$.

If $\sum_i^l m_i = 0$, there is nothing to prove.

If $\sum_i^l m_i = 1$, then, we have $\nu_1 = \lambda - \alpha_{i_0}$ for some $i_0 \in \{1, 2, \dots, l\}$. Now, if ν_1 is dominant, then there is nothing to prove. Otherwise, $\langle \nu_1, \alpha_{i_0} \rangle$ is negative. Since $\nu = \nu_1 + \nu_2$ is dominant, $\langle \nu_2, \alpha_{i_0} \rangle$ is positive. But, we know that $\nu_1 + k\alpha_{i_0}$ is a weight of V_λ for any $k \in \{0, 1, \dots, -\langle \nu_1, \alpha_{i_0} \rangle\}$ and $\nu_2 - k\alpha_{i_0}$ is a weight of V_μ for any $k \in \{0, 1, \dots, \langle \nu_2, \alpha_{i_0} \rangle\}$ [cf [sections (13.4) and (21.3) of [8]]]. In particular, $\nu_1 + \alpha_{i_0} = \lambda$ is a weight of V_λ (which is ofcourse trivial) and $\nu_2 - \alpha_{i_0}$ is a weight of V_μ and we also have $\nu = \lambda + \nu_2 - \alpha_{i_0}$. Therefore, in this case, we take $\lambda_1 = \lambda$ and $\nu_2' = \nu_2 - \alpha_{i_0}$.

Now, let N be a positive integer strictly bigger than 1. Let us assume that the statement is true for all ν such that $\nu = \nu_1 + \nu_2$, where $\nu_1 = \lambda - \sum_{i=1}^l m_i \alpha_i$ being a weight of V_λ with the sum $\sum_{i=1}^l m_i \leq N - 1$ and ν_2 being a weight of V_μ .

Let ν be a dominant weight such that $\nu \leq \lambda$ and $\nu = \nu_1 + \nu_2$ where ν_1 is a weight of V_λ , ν_2 is a weight of V_μ with $\nu_1 = \lambda - \sum_i m_i \alpha_i$ and the sum $\sum_{i=1}^l m_i = N$. If ν_1 is dominant, then, there is nothing to prove. Otherwise, there is a simple root α_{i_0} such that $\langle \nu_1, \alpha_{i_0} \rangle < 0$.

By a proof similar to the case when $\sum_i^l m_i = 1$, it is easy to see that $\nu_1 + \alpha_{i_0}$ is a weight of V_λ and $\nu_2 - \alpha_{i_0}$ is a weight of V_μ . Also, we have $\nu = (\nu_1 + \alpha_{i_0}) + (\nu_2 - \alpha_{i_0})$. Therefore, by induction, there exist a dominant weight $\lambda_1 \leq \lambda$ and a weight ν_2' of V_μ such that

$$\nu = \lambda_1 + \nu_2'.$$

We fix our choice of λ_1 and ν_2' as above. Since ν_2' is a weight of V_μ , $w(\nu_2')$ is also a weight of V_μ for every $w \in W$. Thus, the unique dominant weight (say $\mu_1 = w^{-1}(\nu_2')$) in the W orbit of ν_2' must also be a weight of V_μ .

Hence, we have $\nu = \lambda_1 + w(\mu_1)$ where λ_1 is a dominant weight satisfying $\lambda_1 \leq \lambda$, μ_1 is a dominant weight satisfying $\mu_1 \leq \mu$ and w is an element of the Weyl group W .

Now, the fact that V_ν is a $\widehat{G}$ -submodule of the tensor product $V_{\lambda_1} \otimes V_{\mu_1}$ follows from PRV Conjecture [cf [9]].

This completes the proof of this Lemma.

We have

Corollary 3.3. *Let λ and μ be two dominant weights. Then, the multiplication map*

$$H^0(X, L_\lambda) \otimes H^0(X, L_\mu) \longrightarrow H^0(X, L_{\lambda+\mu}) \text{ is onto.}$$

Proof: Proof of this is an immediate consequence of Lemmas 3.1 and 3.2.

Theorem 3.4. *Let G be a semisimple adjoint group over the field of complex numbers. Let $\pi : \widehat{G} \longrightarrow G$ be a simply connected covering of G . Let X denote the wonderful compactification of G . Let L be an ample line bundle over X . Consider the embedding of X in $\mathbb{P}(H^0(X, L)^*)$. Let $\widehat{X}$ denote the cone over X with respect to this embedding of X . Then, $\widehat{X}$ is normal.*

Proof: By [4], we have $L = L_\lambda$ for some regular dominant weight λ . By [II (5.14)(d)] of [7], it suffices to prove that the multiplication map

$$H^0(X, L_\lambda)^{\otimes N} \longrightarrow H^0(X, L_{N\lambda}) \tag{3.4.1}$$

is surjective for every positive integer N . The proof of this is an immediate consequence of Corollary 3.3.

We have

Corollary 3.5. *Let w be an element of the Weyl group W . Let X_w denote the scheme theoretic closure of the cell BwB in X . Let L be a $\widehat{G} \times \widehat{G}$ linearised very ample line bundle over X and let $\widehat{X}_w$ denote the cone over the embedding given by L . Then, $\widehat{X}_w$ is normal.*

Proof: By [Corollary 3, [3]], the restriction map $H^0(X, L^{\otimes n}) \longrightarrow H^0(X_w, L^{\otimes n})$ is surjective for every positive integer $n \cdots$ (3.5.1).

Now, consider the following commutative diagram of maps:

$$\begin{array}{ccc} H^0(X, L)^{\otimes n} & \longrightarrow & H^0(X, L^{\otimes n}) \\ \downarrow & & \downarrow \\ H^0(X_w, L)^{\otimes n} & \longrightarrow & H^0(X_w, L^{\otimes n}) \end{array} \tag{3.5.2}$$

Here the horizontal arrows are the natural multiplication maps and the vertical arrows are restriction maps.

The assertion of the corollary follows from the observations 3.5.1, 3.5.2 and Corollary 3.3.

4 Appendix:

In this section, we prove the analogue of Theorem 3.4 when the base field is of positive characteristic.

Let k be an algebraically closed field of positive characteristic. Let G be a semisimple adjoint group over k . Let $\widehat{G}$ be a simply connected covering of G . Let X denote the wonderful compactification of G . Let $T, B, B^-, \widehat{T}, \widehat{B}, \widehat{B}^-$ be as in section 2.

With the notations as in section 2, we introduce some more notations as follows: Let χ denote the weight lattice of G with respect to T . So, $\chi \times \chi$ is the weight lattice of $G \times G$ with respect to $T \times T$.

For any weight $\lambda \in \chi$, set $V_\lambda := H^0(G/B, L_{w_0(-\lambda)})$.

Definition 4.1. Let $\widehat{G}$ be as above. Let V be a $\widehat{G}$ module. A filtration of $\widehat{G}$ -submodules $(0) = V_0 \subset V_1 \subset V_2 \subset \cdots \subset V_n = V$ is a good filtration if each V_{i+1}/V_i is a direct sum of $V(\lambda_{i(j)})$'s for some dominant weights $\lambda_{i(j)}$'s.

As in section 2, let $\Delta = \{\alpha_1, \dots, \alpha_l\}$ denote the set of simple roots with respect to T and B .

Let λ and μ be two dominant weights in χ .

Let L_λ (resp. L_μ) denote the line bundle over X associated to $(\lambda, -\lambda)$ (resp. to $(\mu, -\mu)$). Let ν be a dominant weight such that $\nu \leq \lambda + \mu$. Let $\phi : H^0(X, L_\lambda) \otimes H^0(X, L_\mu) \longrightarrow H^0(X, L_{\lambda+\mu})$ denote the multiplication map.

Then, we have

Lemma 4.2. *The $\widehat{G} \times \widehat{G}$ -module $V_{w_0(-\nu)} \otimes V_\nu$ is a $\widehat{G} \times \widehat{G}$ -subquotient of the image $\phi(H^0(X, L_\lambda) \otimes H^0(X, L_\mu))$.*

Proof: By Donkin's conjecture [cf Theorem 4.4.3, [11]], a tensor product of two modules with good filtrations has a good filtration. By Theorem 5.10 of [5], $H^0(X, L_{\lambda_1})$ (resp. $H^0(X, L_{\mu_1})$) has a good filtration of $\widehat{G} \times \widehat{G}$ -submodules such that the successive quotients look like $V_{w_0(-\nu)} \otimes V_\nu$, $\nu \leq \lambda_1$ (resp. $\nu \leq \mu_1$). Therefore, the tensor product $H^0(X, L_{\lambda_1}) \otimes H^0(X, L_{\mu_1})$ has a good filtration of $\widehat{G} \times \widehat{G}$ -submodules.

Now, let H be a semisimple Chevalley algebraic group over the field of complex numbers. Let H_p be the corresponding algebraic group in characteristic p . Now, if V_p is a H_p -module having a good filtration in characteristic p such that it comes from a H -module V modulo p ,

then the character of V is same as the character of V_p and so the H_p - subquotients of V_p are precisely the H - direct summands of V .

Now, let M_ν (resp. M'_ν) be the largest $\widehat{G} \times \widehat{G}$ submodule of the tensor product $H^0(X, L_{\lambda_1}) \otimes H^0(X, L_{\mu_1})$ all of whose $B \times B$ - weights are $\leq (-w_0(\nu), \nu)$ (resp. $< (-w_0(\nu), \nu)$). Also, let N_ν and N'_ν be the corresponding submodules of $H^0(X, L_{\lambda_1 + \mu_1})$. Therefore, from Lemmas 3.1 and 3.2, and by the observation above, ϕ induces a non zero $\widehat{G} \times \widehat{G}$ - module homomorphism from the quotient $(V_{-w_0(\nu)} \otimes V_\nu)^m = M_\nu/M'_\nu \rightarrow N_\nu/N'_\nu = V_{-w_0(\nu)} \otimes V_\nu$ for some positive integer m . Therefore, by the Frobenius reciprocity [cf Proposition 2.1.6 [11]], the assertion follows.

The Theorem (3.4) for positive characteristic follows from Lemma(4.2).

Acknowledgements: We thank Professor C.De Concini for suggesting the problem. We thank Professors M.S.Narasimhan and C.Procesi for their encouragement and useful discussions. We thank ICTP for the hospitality during the stay. We thank the referee for the useful comments and for suggesting the problem for positive characteristics.

References

- [1] A.Borel, Linear representations of semi-simple algebraic groups, AMS, Proceedings of Symposia in Pure Mathematics, volume 29, 1974.
- [2] N.Bourbaki, Groupes et Algebres de Lie, Ch 4, 5 et 6, Paris: Hermann, 1968.
- [3] M.Brion and P.Polo, Large Schubert Varieties, Representation Theory 4 (2000), 97-126.
- [4] C.De Concini and C.Procesi, Complete symmetric varieties, pp 1-43, Invariant Theory, 996, Lecture Notes in Mathematics.
- [5] C.De Concini and T.A.Springer, Compactification of symmetric varieties, Transformation Groups, Vol.4, No.2-3, 1999, pp. 273-300.
- [6] G.Faltings, Explicit resolution of local singularities of moduli spaces, J. reine angew. Math. 483 (1997) pp. 183-196.
- [7] R.Hartshorne, Algebraic Geometry, Springer-Verlag, 1977.
- [8] J.E.Humphreys, Introduction to Lie algebras and representation theory, Springer - Verlag, 1972.

- [9] S.Kumar, Proof of the Parthasarathy-Ranga Rao-Varadarajan Conjecture, Invent. Math. 93, 117-130 (1988).
- [10] E. Strickland, A Vanishing result for group compactifications, Math. Ann. 277 (1987), 165-171.
- [11] Wilberd van der Kallen, On Frobenius Splittings and B-Modules, TIFR Lecture notes in Maths, Springer-Verlag, 1993.