Tau Physics at LEP

Randall Sobie

Institute for Particle Physics and University of Victoria

Introduction

Topological branching ratios

Leptonic decay modes

Tau polarization

Summary

1974

Discovery of the tau

1980's

First measurements of tau properties

1990's

Precision measurements at LEP and CLEO

2000's

High statistics samples at the B-factories

LEP was a unique environment for studying the tau

$$e^+e^- \rightarrow Z^0 \rightarrow \tau^+\tau^-$$

Two highly collimated tau jets

200K tau pairs per LEP experiment

Little background ($\sim 1\%$)

- ALEPH: tau polarization (2001)
- ALEPH: Michel parameters (2001)
- L3: tau leptonic branching ratios (2001)
- L3: topological branching ratios (2001)
- DELPHI: topological branching ratios (2001)
- OPAL: tau polarization (2001)

Topological branching ratios

The tau “1-prong” problem of the 1980’s.

Is the tau a heavy lepton?

Was it decaying into a new particle?

$$B_{\text{inclusive}} = \sum B_{\text{exclusive}}$$

$$85.94 \pm 0.23\% \quad 82.0 \pm 1.2\%$$

The source of the 1-prong problem was found to be due to tau decays with neutral pions.

2001 PDG B(excl):

$$\sum B_{\text{excl}} = 85.32 \pm 0.13$$

Pizero reconstruction

New topological BR results

First new measurements in a decade

2001 PDG $B(\text{excl})$:

$$\sum B_{\text{excl}} = 85.32 \pm 0.13$$

DELPHI

$$B(1p) = (85.316 \pm 0.093 \pm 0.019)\%$$

$$B(3p) = (14.569 \pm 0.093 \pm 0.018)\%$$

$$B(5p) = (0.115 \pm 0.013 \pm 0.006)\%$$

L3

$$B(1p) = (85.274 \pm 0.105 \pm 0.073)\%$$

$$B(3p) = (14.556 \pm 0.105 \pm 0.076)\%$$

$$B(5p) = (0.170 \pm 0.022 \pm 0.026)\%$$

Challenging measurement to do accurately

photon conversion
hadronic interactions
neutral K decays

Reconstructed
nuclear
interactions

Lorentz structure in tau decays

When the final state polarization is integrated out, the lepton momentum spectrum can be described by the 4 Michel parameters

with SM values of

$$\frac{1}{\Gamma} \frac{d\Gamma}{dx} = f_0(x) + \rho f_1(x) + \eta \frac{m_l}{m_\tau} f_2(x) - P_\tau [\xi g_0(x) + \xi \delta g_2(x)]$$

$$(\eta, \rho, \xi, \xi \delta) = (1, 0.75, 1, 0.75)_{SM}$$

ALEPH (2001) lepton momentum spectra

Lorentz structure in tau decays (2)

In the case of tau hadronic decays, the momentum spectrum depends only on a single parameter ξ which is related to the neutrino helicity

$$\tau^- \rightarrow h^- \nu_\tau \quad h = \{\pi, \rho, a_1\}$$

$$\frac{1}{\Gamma} \frac{d\Gamma}{d\omega_h} = f_h(\omega_h) + \xi_h P_\tau g_h(\omega_h)$$

$$\text{eg. } \omega_\pi = \cos \theta \simeq 2x_\pi - 1$$

LEP experiments exploit the correlation between the two tau decays

2001 ALEPH: both leptonic and hadronic decay modes

OPAL, DELPHI, L3: tau leptonic modes

LEP results

PDG2001 Fit Values

$$\rho = 0.747 \pm 0.009$$

$$\eta = 0.011 \pm 0.031$$

$$\xi = 0.997 \pm 0.032$$

$$\delta\xi = 0.746 \pm 0.023$$

Does not include
the new ALEPH
result

Tau neutrino helicity

$$\xi_h = 1.000 \pm 0.008 \text{ (PDG2001)}$$

$$\xi_h = 0.992 \pm 0.011 \text{ (ALEPH)}$$

Consistent with the SM.
Consistent with lepton universality.

Leptonic decay width

The integrated leptonic decay width is given by

$$\Gamma = \frac{G^2 m_\tau^5}{192\pi^3} \left(1 + 4\eta \frac{m_l}{m_\tau} \right)$$

$$\Gamma(\tau \rightarrow l\nu\nu) = \left(\frac{g_\tau g_l}{8m_w^2} \right)^2 \frac{m_\tau^5}{96\pi^3} f\left(\frac{m_l^2}{m_t^2}\right) (1 + \delta_{RC}^\tau)$$

mass correction

photon radiation and W
propagator correction

The “low energy” parameter.
Can be used to put limits on
charged Higgs mass

Measurement of the strong
coupling constant

Comparison of the tau and muon
leptonic BR give precise tests of
lepton universality

Limit on the mass of the
tau neutrino

Limit on the mass of
excited leptons

Tau leptonic branching ratios

2001: L3 electron and muon BR

World Averages:

$$B_e = 17.83 \pm 0.06$$

$$B_\mu = 17.37 \pm 0.07$$

Lepton Universality

Are the couplings identical for all leptons?

A comparison of tau and muon electronic decays yields

$$\frac{g_\tau^2}{g_e^2} = 0.9996 \ B(\tau \rightarrow \mu \nu \nu) \frac{m_\mu^5}{m_\tau^5} \frac{\tau_\mu}{\tau_\tau}$$

$$\frac{g_\tau^2}{g_\mu^2} = 0.9996 \ B(\tau \rightarrow e \nu \nu) \frac{m_\mu^5}{m_\tau^5} \frac{\tau_\mu}{\tau_\tau}$$

Tau hadronic decays give $g(\text{tau})/g(\text{mu})$ at the 0.5% level

Assumes $g(e)=g(\mu)$

Lepton Universality

A comparison of tau leptonic decay modes gives $\frac{g_\mu^2}{g_e^2} = 1.0028 \frac{B(\tau \rightarrow \mu \nu \nu)}{B(\tau \rightarrow e \nu \nu)} = 1.0010 \pm 0.0020$

W decay modes give 1-2% precision.

Pion decay modes give 1.6% precision.

However, the pion decays are sensitive to a longitudinal W whereas tau decays are sensitive to a transverse W.

Tau decays can put limits on the mass of the charged Higgs

A comparison of tau electron and muon decays

$$\Gamma(\eta) = \Gamma^{SM} \left(1 + 4\eta \frac{m_l}{m_\tau} + \dots \right)$$

$$\frac{\Gamma(\tau \rightarrow \mu \nu \nu)}{\Gamma(\tau \rightarrow e \nu \nu)} = \frac{\Gamma_{SM}(\tau \rightarrow \mu \nu \nu)}{\Gamma_{SM}(\tau \rightarrow e \nu \nu)} \left(1 + 4\eta \frac{m_\mu}{m_\tau} \right) \quad \eta = 0.007 \pm 0.021$$

Limit on the mass of the charged Higgs

The Michel parameter, eta, is expected to be zero in the Standard Model but it can be used to place limits on a scalar coupling (A.Stahl)

$$\eta = \frac{1}{2} \text{Re}(g_{LL}^V g_{RR}^{S*} + \dots)$$

For a scalar coupling

$$\eta = \frac{g_{RR}^S}{2}$$

$$g_{RR}^S = -m_l m_\tau \left(\frac{\tan \beta}{m_{H^-}} \right)^2$$

Swain et al

$$m_{H^-} > 1.9 \tan \beta$$

Measurement of the strong coupling constant

The leptonic branching ratios and tau lifetime have sensitivity to the total hadronic width

$$\delta_{QCD} \sim 20\%$$

$$R_\tau = \frac{\Gamma(\tau \rightarrow \nu_\tau + \text{hadrons})}{\Gamma(\tau \rightarrow e \nu_\tau \nu_e)} \simeq 3(1 + \delta_{QCD})$$

Using the Operator Product Expansion

$$R_\tau = 3 (|V_{UD}|^2 + |V_{US}|^2) S_{EW} (1 + \delta_{EW} + \delta^0 + \sum_{D=2,4,\dots} \delta^D)$$

electroweak corrections

Non-perturbative and quark-mass corrections

The perturbative expansion is known to

$$\delta^0 = \frac{\alpha_S}{\pi} + 5.2023 \left(\frac{\alpha_S}{\pi} \right)^2 + 26.366 \left(\frac{\alpha_S}{\pi} \right)^3 + 78 \left(\frac{\alpha_S}{\pi} \right)^4$$

Strong coupling constant

R_τ	can be extracted from the tau leptonic branching ratios and the tau lifetime	3.645 ± 0.020	tau lifetime
		3.629 ± 0.015	leptonic branching ratios

Including a 40% theoretical error on the third order term and all non-perturbative contributions gives

$$\alpha_s(M_\tau) = 0.35 \pm 0.03$$

Extrapolating to the Z-mass gives

$$\alpha_s(M_Z) = 0.121 \pm 0.03$$

PDG average

$$\alpha_s(M_Z) = 0.117 \pm 0.02$$

Strong coupling constant

Clear evidence of the running of the strong coupling constant

Tau width, upsilon decays, deep inelastic scattering, electron-positron event rate at 25 GeV, event shapes at TRISTAN, Z width, e+e- event shapes at Z-mass, 135 and 189 GeV

Neutral currents - tau polarization

The nature of the weak interaction results in a polarization of the outgoing fermions.

Define Lepton Asymmetry

$$A_l = \frac{2v_l a_l}{|v_l|^2 + |a_l|^2}$$

pol	$A_\tau^{pol} = -A_t$
FB	$A_l^{FB} = 0.75 A_e A_l$
FB-pol	$A_l^{FB-POL} = 0.75 A_e$
LR	$A_l^{LR} = A_e$
FB-LR	$A_l^{FB-LR} = 0.75 A_l$

$$\frac{v}{a} = 1 - 4 \sin^2 \theta_W^{eff}$$

Tau polarization

The tau polarization can be measured by studying the kinematics of the tau decay products.

momentum spectra $e \mu \pi$

optimal observables ρa_1

Utilizes the correlation between both tau decays in a single event

Tau polarization results – OPAL and ALEPH (2001)

$$P_\tau = \frac{A_\tau (1 + \cos^2 \theta) + 2A_e \cos \theta}{(1 + \cos^2 \theta) + 2A_e A_\tau \cos \theta}$$

ALEPH

OPAL

$$A_e = 0.1504 \pm 0.0068 = 0.1456 \pm 0.0095$$

$$A_\tau = 0.1451 \pm 0.0059 = 0.1454 \pm 0.0113$$

Electroweak mixing angle

LEP EWWG:

$$A_e = 0.1498 \pm 0.0049$$

$$A_\tau = 0.1439 \pm 0.0043$$

$$A_l = \frac{2v_l a_l}{|v_l|^2 + |a_l|^2}$$

$$\frac{v}{a} = 1 - 4 \sin^2 \theta_W^{\text{eff}}$$

Summary

LEP has made significant experiment measurements of tau properties as well as using the tau to test the Standard Model

There have been a number of precision measurements over the past few years.

Close to the end with only a few more results expected.

OPAL tau muonic branching ratio

OPAL strange spectral functions