CMS Conference Report

29 Apr 2002

Squarks and Gluino Searches with CMS at LHC

G. Segneri

Università degli Studi di Pisa and INFN

Abstract

In this paper the studies about squarks and gluino searches with the CMS detector are reviewed. Inclusive squarks and gluino searches are described and the 5 sigma discovery contours are shown. The scalar bottom quark (*sbottom* or $\tilde{b}$) and gluino ($\tilde{g}$) reconstructions using the process $\tilde{g} \rightarrow \tilde{b}b$ are discussed and the resolutions on sbottom and gluino masses are presented. The scalar top quark (*stop* or $\tilde{t}$) search is also described and it is shown how the stop signal can be separated from the $t\bar{t}$ background and the background due to the overall supersymmetric production.

Presented at 8th *Adriatic Meeting - Central European Symposia on Particle Physics in the New Millennium*,
Dubrovnik, Croatia, 4-14 September 2001

Squarks and Gluino Searches with CMS at LHC

Gabriele Segneri

Università degli studi di Pisa and INFN

Abstract. In this paper the studies about squarks and gluino searches with the CMS detector are reviewed. Inclusive squarks and gluino searches are described and the 5 sigma discovery contours are shown. The scalar bottom quark (*sbottom* or $\tilde{b}$) and gluino ($\tilde{g}$) reconstructions using the process $\tilde{g} \rightarrow \tilde{b}b$ are discussed and the resolutions on sbottom and gluino masses are presented. The scalar top quark (*stop* or $\tilde{t}$) search is also described and it is shown how the stop signal can be separated from the $t\bar{t}$ background and the background due to the overall supersymmetric production.

1 Introduction

One of the main goals of the experiments at the Large Hadron Collider (LHC) [1] will be to find out new physics beyond the Standard Model. The Supersymmetry (SUSY) [2] and in particular the Minimal Supersymmetric Standard Model (MSSM) are possible extensions of this theory. Thanks to the centre of mass energy of 14 TeV which will be available at the LHC, it will be possible to extend the searches of the particles predicted by SUSY up to masses of 2.5 TeV.

The paper is dedicated to the searches with the CMS detector [3] of the new strongly interacting SUSY particles, the squarks and the gluino. The first part of the paper provides some general remarks about the MSSM-mSUGRA model which is assumed in the analyses. The inclusive squarks and gluino searches are described and for each of them the reach of the CMS experiment is provided. The reconstruction of the decay $\tilde{g} \rightarrow \tilde{b}b$ and the stop search are also discussed and some preliminary results are shown.

2 General Remarks

These studies are performed in the framework of the Minimal Supergravity (mSUGRA) model, a constrained version of the MSSM model. In this model Supersymmetry breaking occurs through the mediation of the universal gravitational interaction. At the Grand Unification (GUT) scale ($m_{GUT} \sim 2 \cdot 10^{16}$ GeV) gauginos and scalars have common masses and couplings. Masses and couplings at the Electroweak scale are derived from those at GUT scale through Renormalisation Group Equations. The independent parameters of the model are: the common scalar mass m_0 , the common gaugino mass $m_{1/2}$, the common trilinear coupling among the scalars A_0 , the ratio between the vacuum expectation values

Fig. 1. Total mSUGRA cross-section contours as function of m_0 and $m_{1/2}$ for $A_0 = 0$, $\mu > 0$ and $\tan \beta = 2$ (left figure) and $\tan \beta = 35$ (right figure) [5]

of the two Higgs doublets $\tan \beta$ and the sign of the higgsino mixing parameter μ .

An additional requirement is the R-parity conservation: as a consequence of this assumption, the SUSY particles can only be produced in even numbers at accelerator experiments. Moreover, the lightest SUSY particle (LSP) should be stable and each SUSY particle would decay into Standard Model particles and one LSP. The LSP in this model is the lightest neutralino $\tilde{\chi}_1^0$.

Contours corresponding to constant cross sections obtained with ISAJET 7.32 [4] are shown in fig.1 for $A_0 = 0$, $\mu > 0$, different values of m_0 and $m_{1/2}$ and $\tan \beta = 2$ (left plot) and $\tan \beta = 35$ (right plot) [5]. The $\mu < 0$ case is very similar and it is omitted. The overall SUSY production cross section (continuous line) is compared with the total cross section for processes with at least one strongly interacting SUSY particle (dashed line). At low values of m_0 and $m_{1/2}$ the two lines are very close since squarks and gluino are relatively light and dominate the SUSY particles production. Contours separate at higher m_0 and $m_{1/2}$ as squarks and sleptons become more massive.

3 Inclusive Searches

The inclusive searches are based on peculiar features which are common to all the SUSY processes involving squarks and gluino. A very high tagging efficiency can be achieved since the identification of the specific processes is not required.

The most remarkable features of squarks and gluino decays are:

- a large number of leptons produced by $Z^0 \rightarrow \ell^+\ell^-$, $W^\pm \rightarrow \ell\nu$, $\tilde{\chi}_2^0 \rightarrow \ell\tilde{\ell}, \ell\tilde{\ell}^*$, $\tilde{\ell} \rightarrow \ell\tilde{\chi}_1^0$ decays
- high E_T hadron jets from squarks and gluino decays ($\tilde{q} \rightarrow q\tilde{\chi}_i^0$, $\tilde{q} \rightarrow q'\tilde{\chi}_i^\pm$, $\tilde{g} \rightarrow q\bar{q}$)
- large missing transverse momentum due to LSPs which do not interact with the detector
- a large number of b quarks (from decays of $\tilde{t}, \tilde{b}, h, t$) and τ leptons (specially at high $\tan\beta$).

Several search strategies based on some of these peculiarities are investigated [5] and for each of them the region of sensitivity of the CMS experiment is calculated.

Signal events are generated for several m_0 and $m_{1/2}$ values, $\tan\beta = 2$ and 35, $A_0 = 0$ and both signs of μ . The SUSY processes are simulated with ISAJET 7.32 and the backgrounds ($t\bar{t}$, Z^0 plus jets, $W^\pm$ plus jets, QCD) with PYTHIA 5.7[6]. The detector response is reproduced with a fast detector simulation package (CMSJET[7]). For each of these points the SUSY selection cuts are optimised in order to achieve the best sensitivity. Sensitivity is defined as $S/\sqrt{S+B}$ being S and B the expected SUSY signal and the Standard Model background.

Five analyses, called 0l, 1l, 2lOS, 2lSS, 3l are based on the requirements of no leptons, at least one lepton, two leptons with opposite charge, two leptons with the same charge and three lepton respectively and one analysis is based only on missing transverse momentum (E_T^{miss}). The basic requirements of $E_T^{miss} > 200$ GeV and at least 2 jets with $E_T^{jet} > 40$ GeV and $|\eta^{jet}| < 3$ are common to all the analyses.

Fig. 2 shows the five sigma discovery contours (corresponding to $S/\sqrt{S+B} = 5$) for an integrated luminosity $\int Ldt = 100 \text{ fb}^{-1}$, $A_0 = 0$, $\mu > 0$, $\tan\beta = 2$ (left plot) and 35 (right plot). The explorable regions lie below the contours. The $\mu < 0$ case is very similar. The isomass contours of some SUSY particles and countours corresponding to $\Omega h^2 = 1$ and 0.4 are also superimposed. The E_T^{miss} search, which has the best performance, allows to explore regions with $m_{\tilde{q}} \sim 2.5$ TeV. The muon isolation, pile-up and uncertainty on cross-sections effects have been investigated as well and do not show a dramatic degradation in performance.

4 Scalar Bottom and Gluino Reconstructions

The gluino and the sbottom are the strongly interacting SUSY particles which are the simplest to reconstruct. The decay $\tilde{g} \rightarrow b\bar{b}$, where $\tilde{b} \rightarrow b\tilde{\chi}_2^0$, $\tilde{\chi}_2^0 \rightarrow \tilde{\ell}\bar{\ell}(\tilde{\ell}\ell) \rightarrow \tilde{\chi}_1^0\ell^+\ell^-$ would allow to reconstruct both particles simultaneously.

In this decay two b -jets, two same flavour and opposite charge leptons and large missing transverse momentum are produced and leptons have a peculiar $\ell^+\ell^-$ invariant mass distribution with a sharp edge. If $m_{\tilde{\chi}_2^0} < m_{\tilde{\ell}} + m_{\ell}$ the $\tilde{\chi}_2^0$ decay would be a three body decay mediated by a virtual slepton and the edge would be placed at $m_{\tilde{\chi}_2^0} - m_{\tilde{\chi}_1^0}$. An example of lepton invariant mass distribution

Fig. 2. Five sigma discovery contours for various final states in $\int L dt = 100 \text{ fb}^{-1}$ for $A_0 = 0$, $\mu > 0$ and $\tan \beta = 2$ (left figure) and $\tan \beta = 35$ (right figure) [5]

is shown in the left plot of fig.3. In the case $m_{\tilde{\chi}_2^0} > m_{\tilde{t}} + m_{\ell}$ the neutralino decay is a two body decay and the edge would be placed at:

$$m_{\ell+\ell-} = \frac{\sqrt{(m_{\tilde{\chi}_2^0}^2 - m_{\tilde{t}}^2)(m_{\tilde{t}}^2 - m_{\tilde{\chi}_1^0}^2)}}{m_{\tilde{t}}}. \quad (1)$$

The point corresponding to the values of the mSUGRA parameters $m_0 = 200$ GeV, $m_{1/2} = 100$ GeV, $\tan \beta = 2$, $A_0 = 0$ and $\mu < 0$ is examined. The signal events as well as the main background events are generated with PYTHIA 6.136 and detector simulation is performed with CMSJET. The statistics corresponds to an integrated luminosity of 3 fb^{-1} .

The $\tilde{b}$ reconstruction is performed in two steps. Opposite charge leptons with an invariant mass distribution close to the edge are selected. This requirement allows to select a kinematical condition in which the leptons are emitted back to back in the $\tilde{\chi}_2^0$ reference frame. In this condition and under the assumption $m_{\tilde{\chi}_2^0} \sim 2m_{\tilde{\chi}_1^0}$, which is usually valid, the $\tilde{\chi}_2^0$ momentum is reconstructed through the formula:

$$\vec{P}_{\tilde{\chi}_2^0} = \left(1 + \frac{m_{\tilde{\chi}_1^0}}{m_{\ell+\ell-}}\right) \vec{P}_{\ell+\ell-}. \quad (2)$$

The $\tilde{\chi}_2^0$ momentum is then summed with the momentum of the highest E_T b -tagged jet and the $\tilde{b}$ can be reconstructed. The gluino is reconstructed from the sbottom momentum and that of the closest b -tagged jet.

Fig. 3. $m_{\ell^+\ell^-}$ distribution for leptons coming from $\tilde{\chi}_2^0$ decay (left side) and invariant $\tilde{b}$ and $\tilde{g}$ mass distributions (right side) at $m_0 = 200$ GeV, $m_{1/2} = 100$ GeV, $\tan\beta = 2$, $A_0 = 0$ and $\mu < 0$

The mass distributions for these mSUGRA parameters values are shown in the right plot of fig.3. The resolutions on the $\tilde{b}$ and $\tilde{g}$ masses are 32 GeV and 40 GeV respectively. The measured values of $m_{\tilde{b}}$ and $m_{\tilde{g}}$ are 263 GeV and 295 GeV which are not far from the corresponding Monte Carlo input values 275 GeV and 299 GeV.

5 Scalar Top Quark Search

The stop eigenstate with the lowest mass would be the lightest scalar quark for a large set of values of the mSUGRA parameters. It can be abundantly produced at LHC and has special decay topologies. The possibility to observe a stop signal over the Standard Model and SUSY backgrounds is here discussed[9].

The main production processes would be the $t\bar{t}$ pair production and the production from the gluino decay $\tilde{g} \rightarrow t\bar{t}$ ($t\bar{t}$). The study is concentrated on the $t\bar{t}$ process since it would be the simplest one: despite its lower cross section it would provide a simpler event topology.

The dependence of the $\tilde{t}$ mass with respect to m_0 is shown in the left plot of fig.4 for different values of $m_{1/2}$ at $\tan\beta = 10$, $\mu > 0$ and $A_0=0$: no remarkable dependence is foreseen on $\tan\beta$ and the sign of μ .

The $t\bar{t}$ production cross-section is shown in the right plot of fig.4 as a function of the $\tilde{t}$ mass. It rapidly decreases with $m_{\tilde{t}}$ and depends on other parameters only through higher order corrections[8].

The $\tilde{t} \rightarrow t\tilde{\chi}_1^0$ (with $t \rightarrow bW^\pm$) and $\tilde{t} \rightarrow b\tilde{\chi}_1^\pm$ ($\tilde{\chi}_1^\pm \rightarrow W^\pm\tilde{\chi}_1^0$) decay modes are considered in the analysis since they would dominate in most of the parameter space points examined. Both channels provide one b -jet, one $W^\pm$ and one $\tilde{\chi}_1^0$. The analysis is concentrated on $t\bar{t}$ events where one of the two $W^\pm$ (from t

Fig. 4. $\tilde{t}$ mass with respect to m_0 for different values of $m_{1/2}$ (left) and $t\bar{t}$ production cross section as a function of $m_{\tilde{t}}$ (right)[9]

or $\tilde{\chi}_1^\pm$) decays leptonically to provide lepton trigger and the other one decays hadronically to achieve high statistics.

This study is performed with PYTHIA 6.152 and detector simulation is performed with CMSJET. The analysis is split in two parts: in the first part the possibility to discriminate this signal from the known Standard Model backgrounds is investigated, while in the second part the backgrounds due to the other SUSY processes are also included.

The main Standard Model background is $t\bar{t}$ since it has the same final state particles of the signal. $W^\pm$ plus jets, Z^0 plus jets, $W^\pm W^\mp$, $W^\pm Z^0$ backgrounds are also considered. The variables which are used to perform the Standard Model backgrounds rejection are E_T^{miss} , the total transverse mass (m_T^{tot}) and the transverse mass of the highest p_T lepton- E_T^{miss} system ($m_T^{\ell\nu}$). The distributions of E_T^{miss} and $m_T^{\ell\nu}$ are shown in fig.5 for the $t\bar{t}$ signal and $t\bar{t}$ background: the distributions are broader for signal events. The Standard Model backgrounds rejection is accomplished by lower cuts on these variables.

The largest contribution to SUSY backgrounds would come from from $\tilde{q}\tilde{q}$ and $\tilde{g}\tilde{g}$ production. A large number of these events is expected to pass the $t\bar{t}$ selection because squarks and gluino would produce a large number of hadron jets, leptons and E_T^{miss} in their decays. These backgrounds can be rejected only using the general features of the event because the number of decay topologies is very large.

Since squarks and gluino are heavier than $\tilde{t}$ for almost all points in the parameter space considered, the E_T^{miss} and m_T^{tot} distributions for the SUSY backgrounds are concentrated at higher values with respect to the signal. SUSY backgrounds are reduced by upper cuts on these variables. An additional E_T cut on the hardest jet in the event is also applied because very high E_T jets are produced in the $\tilde{q} \rightarrow q\tilde{\chi}_i^0$ and $\tilde{q} \rightarrow q'\tilde{\chi}_i^\pm$ decays. The existence of a $\tilde{t}$ -like

Fig. 5. E_T^{miss} (left side) and $m_T^{l\nu}$ (right side) distributions for the $\tilde{t}\tilde{t}$ signal and the $t\bar{t}$ background at $m_0=400$ GeV, $m_{1/2}=400$ GeV, $A_0 = 0$, $\tan\beta = 10$ and $\mu > 0$ [9]

particle can thus be indicated by an excess of events in the intermediate mass range.

For each point examined cuts are tuned in order to maximise the signal significance defined by $S/\sqrt{B}$, being S and B the expected signal and background respectively. The signal significance is calculated with the Standard Model backgrounds and the Standard Model backgrounds together with the SUSY ones separately. Fig.6 shows the highest signal significance which can be obtained with this analysis for the low luminosity period ($\int Ldt = 30 \text{ fb}^{-1}$) as a function of m_0 and $m_{1/2}$ in the Standard Model background case (left) and in the Standard Model plus SUSY backgrounds case (right). The region excluded by theory and the LEP Higgs mass bound is also displayed.

The signal is visible over the Standard Model backgrounds with a significance higher than 5 for $m_{1/2} < 450$ GeV and $m_0 < 700$ GeV. At higher m_0 values the signal significance decreases because the $\tilde{t} \rightarrow \tilde{\chi}_2^0 b$ branching ratio becomes higher than that of the signal processes considered. At higher $m_{1/2}$ the low value of the significance is due to a lower cross section.

The effect of $\tan\beta$ on the expected signal is negligible since the $\tilde{t}$ mass is almost independent on it. The only difference would be observed at low $m_{1/2}$ values where the $\tilde{\chi}_1^\pm \rightarrow \tilde{\tau}\nu_\tau$ decay mode is enhanced. A different search is required in this case.

In the presence of SUSY backgrounds, the $\tilde{t}$ discovery with a signal significance of 5 would not be possible in the low luminosity period for most of the parameter space points examined. However, assuming a comparable detector performance at high luminosity, the 5-sigma discovery would be possible with an integrated luminosity of 500 fb^{-1} in most of the points. At a large A_0 , the $\tilde{t}$ would be lighter and the $\tilde{t}\tilde{t}$ cross section would be high enough to allow the 5-sigma observability even in the low luminosity period[9].

Fig. 6. $S/\sqrt{B}$ as a function of m_0 and $m_{1/2}$ for only Standard Model backgrounds (left) and Standard Model plus SUSY backgrounds (right) [9]

6 Conclusions

The CMS detector will allow the discovery of Supersymmetry in a very large range of mSUGRA parameters up to masses of 2.5 TeV. The squarks and gluino decays offer many special signatures to discriminate the SUSY processes from the Standard Model ones. Although studies are still preliminary, it will be possible in many cases to separate some signals from the overall SUSY production and perform measurements. In particular the decay $\tilde{g} \rightarrow \tilde{b}b$ would allow to reconstruct both sbottom and gluino masses. An excess of stop-like events can be observed over the Standard Model backgrounds and in some cases also over the SUSY backgrounds.

References

1. The LHC Study Group, CERN/AC/95-05
2. H.P. Nilles, Phys. Rep. **110**, 1 (1984)
3. The CMS Collaboration, CERN/LHCC 94-038
4. H. Baer, F. Paige, S. Protopopescu and X. Tata, hep-ph/9305342
5. S. Abdullin and F. Charles, Nucl. Phys. B **547**, 60 (1999)
6. T. Sjöstrand, Comp. Phys. Comm. **82**, 74 (1994)
7. S. Abdullin, A. Khanov and N. Stepanov, CMS-TN 94-180
8. W. Beenakker, M. Krämer, T. Plehn, M. Spira, P.M. Zerwas, Nucl. Phys. B **515** (1998), 3
9. G. Segneri: Scalar Top Quark Search at LHC with the CMS Detector, Ph.D. Thesis, Università di Pisa (2002)