CERN LIBRARIES, GENEVA

CM-P00061560

Ref.TH.2045-CERN

WHY IS THE DIFFRACTION PEAK A PEAK ? I

H. Cornille and A. Martin

CERN -- Geneva

ABSTRACT

We prove that the high energy differential cross-section for an elastic process has a maximum exactly in the forward direction and that the slope of the diffraction peak is at most $(\log s)^2$. We compare the width of the diffraction peaks defined by the absorptive part and the differential cross-section. Our assumptions are that the amplitude is dominated by the even signature amplitude and that the total cross-section, if it decreases, decreases less fast than $s^{-\frac{1}{2}}$. Strictly speaking our results hold only for a sequence of energies approaching infinity. The proofs are given for the spin 0 case, but it is not unreasonable to hope that they can be generalized to arbitrary spins.

I. INTRODUCTION

One of the most familiar facts in particle physics is the existence of a peak of the differential cross-sections for high energy elastic scattering exactly in the forward direction. This fact is so familiar that it is only very seldom that one questions the existence of this peak, called diffraction peak.

It has been understood long ago that the cross-section must be concentrated near the forward direction, i.e., at asymptotically small angles ¹⁾. For instance we know from consequences of local field theory and unitarity that the fixed angle amplitude is bounded by $(s^{\frac{3}{4}}(\log s)^{\frac{3}{2}})/\sqrt{\sin \Theta}$ ²⁾ if s is the square of the c.m. energy, while the forward amplitude is larger in modulus than σ_{total} . So, if the cross-section decreases less fast than $s^{-1/4}$, we see that the fixed angle amplitude becomes negligible as compared to the forward amplitude. On the other hand, if the total cross-section goes to infinity, the ratio of the fixed t amplitude to the forward one goes to zero for almost every t value ³⁾. What is more difficult to understand is that the elastic cross-section is maximum exactly at $\Theta=0$ or $t=0$ [with $t=-2k^2(1-\cos\Theta)$]. The standard argument produced is that the absorptive or imaginary part of the scattering amplitude is maximum in the forward direction, because of its positivity properties. Then it is argued that high energy scattering is dominated by shadow effects of inelastic processes and "therefore" purely imaginary. However, the logical connection between the two parts of the latter sentence is illusory. At this stage there is absolutely no need for the real part to be small.

A somewhat better argument consists in saying that if the total cross-section approaches a constant sufficiently smoothly or, more generally, has a behaviour like $(\log s)^\alpha$ and if the odd signature amplitude is negligible, Khuri and Kinoshita ⁴⁾ predict that the real part of the forward scattering amplitude is small. This is what happens in explicit models like the Regge model which are infinitely smooth by construction. However, if we do not want to tie ourselves with this super smoothness assumption, we have no guarantee that the real part will remain small. We have no guarantee that the derivative of the real part with respect to t , which controls the width of the real contribution to the diffraction peak, will be small. In fact, respectable physicists have proposed models of oscillating total cross-sections ⁵⁾ and it is desirable to know if in this case the differential cross-section is still maximum in the forward direction.

There is a second and most important problem. Suppose it is established that the cross-section has a maximum in the forward direction. Will the slope of the diffraction peak, i.e.,

$$\eta = \frac{1}{2} \frac{\partial}{\partial t} \left(\log \left(\frac{d\sigma}{d\Omega} \right) \right) \Big|_{t=0}$$

satisfy the same bounds as the corresponding quantity η_{abs} defined by the absorptive part alone ? Namely, is it true that this quantity is bounded by $(\log s)^2$? Bessis ⁶⁾ has established this result under the assumption that at all angles, the differential cross-section is less than some factor times the forward differential cross-section, but this is precisely one of the things we do not know.

Finally, we can ask ourselves a more detailed question. Stimulated by the experimental evidence of a shrinking diffraction peak ⁷⁾, theoreticians have produced models for the absorptive part of the amplitude which exhibit shrinking ⁸⁾. The question is, will the width of the differential cross-section follow the width of the absorptive part ?

We try to give an answer to all these questions, using on the one hand the known properties of the absorptive part and, on the other hand, dispersion relations for the forward amplitude and the derivative of the forward amplitude. We do not make any smoothness assumption, except continuity of the scattering amplitude with respect to energy. The only important and unpleasant assumption we have to make is that the scattering amplitude is dominated by the even signature, i.e., crossing symmetric with respect to the interchange of s and u (it will appear clearly that if odd signature contributions are smaller by a power of the energy, our results survive). Then we succeed in giving a positive answer to all the questions we have raised with the usual restriction that the results hold in an average sense or, if one prefers, for an infinite sequence of energies approaching infinity. An important finding is that the real part does not need to be small. An oscillating cross-section like $\sigma_t = A + B \sin(\lambda s)$ or a decreasing cross-section like $\sigma_t = s^{-1/5}$ will still produce a diffraction peak with all the required properties.

The paper is organized as follows : in Section II, we recall the important properties of the absorptive part that we shall use later. In Sections III and IV we establish via dispersion relations upper and lower bounds for the slope of the diffraction peak. In Section V we prove that η and η_{abs} have the same behaviour with energy if η_{abs} is sufficiently slowly varying at high energies, in a way which we specify.

II. THE ABSORPTIVE PART

The absorptive part of a spin 0-spin 0 scattering amplitude can be written

$$A_s(s, t) = \sum (2l+1) \operatorname{Im} f_l(s) P_l\left(1 + \frac{t}{2k^2}\right) \quad (1)$$

where $t = -2k^2(1 - \cos\theta)$ is negative in the physical region. Unitarity can be expressed as

$$\operatorname{Im} f_l(s) \geq \frac{2k}{\sqrt{s}} |f_l(s)|^2 > 0 \quad (2)$$

From unitarity alone, a certain number of inequalities have been obtained and in particular ⁹⁾

$$\frac{1}{2} \eta_{\text{abs.}} = \frac{\frac{d}{dt} A_s(s, t)}{A_s(s, t)} \Big|_{t=0} \geq \frac{1}{8} \left[\frac{\sigma_{\text{Tot.}}}{4\pi} - \frac{1}{k^2} \right] \quad (3)$$

with

$$A_s(s, t=0) = \frac{k\sqrt{s}}{8\pi} \sigma_{\text{Tot.}} \quad (4)$$

Therefore

$$\frac{d}{dt} A_s(s, t=0) > \text{const.} \frac{|A_s(s, t=0)|^2}{s} \quad (5)$$

This is as far as one can go using unitarity alone. If one uses the fact that $A_s(s, t)$ is analytic in a well-defined ellipse in the t plane and combines with positivity and unitarity properties, one finds that there is an effective cut-off in the partial wave distribution at $L_{\text{max}} = k/m_\pi \log s$.

Hence

$$\begin{aligned} \left. \frac{d A_s(s, t)}{dt} \right|_{t=0} &= \frac{1}{2k^2} \sum_{l=0}^{L_{\max}} (2l+1) \operatorname{Im} f_l(s) \frac{l(l+1)}{2} \\ &\approx \frac{1}{2k^2} \sum_{l=0}^{L_{\max}} (2l+1) \operatorname{Im} f_l(s) \frac{l(l+1)}{2} \\ &\leq A(s, t) \Big|_{t=0} \frac{(\log s)^2}{2m_\pi^2} \end{aligned} \quad (6)$$

Therefore we have now upper and lower bounds on η_{abs} :

$$c_1 \frac{A_s(s, t=0)}{s} < \eta_{\text{abs.}} < c_2 (\log s)^2 \quad (7)$$

C_1 and C_2 being constants which can be determined.

A qualitatively similar double inequality can be written for arbitrary spins if one replaces A by the square root of the sum of the squares of the various absorptive parts contributing to the differential cross-sections. This will be established in a forthcoming publication ¹⁰⁾.

The conclusion of this section is that if the real part of the scattering amplitude is negligible, the cross-section is indeed maximum in the forward direction. In addition, the inverse width of the diffraction peak cannot increase faster than $(\log s)^2$. This is as far as one can go without specific models.

III. REAL PART EFFECTS : HEURISTIC ARGUMENTS

The quantity on which we want to obtain information is the slope of the diffraction peak η defined by

$$\eta = \frac{1}{2} \frac{d}{d\log s} \log \left(\frac{d\sigma}{dt} \right) \Big|_{t=0} = \frac{A \frac{dA}{dt} + R \frac{dR}{dt}}{|F|^2} = \operatorname{Re} \left(\frac{\frac{dF}{dt}(s, t)}{F(s, t)} \right) \Big|_{t=0} \quad (8)$$

where $F = R + iA$.

Perhaps it is useful to present first a rather crude argument applying only to smooth functions for which the Khuri-Kinoshita theorems⁴⁾ apply. First we ask if η is positive as well as η_{abs} . If A and dA/dt are smooth, we can apply the standard results, noticing, as will appear more clearly later, that $(dF(s,t)/dt)|_{t=0}$ approximately coincides for large positive s with a crossing symmetric function in the interchange $s \rightleftharpoons u$.

Assume first

$$A \approx s (\log s)^\gamma. \quad (9)$$

From the Froissart bound $\gamma \leq 2$. From the inequalities of the previous section, we get

$$s (\log s)^{2\gamma} < \frac{dA}{dt} < s (\log s)^{\gamma+2} \quad (10)$$

so, if $dA/dt \approx s (\log s)^\beta$ we have

$$2\gamma < \beta < \gamma+2. \quad (11)$$

Then the corresponding real parts, for even signature amplitudes, are

$$R \sim \frac{\gamma\pi}{2} s (\log s)^{\gamma-1}$$

$$\frac{dR}{dt} \sim \frac{\beta\pi}{2} s (\log s)^{\beta-1}$$

If $\gamma > 0$, $R(dR/dt)$ is positive and of the order of $(\log s)^{-2}$ as compared to $A(dA/dt)$. If $\gamma < -2$, $R(dR/dt)$ is also positive because both R and dR/dt are negative. If $-2 < \gamma < 0$, $R(dR/dt)$ has no definite sign but is negligible as compared to $A(dA/dt)$. If, on the other hand, we assume for $t=0$

$$A \approx s^\alpha \quad \alpha < 1 \quad (12)$$

corresponding to a cross-section decreasing like a power of the energy, and

$$\frac{dA}{dt} \approx s^\beta \quad (13)$$

we have of course

$$2\alpha - 1 \leq \beta \leq \alpha \quad (14)$$

from inequality (5), and

$$R \frac{dR}{dt} \sim \left(\cot \alpha \cdot \frac{\pi d}{2} \right) \left(\cot \alpha \cdot \frac{\pi \delta}{2} \right) s^{\alpha + \delta} \quad (15)$$

If $\alpha > \frac{1}{2}$ we have $\delta > 0$ and $R(dR/dt) > 0$, which is what we want. If

$$\alpha \gg \frac{1}{2} + \epsilon \quad (16)$$

we get

$$\eta / \eta_{abs} < C \quad (17)$$

Notice that such a case corresponds to a rather unfamiliar behaviour $A(s, t) \sim s^\alpha f(t/s^{\alpha-\delta})$ which is not obviously forbidden by analyticity and s channel unitarity.

IV. REAL PART EFFECTS : RIGOROUS RESULTS

The strategy we shall adopt is the following : we know from Eq. (8) that

$$\eta = \operatorname{Re} \frac{\frac{d}{dt} F(s, t)}{F(s, t)} \Big|_{t=0}$$

We shall try to get upper and lower bounds on $(dF/dt)/F$ for $s \rightarrow i\infty$ by using ordinary dispersion relations separately for dF/dt and F in conjunction with the restrictions derived in Section II on dA/dt and A . Once we have these bounds, we have to make sure that they are compatible with the asymptotic behaviour of $\operatorname{Re}((dF/dt)/F)$ along the s cut. To do this we write inverse dispersion relations for $(dF/dt)/F$. The constraints will appear as conditions on the asymptotic behaviour of a regular integral involving $\operatorname{Re}((dF/dt)/F)$ when a certain parameter goes to infinity. If $(dF/dt)/F$ is assumed to be continuous, this will imply restrictions on η at least for a sequence of energies approaching infinity.

The scattering amplitude, which is supposed to be $s \rightleftharpoons u$ crossing symmetric, satisfies the dispersion relation

$$F(s, t) = C(t) + s^2 \int_{s_0}^{\infty} \frac{A(s', t) ds'}{s'^2 (s' - s)} + u^2 \int_{u_0}^{\infty} \frac{A(s', t) ds'}{s'^2 (s' - u)} \quad (18)$$

where $s+t+u = 2(M_A^2 + M_B^2) = \Sigma$, $u_0 = s_0 = (M_A + M_B)^2$. The derivative $F'(s, 0)$ can be written $dF/dt = F' = F'_S + F'_C$, where

$$F'_S(s, 0) = C'(0) + s^2 \int_{s_0}^{\infty} \frac{A'(s', 0) ds'}{s'^2 (s' - s)} + (\Sigma - s)^2 \int_{u_0}^{\infty} \frac{A'(s', 0) ds'}{s'^2 (s' - \Sigma + s)} \quad (19)$$

is even symmetric under $s \leftrightarrow u$ crossing, whereas the corrective term

$$F'_C(s, 0) = - \left[(\Sigma - s)^2 \int_{s_0}^{\infty} \frac{A'(s', 0) ds'}{s'^2 (s' - \Sigma + s)^2} + 2(\Sigma - s) \int_{s_0}^{\infty} \frac{A'(s', 0) ds'}{s'^2 (s' - \Sigma + s)} \right] \quad (20)$$

has no right-hand cut, and can be bounded for $s \rightarrow +\infty$ by using the Froissart bound. One finds

$$|F'_C(s, 0)| < (\log s)^3 \quad (21)$$

It will turn out that in all interesting situations this corrective term is negligible. Introducing now the symmetric variable

$$z = \left(\frac{s - M_A^2 - M_B^2}{2 M_A M_B} \right)^2 \quad (22)$$

we can write dispersion relations for F and F'_S

$$F(s, 0) = F(z) = F(0) + \frac{z}{\pi} \int_1^{\infty} \frac{A(x) dx}{x(x-z)} \quad (23)$$

with $A(x) = A(s, 0)$;

$$F'_s(s,0) = F'_s(z) = F'_s(0) + \frac{z}{\pi} \int_1^{\infty} \frac{A'(x) dx}{x(x-z)} \quad (24)$$

with $A'(x) = ((dA(s,t))/dt)|_{t=0}$.

We now use the double inequality (7) from Section II to get upper and lower bounds for

$$(F'_s(z) - F'_s(0)) / (F(z) - F(0))$$

for $z \rightarrow -\infty$.

a) from $\eta_{\text{abs}} < \text{const}(\log s)^2$, and

$$A(x) < \sqrt{x} (\log x)^2, \quad A'(x) < \sqrt{x} (\log x)^4$$

we get

$$\begin{aligned} |z| \int_1^{\infty} \frac{A'(x) dx}{x(x+|z|)} &< \left(|z| \int_1^{|z|} \frac{A(x) dx}{x(x+|z|)} (\log |z|)^2 + |z| \int_{|z|}^{\infty} \frac{\sqrt{x} (\log x)^4 dx}{x^2} \right) \\ &< \left[|z| \int_1^{\infty} \frac{A(x) dx}{x(x+|z|)} \right] (\log |z|)^2 + O\left(\frac{\log^4 |z|}{|z|}\right) \end{aligned}$$

Since the first bracket in the right-hand side is in any case larger than a positive constant, we get

$$\frac{F'_s(-|z|) - F'_s(0)}{F(-|z|) - F(0)} < \text{const.} (\log |z|)^2 \quad (25)$$

b) From

$$A'(x) > \text{const.} \frac{(A(x))^2}{\sqrt{x}}$$

we get, using the dispersion relations in conjunction with Schwarz inequality

$$\frac{F'_s(-|z|) - F'_s(0)}{F(-|z|) - F(0)} > \text{const.} \left[|z|^{\frac{1}{2}} \int_1^{\infty} \frac{A(x) dx}{x(x+|z|)} \right] \quad (26)$$

Let us now remark that we can get rid of the subtraction constants appearing in these inequalities if $F'_S(-|z|)$ and $F(-|z|)$ go to infinity for $|z| \rightarrow \infty$. This is the case if the integrals

$$\int \frac{A(x) dx}{x}, \quad \int \frac{A'(x) dx}{x}$$

are both divergent. From inequality (5) this is guaranteed if $A(x) > x^{1/4}$, i.e.,

$$\sigma_{TOT.} > \text{const. } s^{-1/2} \quad (27)$$

The heuristic argument of Section III shows that anyway it is hopeless to get any interesting result if σ_t decreases faster than $s^{-1/2}$.

Therefore, from (25), (26) and condition (27), we get

$$\text{const.} (\log |z|)^2 > \frac{F'_S(z)}{F(z)} > c |z|^{1/2} \int_1^\infty \frac{A(x) dx}{x(x+|z|)} \quad (28)$$

for $z_{\text{real}} < -M$.

We write now an inverse dispersion relation. Here we shall remember that $F(z)$ has at most one zero. If this zero exists, it will contribute by a pole term in the dispersion relation. In addition, using unitarity and $L_{\text{max}} = \sqrt{s \log s}$ one gets

$$\left| \frac{F'_S}{F} \right| < \left| \frac{F'_S}{A} \right| < s^{1/2} \frac{(\log s)^3}{A^{1/2}}$$

and, with condition (27)

$$\left| \frac{F'_S}{A} \right| < s^{1/4} (\log s)^3 = x^{1/8} (\log x)^3 \quad (29)$$

Therefore, we can write an unsubtracted dispersion relation for

$$\frac{1}{\sqrt{1-z}} \frac{F'_S(z)}{F(z)}$$

which reads, for $z < z_0$ (if there is a pole)

$$\frac{1}{\sqrt{1+|z|}} \frac{F'_s(-|z|)}{F(-|z|)} = \frac{c}{z_0 + |z|} + \frac{1}{\pi} \int_1^{\infty} \frac{dx}{\sqrt{x-1}} \frac{\operatorname{Re} \left(\frac{F'_s(x)}{F(x)} \right)}{x + |z|} \quad (30)$$

We now want to make sure that the behaviour of $\operatorname{Re}(F'(x)/F(x))$ for $x \rightarrow \infty$ is compatible with the behaviour of $F'_S(z)/F(z)$ for $z \rightarrow -\infty$. Remember that

$$\left| \operatorname{Re} \frac{F'(x)}{F(x)} - \operatorname{Re} \frac{F'_s(x)}{F(x)} \right| < \frac{|F'_c(x)|}{A(x)}$$

a) If $\operatorname{Re} \frac{F'(x)}{F(x)} > (\log x)^2 \nu(x)$

for $x > x_0$, where $\nu(x) \rightarrow \infty$, we get from (21) and (27) :

$$\operatorname{Re} \frac{F'_s(x)}{F(x)} > (\log x)^2 \nu(x) - \frac{(\log x)^3}{x^{1/4}}$$

The corrective term is therefore negligible.

Inserting this into the dispersion relation, we find

$$\frac{F'_s(-|z|)}{F(-|z|)} > \operatorname{const.} \nu(|z|) (\log |z|)^2$$

which contradicts the left-hand side of (28). We conclude that if $\operatorname{Re}(F'/F)$ is continuous, there is at least a sequence of energies for which

$$\eta(s_n) = \operatorname{Re} \frac{F'}{F} < \nu(s_n) (\log s_n)^2 \quad (31)$$

where $\nu(s_n)$ is a function approaching infinity arbitrarily slowly.

b) We also want to show that at least for a sequence of energies $\operatorname{Re}(F'/F) > 0$. Here a slight refinement is necessary :

$$b1) \text{ if } \int \frac{A(x)dx}{x^{3/2}} \quad \text{or} \quad \int \frac{A(s,0)ds}{s^2}$$

is convergent, we see from (19) that $|F'_C| < \text{const}$ for $x \rightarrow \infty$. Then assume

$$\operatorname{Re} \frac{F'}{F} < x^{-1/4}$$

we also get

$$\operatorname{Re} \frac{F'_s}{F} < x^{-1/4} + \frac{\text{const.}}{A} < \text{const.} x^{-1/4}$$

Inserting this into the dispersion relation, we find from (30) and the right-hand side of (28)

$$|z|^{1/2} \int \frac{A(x)dx}{x(x+|z|)} < \text{const.} |z|^{-1/4} \quad (32)$$

If we assume

$$A(x) > v(x) x^{1/4}$$

with $v(x) \rightarrow \infty$ for $x \rightarrow \infty$ we get a contradiction. We conclude that if

$$s^{1/2} \sigma_{\text{TOT}}(s) \rightarrow \infty \quad (33)$$

there exists at least a sequence of energies s_n , such that

$$\eta(s_n) > s_n^{-1/2} \quad (34)$$

b2) If $\int (A(x)dx)/x^{3/2}$ is divergent, it is no longer true that $|F'_C/F| < \text{const} x^{-1/4}$. In fact, we have

$$\left| \frac{F'_C}{F} \right| < (\log x)^3 x^{-1/4}$$

However, then the contradiction is still worse, for, assume that

$$\operatorname{Re}(F'/F) < x^{-1/4}$$

We conclude

$$\operatorname{Re} \left(\frac{F'_j}{F} \right) < x^{-1/4} (\log x)^3 < x^{-1/8}$$

for x large, and we obtain

$$|z|^{1/2} \int \frac{A(x) dx}{x(x+|z|)} < |z|^{-1/8}$$

We also have

$$\int_1^\infty \frac{A(x) dx}{x(x+|z|)} < C$$

So

$$\int_0^\infty \frac{d|z|}{|z|^{1/2}} \int_1^\infty \frac{A(x) dx}{x(x+|z|)} < C \int_0^\infty \frac{d|z|}{|z|^{1/2}} + \int_A^\infty \frac{d|z|}{|z|^{9/8}} \quad (35)$$

The double integral on the left-hand side is therefore absolutely convergent. Hence, it can be integrated in an arbitrary order and $\int_1^\infty (A(x) dx) / x^{3/2}$ is therefore convergent, which contradicts our assumption.

If we combine all these results and use continuity of $\eta(s)$ we find the following theorem :

if $s^{1/2} \sigma_{\text{total}}(s) \rightarrow \infty$ for $s \rightarrow \infty$ there exists a sequence of energies (s_n) , $s_n \rightarrow \infty$ (when $n \rightarrow \infty$) such that

$$s_n^{-1/4} < \eta(s_n) < (\log s_n)^2 \nu(s_n) \quad (36)$$

where $\nu(s_n)$ is an arbitrarily slowly increasing function of s .

This means that there is a sequence of energies for which the slope of the diffraction peak is positive and has the same upper bound as the slope of the absorptive part. Let us remark that if η is continuous, there will be around s_n an interval of values $|s - s_n| < \delta_n$ such that η is positive and has the same qualitative upper bound. It is, however, difficult to estimate the asymptotic density of these intervals.

If one wants to obtain statements involving all energies, it is perhaps better to obtain sum rules. For this purpose it seems more appropriate to work with $\log F$ and $\log F'$. What one finds is that the phases δ of $F-F(0)$ and ϕ of $F'_S-F'_S(0)$ have to be close enough in some averaged sense. This indicates that $\cos(\delta-\phi)$, as well as $\operatorname{Re}(F'_S/F)$ are positive in an averaged sense. One finds

$$\left| \int_{\text{const.}}^{\infty} w(x) \left[\frac{\delta-\phi}{\pi} - \tau(x) \right] dx \right| < \infty \quad (37)$$

with $w(x) = x^{-1}(\log x)^{-1} v(x)$, $v(x)$ increasing arbitrarily slowly

$$-\frac{1}{4} < \tau(x) < 2 \frac{\log \log x}{\log x}$$

We do not want to give here the details about the sum rules (see Appendix). The method followed is the same as in Ref. 11), the input is furnished by the bounds (28) on $F'(-|z|)/F(-|z|)$.

V. THE CASE OF η_{abs} SMOOTH : COMPARISON OF η AND η_{abs}

If η_{abs} is a slowly varying function, one can hope to get more stringent conditions on η . For instance, in the case of ordinary Regge behaviour, $\eta_{\text{abs}} = \log s$, and one expects $\eta = \log s$ too. We shall consider the following class of smooth functions Σ , $f \in \Sigma$, if

$$0 < c_p < \frac{f(x^p)}{f(x)} < C_p \quad x > x_0 \quad (38)$$

for any $p > 1$. For instance $(\log x)^n$, $\exp \sqrt{\log \log x}$, $1 + \sin x / \log \log x$, belong to the class Σ . A function belonging to this class increases and decreases less fast than any power.

We assume that η_{abs} belongs to the class Σ . Then we try to estimate

$$\frac{F'_S(-|z|)}{F(-|z|)} \approx \frac{\int_1^{\infty} \frac{\eta_{\text{abs.}}(x) A(x) dx}{x(x+|z|)}}{\int_1^{\infty} \frac{A(x) dx}{x(x+|z|)}}$$

[under the assumption that $\int (A(x)dx)/x$ and $\int (A'(x)dx)/x$ diverge]. We find using

$$A(x) < x^{\frac{1}{2}} / (\log x)^2 < x^{3/4}$$

$$\eta(x) A(x) < x^{3/4}$$

$$A(x) > x^{1/4}$$

$$\begin{aligned} \int_1^{\infty} \frac{\eta_{abs}(x) A(x) dx}{x(x+|z|)} &> \int_{|z|e}^{|z|N} \frac{\eta_{abs}(x) A(x) dx}{x(x+|z|)} \\ &> C_1 \eta_{abs}(|z|) \int_1^{\infty} \frac{A(x) dx}{x(x+|z|)} - |z|^{\frac{3\epsilon-1}{4}} - |z|^{-N/4} \end{aligned} \quad (39)$$

and

$$\int_1^{\infty} \frac{\eta_{abs}(x) A(x) dx}{x(x+|z|)} < |z|^{\frac{3\epsilon-1}{4}} + |z|^{-N/4} + C_2 \eta_{abs}(|z|) \int_1^{\infty} \frac{A(x) dx}{x(x+|z|)} \quad (40)$$

If we now use the fact that from assumption (27)

$$\int_1^{\infty} \frac{A(x) dx}{x(x+|z|)} > |z|^{-3/4}$$

and that $\eta_{abs}(|z|)$ is larger than any negative power of $|z|$ we see that we can neglect the corrective terms in (39) and (40) and get :

$$0 < C_1 \leq \frac{F'_s(-|z|)}{\eta_{abs}(|z|) F(-|z|)} \leq C_2 \quad (41)$$

or

$$0 < C_1 < \frac{\sqrt{|z|+1} \frac{1}{\pi} \int \frac{\eta(x) dx}{\sqrt{x-1} (x+|z|)}}{\eta_{abs}(|z|)} < C_2 \quad (42)$$

for $|z| \rightarrow \infty$.

This in itself is a strong constraint on $\eta(x)$ (corrective terms for lack of crossing symmetry of F' can be neglected). If a specific law for η_{abs} is assumed, say $\eta_{\text{abs}} = (\log s)^\alpha$, we find that there is a sequence of energies for which

$$0 < C_1 < \frac{\eta(s_n)}{(\log s_n)^\alpha} < C_2 \quad (43)$$

If no specific law is assumed but if we postulate that $\eta(x)$ also belongs to the class Σ , we can, as was done previously, split the integral in (42) into

$$\int_1^{|z|^\epsilon} \frac{|z|^N}{|z|^\epsilon} \int_{|z|^\epsilon}^{|z|^N} \frac{|z|^N}{|z|^\epsilon} \int_{|z|^N}^\infty \frac{|z|^N}{|z|^\epsilon}$$

and prove in this way that for $x \rightarrow \infty$

$$0 < C_1 < \frac{\eta(x)}{\eta_{\text{abs.}}(x)} < C_2 \quad (44)$$

One can restrict $\eta_{\text{abs}}(x)$ further and assume

$$\eta_{\text{abs.}}(x) \rightarrow L$$

for $x \rightarrow \infty$. Then, given ϵ we can find x_0 such that

$$|\eta_{\text{abs.}}(x) - L| < \epsilon$$

for $x > x_0$. Hence

$$\frac{F'(-|z|) - F'(0)}{F(-|z|) - F(0)} > \frac{\int_1^{x_0} \frac{A(x) \eta_{\text{abs.}}(x) dx}{x(x+|z|)} + (L-\epsilon) \int_{x_0}^\infty \frac{A(x) dx}{x(x+|z|)}}{\int_1^{x_0} \frac{A(x) dx}{x(x+|z|)} + \int_{x_0}^\infty \frac{A(x) dx}{x(x+|z|)}}$$

Then, if we assume $\int_{x_0}^\infty (A(x) dx)/x$ divergent

$$|z| \int_{x_0}^\infty \frac{A(x) dx}{x(x+|z|)} \rightarrow \infty$$

for $|z| \rightarrow \infty$ and

$$\lim_{|z| \rightarrow \infty} \frac{F'(-|z|) - F'(0)}{F(-|z|) - F(0)} \gg L - \epsilon$$

ϵ can be taken arbitrarily small. On the other hand, $F(-|z|)$ and $F'(-|z|)$ tend to infinity. Hence

$$\lim_{|z| \rightarrow \infty} \frac{F'(-|z|)}{F(-|z|)} \gg L \quad (45)$$

Majorizing now η_{abs} by $L + \epsilon$ for $x > x_0$, we get the opposite inequality and thus

$$\lim_{|z| \rightarrow \infty} \frac{F'(-|z|)}{F(-|z|)} = L \quad (46)$$

Then, from the inverse dispersion relation (30), we find a contradiction if $\eta(x) > L + \epsilon$ for $x > x_0$ or if $\eta(x) < L - \epsilon$ for $x > x_0$. The conclusion is that there is at least a sequence s_n for which $\eta(s_n) \rightarrow L$. In particular if η has a limit, this limit coincides with the limit of η_{abs} .

It is worth noticing that all this argument can be applied to $L = \infty$. More precisely if we replace $\lim \eta_{\text{abs}} = L$ by $\eta_{\text{abs}} \rightarrow \infty$ we prove $F'(z)/F(z) \rightarrow \infty$ for $z \rightarrow -\infty$, and we find a contradiction with the dispersion relation (30) if $\eta(x) < M$ for $x > 1$. We conclude that there is at least a sequence for which $\eta(s_n) \rightarrow \infty$.

Conversely if η has a limit L , the inverse dispersion relation (30) shows that

$$\lim_{z \rightarrow -\infty} \frac{F'_s(z)}{F(z)} = L$$

and thus

$$\lim_{|z| \rightarrow \infty} \frac{\int_1^\infty \frac{\eta_{\text{abs}}(x) A(x) dx}{x(x+|z|)}}{\int_1^\infty \frac{A(x) dx}{x(x+|z|)}} = L$$

implies that η_{abs} approaches L at least for a sequence of energies. In particular if $\eta \rightarrow \infty$, there is at least a sequence for which $\eta_{\text{abs}} \rightarrow \infty$.

VI. CONCLUSION

We have proved two facts about the forward diffraction peak :

- i) that it is a peak, in the sense that the cross-section is maximum at $t=0$;
- ii) that the slope of this peak cannot be arbitrarily large, i.e., less than $(\log s)^2$; like the slope of the diffraction peak defined by the absorptive part alone. If one makes more specific smoothness assumptions on the slope of the absorptive diffraction peak, one can prove that the slopes, defined by the differential cross-section and by the absorptive part, have qualitatively the same energy dependence at high energies.

Let us discuss now the assumptions. One assumption is that the total cross-section decreases less fast than $s^{-\frac{1}{2}}$. Though this is not at all an axiomatic lower bound, we find that it is quite acceptable. In fact, it is a good surprise to see that our results hold in situations where the forward real part and its derivative with respect to t are not negligible for $t=0$, for instance if $\sigma_t \sim s^{-1/4}$. It is also a good surprise to find that finite amplitude oscillations of the total cross-section around a constant value will not affect the results, in spite of the fact that the Khuri-Kinoshita theorems no longer apply.

The second assumption is more serious. It is that the odd signature amplitude is negligible. In fact all our theorems have been established assuming a pure even signature amplitude. We can certainly allow a small odd signature amplitude with an energy dependence in s^α , $\alpha \leq \frac{1}{2}$. Then if σ_t decreases less fast than $s^{-1/4}$ all the results survive. However, there is no axiomatic basis for this assumption, whose only support is of experimental origin ¹²⁾.

Finally, let us note that our theorems either hold for sequences of energies approaching infinity or appear as sum rules. This is unfortunate but unavoidable in all questions involving dispersion relations.

Let us also indicate that it is possible to show that there is a sequence of energies for which $|F(s_n, t)/F(s_n, 0)| \leq 1$ for some fixed negative t . However, it seems very difficult to prove the existence of a sequence of energies for which this holds uniformly in t $-T \leq t \leq 0$.

Our results hold for spin 0 particles. It is not hopeless to extend them to arbitrary spins. A preliminary step consists in showing that the sum of the absorptive contributions to the differential cross-section has properties very similar to the square of the absorptive part in the scalar case. This is very probably true and will be presented in a forthcoming publication¹⁰⁾.

ACKNOWLEDGEMENTS

We would like to thank B. Bonnier for his interest in this work.

A P P E N D I X

1) The positivity of $\text{Re}(F'_S/F)$ deduced from the phase representation

Let us define

$$\begin{aligned} G(z) &= F(z) - F(0), G(x+io) = |G|e^{i\delta} = A + iR_G \text{ for } x > 1 \\ G'(z) &= F'_S(z) - F'_S(0), G'(x+io) = |G'|e^{i\phi} = A' + iR_{G'} \text{ for } x > 1 \\ |G|^2 \text{Re}\left(\frac{G'}{G}\right) &= AA' + R_G R_{G'} = |G||G'| \cos(\phi - \delta) \\ \left(\frac{|F|^2}{|G||G'|}\right) \text{Re}\left(\frac{F'_S}{F}\right) &= \cos(\phi - \delta) + \frac{1}{|G||G'|} \left\{ R_G F'_S(0) + R_{G'} F(0) \right\} \end{aligned} \quad (\text{A.1})$$

1a) $\text{Re}(G'/G) > 0$ or, equivalently, $\cos(\phi - \delta) > 0$. G and G' satisfy

$$\begin{aligned} G(z) &= z B_G \exp\left(\frac{z}{\pi} \int_1^\infty \frac{\delta(x) dx}{x(x-z)}\right), B_G = \frac{\partial G}{\partial z} \Big|_{z=0} \\ G'(z) &= z B_{G'} \exp\left(\frac{z}{\pi} \int_1^\infty \frac{\phi(x) dx}{x(x-z)}\right), B_{G'} = \frac{\partial G'}{\partial z} \Big|_{z=0} \\ \Psi(z) &= \log\left(\frac{G'}{G}\right) = \log\left[\frac{\int_1^\infty \frac{A' dx}{x(x-z)}}{\int_1^\infty \frac{A dx}{x(x-z)}}\right] \\ \Psi(z) &= \log\left(\frac{B_{G'}}{B_G}\right) + \frac{z}{\pi} \int_1^\infty \frac{\phi - \delta}{x(x-z)} dx \end{aligned} \quad (\text{A.2})$$

We get from Eqs. (25) and (26) : if

$$s^{\frac{1}{2}} \sigma_{\text{TOT}}(s) \xrightarrow{s \rightarrow \infty} \infty$$

then

$$-\frac{1}{4} \log |z| < \Psi(-|z|) \leq 2 \log \log |z|, \quad (\text{A.3})$$

$|z|$ large. For $x > 1$, ϕ and δ are bounded because $A > 0$, $A' > 0$ and we furthermore assume that ϕ and δ are continuous functions.

We want to establish sum rules similar to those of Ref. 9). Let us put $z = Re^{i\theta}$ in the last Eq. (A.2). For R large, it is easy to get $|\Psi(z)| < \text{const} \log R$ if $|\theta| > \theta_0 > 0$ (θ_0 small but fixed) and $|\Psi(z)| < |\log(\theta)| \log R$ for $|\theta| < \theta_0$. We apply the Cauchy theorem to $\Psi(z)(\log z)^{-1}$ and take for the logarithmic term a cut from 0 to $-\infty$. The logarithmic determination is chosen $\log|z|$ for $z > 0$ and $\log|z| \pm i\pi$ for $z = -|z| \pm i0$. The contribution along the great circle in the z plane is bounded and we get :

$$\left| \int_{\text{const.}}^R \overline{W(x)} \left(\frac{\phi - \frac{1}{2}}{\pi} - \tau(x) \right) dx \right| < \infty$$

$$\bar{W} = x^{-1} (\log x)^{-1}, \quad -\frac{1}{4} < \tau(x) \leq 2 \frac{\log \log x}{\log x}$$

where R is arbitrarily large but finite. Let us introduce $v(x) > 0$ increasing arbitrarily slowly, and define

$$W = x^{-1} (\log x)^{-1} v^{-1}, \quad \int_{\text{const}}^{\infty} W(x) dx = \infty;$$

From the second mean value theorem we get :

$$\left| \int_{R_1}^{R_2} W(x) \left(\frac{\phi - \frac{1}{2}}{\pi} - \tau(x) \right) dx \right| = v^{-1}(R_1) \left| \int_{R_1}^{R_3} \bar{W}(x) \left(\frac{\phi - \frac{1}{2}}{\pi} - \tau(x) \right) dx \right| \\ < v^{-1}(R_1) \text{const.}$$

where R_3 is some number between R_1 and R_2 . It follows that

$$\left| \int_{\text{const}}^{\infty} dx W(x) \left(\frac{\phi - \frac{1}{2}}{\pi} - \tau(x) \right) \right| < \infty \quad (\text{A.4})$$

Because the weight function W alone leads to a divergent integral, we have in an average meaning that

$$\left\langle \frac{\phi - \frac{1}{2}}{\pi} - \tau(x) \right\rangle \rightarrow 0 \quad \text{or} \quad \langle \cos(\phi - \frac{1}{2}) \rangle > 0$$

1b) $\text{Re}(F'_S/F) > 0$. Let us consider a sequence $(s_n) \rightarrow \infty$ (when $n \rightarrow \infty$) such that

$$-\frac{1}{2} + a \leq \frac{\phi(s_n) - \frac{1}{2}}{\pi} \leq \frac{1}{2} - b \quad (\text{A.5})$$

where a, b are small positive fixed numbers. It follows that $\cos(\theta-\delta) > \lambda$, where λ is a positive fixed number. From (A.1)-(A.3) and Eq. (5), we get

$$\frac{|F|^2}{|G||G'|} \operatorname{Re}\left(\frac{F'}{F}\right) > \lambda - \left(\frac{\text{const.}}{A} + \frac{\text{const.}}{A'}\right) > \lambda - \epsilon > 0$$

for (s_n) sufficiently large.

2) Forward phase for an even symmetric s-u amplitude

It is worth while noticing that the mathematical tools developed here could be useful in other contents like, for instance, the problems of the forward phase studied previously by Khuri and Kinoshita ⁴⁾.

Firstly we want to show that if $F(z)-F(0)=G(z)$ is such that :

$$\left. \begin{aligned} G(z) &= \frac{z}{\pi} \int_1^\infty \frac{A(x) dx}{x(x-z)} = z B_G \exp \left(\frac{z}{\pi} \int_1^\infty \frac{J(x) dx}{x(x-z)} \right) \\ G(x+i0) &= |G| \exp(i\delta) = R_G + iA \text{ for } x > 1, A = x^{\frac{1}{2}} \sigma_T > 0, \\ &\quad \int \text{continuous} \\ (\log x)^{-\beta_1} &< \sigma_T(x) < (\log x)^{\beta_2}, \end{aligned} \right\} \quad (A.6)$$

then for at least one sequence $(x_n) \rightarrow \infty$ (when $n \rightarrow \infty$) and $\epsilon > 0$ arbitrarily small, we have

$$\frac{-\pi(\beta_2 + \epsilon)}{\log x_n} \leq J(x_n) - \frac{\pi}{2} \leq \frac{\pi(\beta_1 + \epsilon)}{\log x_n} \quad (A.7)$$

From (A.6) we get for $z < 0$ and $|z|$ large

$$|z|^{\frac{1}{2}} (\log |z|)^{-\beta_1} \leq G(-|z|) \leq |z|^{\frac{1}{2}} (\log |z|)^{\beta_2}$$

Let us define : $\Delta(|z|) = \frac{|z|}{\pi} \int_1^\infty \frac{J(x) dx}{x(x+|z|)}$,

$$G(-|z|) = |z| \exp(-\Delta(|z|))$$

- i) $\delta > \pi(\frac{1}{2} + (\beta_1 + \epsilon)/\log x)$ for $x > x_0$ is impossible. In that case $\Delta > \Delta_0 = \frac{1}{2}\log|z| + (\beta_1 + \epsilon)\log\log|z|$ and $G(-|z|) < |z|^{\frac{1}{2}}(\log|z|)^{-(\beta_1 + \epsilon)}$ which contradicts the left-hand side of (A.6).
- ii) $\delta < \pi(\frac{1}{2} - (\beta_2 + \epsilon)/\log x)$ for $x > x_0$ is impossible. In that case $\Delta < \Delta_1 = \frac{1}{2}\log|z| - (\beta_2 + \epsilon)\log\log|z|$ and $G(-|z|) > |z|^{\frac{1}{2}}(\log|z|)^{\beta_2 + \epsilon}$ contradicts the right-hand side of (A.6).
- iii) Taking into account the fact that δ is continuous, the result (A.7) follows.

Coming back to R_G/A we get :

$$\frac{-\pi(\beta_1 + \epsilon)}{\log x_n} \leq \frac{R_G}{A} \leq \frac{\pi(\beta_2 + \epsilon)}{\log x_n}$$

Secondly, if we define $F(z) = G(z) + F(0)$, due to $|F(0)/A| \leq x^{-\frac{1}{2}}(\log x)^\beta \rightarrow 0$ and putting $x = s^2$, $F(x) = F(s, 0) = R_F + iA$ we finally get :

if $(\log s)^{-\beta_1} \leq \sigma_T \leq (\log s)^{\beta_2}$ there exists at least one sequence $(s_n) \rightarrow \infty$ when $n \rightarrow \infty$ such that

$$-\frac{\pi}{2} \frac{(\beta_1 + \epsilon)}{\log s_n} \leq \frac{R_F}{A} \leq \frac{\pi}{2} \frac{(\beta_2 + \epsilon)}{\log s_n}$$

This result constitutes a converse theorem of one of the Jin-MacDowell's ¹³⁾ theorems. The method presented here could be useful in connection with other theorems of Refs. 4)-13).

R E F E R E N C E S

- 1) Y.S. Jin, Phys.Rev. 143, 975 (1966).
- 2) M. Froissart, Phys.Rev. 123, 1053 (1961);
A. Martin, Nuovo Cimento 42, 930 (1966).
- 3) H. Cornille and F.R.A. Simão, Nuovo Cimento 5A, 138 (1971).
- 4) N.N. Khuri and T. Kinoshita, Phys.Rev. 137, B720 (1965); 140, B706 (1965).
- 5) G.F. Chew and D.R. Snyder, Phys.Letters 31B, 75 (1970).
- 6) J.D. Bessis, Nuovo Cimento 45A, 974 (1966).
- 7) V. Bartonev et al., Phys.Rev.Letters 31, 1088 (1973).
- 8) See, e.g., A. Krzywicki, Nuclear Phys. B86, 296 (1975);
P. Grassberger and H. Miettinen, GERN preprint TH.1979 (1975).
- 9) A. Martin, Phys.Rev. 129, 1432 (1963).
- 10) H. Cornille and A. Martin - to be published.
- 11) H. Cornille and A. Martin, CERN preprint TH.1991 (1975) - to appear in
Nuclear Phys.
- 12) A.S. Carroll, Phys.Rev. 33, 928 and 932 (1975).
- 13) Y.S. Jin and S.W. MacDowell, Phys.Rev. 138, B1279 (1965).