Archives

ON THE NUMBER OF SOLUTIONS TO A CROSSING SYMMETRICNEUTRAL $\pi - \pi$ MODEL

O. Brander *)

CERN -- Geneva

A B S T R A C T

The crossing symmetry of the $\pi - \pi$ system has invited many authors to try different versions of the bootstrap hypothesis on it. In the last few years, there has been some hope that the positivity conditions of Martin, Balachandran and Nuyts, Roskies, and, most recently, the Roy physical region constraints, might be sufficient to fix the low $\pi - \pi$ partial waves with very little additional information, like the position and width of the ρ resonance. This hope was recently proved too optimistic by Basdevant, Froggatt and Petersen. A similar result was obtained by the present author in an approximately crossing symmetric, solvable model. In this paper we strengthen this latter result by determining the multiplicity of the exact solution to a crossing symmetric neutral $\pi - \pi$ model. We consider only the S wave, but the multiplicity would increase by the addition of other coupled channels. The analysis is not confined to weak coupling only, and includes all solutions, in particular also a class of logarithmically decreasing ones, which are left out by most other authors.

*) Permanent address : Institute of Theoretical Physics, Göteborg, Sweden.

1. - INTRODUCTION

Some time ago, the present author showed in a simple solvable neutral $\pi \pi$ model ¹⁾ that the positivity conditions, derived by Martin ²⁾ and a number of other authors, are not sufficient to fix the $\pi^0 \pi^0$ S wave when, say, the scattering length is given ³⁾. In fact a number of CDD poles, with arbitrary position and residue, could be added to the expression for $1/f(s)$, without violating seriously the positivity conditions.

One might argue that the model of Ref. 1) only approximately satisfies the positivity conditions, and that the result would be different in a model which satisfies the positivity conditions exactly. The obvious answer to this argument is to construct such a model, which satisfies all possible positivity conditions, and prove that its solution is not unique. This is what we shall do in the present work.

The model we shall consider is a neutral, crossing symmetric $\pi \pi$ model, which is known as the Cini-Fubini ⁴⁾ approximation for the S wave. It is obtained by taking the once subtracted Mandelstam representation with a symmetric spectral function $\varrho(s,t)$, that is

$$f(s,t,u) = \frac{s}{\pi} \rho \int_4^\infty \frac{\varrho(s',t,u) ds'}{s'(s'-s)} + \frac{t}{\pi} \rho \int_4^\infty \frac{\varrho(t',u,s) dt'}{t'(t'-t)} + \frac{u}{\pi} \rho \int_4^\infty \frac{\varrho(u',s,t) du'}{u'(u'-u)} \quad (1.1)$$

with the absorptive part

$$\varrho(s',t,u) = h(s') + \frac{t}{2\pi} \rho \int_{16}^\infty \frac{\varrho(s',t') dt'}{t'(t'-t)} + \frac{u}{2\pi} \rho \int_{16}^\infty \frac{\varrho(u',s') du'}{u'(u'-u)}, \quad (1.2)$$

and assuming that the absorptive part is well approximated by the first few terms of a partial wave expansion, in this case the first term,

$$\varrho(s,t,u) = \sum_{\ell=0}^{\infty} (2\ell+1) \text{Im} f_{\ell}(s) P_{\ell} \left(1 + \frac{2t}{s-4} \right) \approx \text{Im} f(s). \quad (1.3)$$

Adding a unitarity relation for the S wave amplitude $f(s)$, the result is a simple, crossing symmetric model, fulfilling all possible positivity conditions. Of course, it is not a realistic model, since it assumes all higher waves, and in particular the P wave, to be real. However, since crossing and unitarity are exact for the S wave, it is a good testing

ground for hypotheses about the usefulness of the positivity conditions. The model is also internally consistent, and believed to have non-trivial solutions, in contrast to the one obtained by taking also the P wave in Eq. (1.3) different from zero, which Lovelace⁴⁾ proved to have no solution.

Projecting from Eq. (1.1) the S wave results in the dispersion relation

$$\operatorname{Re} f(s) = a_0 + \frac{1}{\pi} P \int_4^\infty \left[\frac{s-s'}{(s'-4)(s'-s)} - \frac{2}{s'} + \frac{2}{s-4} \ln \frac{s'+s-4}{s'} \right] \operatorname{Im} f(s') ds',$$

or, if no subtraction is necessary,

$$\operatorname{Re} f(s) = \frac{1}{\pi} P \int_4^\infty \left[\frac{1}{s'-s} + \frac{2}{s-4} \ln \frac{s'+s-4}{s'} \right] \operatorname{Im} f(s') ds'. \quad \text{A.1}$$

For simplicity we assume that no bound states are present. To the dispersion relation we add the unitarity condition

$$\operatorname{Im} f(s) = \varrho(s) R(s) |f(s)|^2 = \varrho(s) |f(s)|^2 + \frac{1}{4\varrho(s)} [1 - \eta^2(s)] \quad \text{B}$$

to obtain a non-linear problem for the determination of $\operatorname{Re} f$ and $\operatorname{Im} f$. The functions ϱ and R or η are assumed given,

$$\varrho(s) = \sqrt{\frac{s-4}{s}}, \quad R(s) \geq 1, \quad 0 \leq \eta(s) \leq 1, \quad (1.4)$$

and the inelasticity η and the inelasticity ratio R can be expressed in each other by

$$R(s) = \frac{2 - 2\eta(s) \cos 2\delta(s)}{1 + \eta^2(s) - 2\eta(s) \cos 2\delta(s)}, \quad (1.5)$$

or

$$\eta(s) = \frac{R(s)-1}{R(s)} \cos 2\delta(s) \pm \frac{1}{R(s)} \sqrt{1 - [R(s)-1]^2 \sin^2 2\delta(s)}, \quad (1.6)$$

where $\delta(s)$ is the phase shift.

We shall also discuss the non-crossing symmetric model obtained by modifying the dispersion relation A.1 to

$$\operatorname{Re} f(s) = \frac{1}{\pi} P \int_4^{\infty} \left[\frac{1}{s' - s} + \frac{m}{s' + s - 4} \right] \operatorname{Im} f(s') ds', \quad \text{A.2}$$

or to a subtracted version of this equation. The parameter $m \geq 0$ can be varied, which will help us in the analysis. For $m = 2$ the left-hand cuts of the dispersion relations A.1 and A.2 have a similar behaviour for large s' , a fact that we shall also use.

The multiplicity of a class of exact solutions to a dispersion relation like A.1 or A.2 and the unitarity condition B has been studied by Lovelace⁵⁾. He showed that a solution $\delta(s)$ to the present problem, satisfying

$$\delta(s) - \delta(4) = \pi\nu + O(s^{-\mu}), \quad \mu > 0, \quad \text{as } s \rightarrow \infty, \quad (1.7)$$

contains 2ν arbitrary parameters, corresponding to ν CDD poles.

This result of Lovelace already answers, for solutions satisfying Eq. (1.7), the problem posed above. It shows that in order to get a unique solution for the S wave amplitude one must ;

- (i) fix the total variation of the phase shift (to $\pi\nu$) ;
- (ii) determine a number of parameters (here 2ν) from auxiliary conditions ;

besides using analyticity, unitarity and crossing symmetry.

We generalize this result to include all solutions, not only those approaching zero like an inverse power when $s \rightarrow \infty$, as Eq. (1.7) implies. We begin in Section 2 by classifying all possible solutions to the dispersion relation A.1 or A.2 and the unitarity condition B. We prove that there are only two classes of solutions, one satisfying Eq. (1.7) and the other approaching zero at infinity like an inverse power of the logarithm of s . It is the slow decrease of this latter class of solutions which makes the present study difficult.

In Section 3 we study the asymptotic form of the slowly decreasing solutions, deriving their asymptotic series in inverse powers of $\ln s$, and in Section 4 we discuss the exactly solvable case $m = 0$.

Section 5 contains the formulation of our non-linear problem in a Banach space with a norm that allows us to include all solutions. By taking the Fréchet derivative, and using the "implicit variable" theorem by Vainberg and Trenogin ^{6),5)}, we reduce the problem of the local multiplicity of the solutions of the non-linear problem to the problem of determining the index of a linear, singular integral operator.

In Section 6 we determine this index in some solvable cases, and in Section 7 we use theorems on $\emptyset$ operators by Gokhberg and Krein ⁷⁾ to include also the non-solvable cases. Finally, for the benefit of those readers who do not want to follow all the details of the argument, we present in Section 8 the main results and a short discussion. Three Appendices contain some necessary mathematical details.

2. - CLASSIFICATION OF SOLUTIONS

In this section we shall classify all possible solutions to the dispersion relations A.1 or A.2, satisfying the unitarity relation B. We begin by proving that no solution exists, which requires any subtraction.

Assume that

$$\operatorname{Im} f(s) > C (\ln s)^{-\alpha} \text{ for } s > s_a, C > 0. \quad (2.1)$$

Then the dispersion relation A.2, subtracted once, calling the subtraction constant C_0 , implies that for s large enough

$$|\operatorname{Re} f(s)| > \begin{cases} \left| \frac{C}{\pi(1+m)} \ln(\ln s) + C_0 \right| & \text{if } \alpha = 1 \\ \left| \frac{C(1+m)}{\pi(1-\alpha)} (\ln s)^{1-\alpha} + C_0 \right| & \text{if } \alpha \neq 1. \end{cases} \quad (2.2)$$

This follows from the asymptotic analysis of Appendix 1, Eq. (A.1.9). The same result, with $m = 2$, can be proved in the case of dispersion relation A.1.

Now, unitarity demands

$$[\operatorname{Re} f(s)]^2 \leq \frac{1}{g(s)} \operatorname{Im} f(s) \leq \frac{1}{g(s)}, \quad (2.3)$$

which clearly excludes the growing behaviour of $\operatorname{Re} f(s)$ for $\alpha \leq 1$.

Assuming also that

$$\operatorname{Im} f(s) < C (\ln s)^{-\alpha+\varepsilon} \text{ for } s > s_m, \varepsilon > 0, \quad (2.4)$$

which is that the asymptotic behaviour is close to the lower limit in Eq. (2.1), we get from Eq. (2.3) that

$$\left[\frac{C(1+m)}{\pi(1-\alpha)} (\ln s)^{1-\alpha} + C_0 \right]^2 < C (\ln s)^{-\alpha+\varepsilon},$$

which, since we know already that $\alpha > 1$, implies that $C_0 = 0$ and $\alpha > 2 - \varepsilon$. This proves that any solution $\operatorname{Im} f(s)$ to the dispersion relations A.1 or A.2, satisfying the unitarity relation B must be asymptotically smaller than $(\ln s)^{-2+\varepsilon}$, $\varepsilon > 0$. Thus no subtraction is necessary.

From unitarity we also have

$$\begin{aligned} 0 \leq 1 - \eta^2(s) &= 4g(s) \left\{ \operatorname{Im} f(s) - g(s) [\operatorname{Re} f(s)]^2 - g(s) [\operatorname{Im} f(s)]^2 \right\} \rightarrow \\ &\xrightarrow{s \rightarrow \infty} -4g^2(s) [\operatorname{Re} f(s)]^2 \leq 0 \end{aligned} \quad (2.5)$$

as soon as $\operatorname{Im} f(s) \rightarrow 0$ as $s \rightarrow \infty$. Thus, in our model,

$$\operatorname{Re} f(\infty) = 0 \text{ and } \eta(\infty) = 1. \quad (2.6)$$

Next, we shall prove that if a solution exists, behaving asymptotically as an inverse logarithm, then its leading asymptotic behaviour is completely fixed. We thus assume that

$$\operatorname{Im} f(s) \underset{s \rightarrow \infty}{\sim} C (\ln s)^{-\alpha}, \quad C \neq 0, \quad (2.7)$$

where we know from above that $\alpha > 2 - \varepsilon$. Then, according to Appendix 1, the dispersion relation A.2 implies

$$\operatorname{Re} f(s) \underset{s \rightarrow \infty}{\sim} \frac{C(1+m)}{\pi(\alpha-1)} (\ln s)^{1-\alpha}. \quad (2.8)$$

In the case of A.1, the same formula with $m = 2$ is valid.

We now employ unitarity, assuming for the inelasticity ratio $R(s)$ the asymptotic form

$$R(s) \underset{s \rightarrow \infty}{\sim} R_0 (\ln s)^\gamma, \quad \gamma \geq 0. \quad (2.9)$$

This gives a reasonable generality to our results, since experimentally, $R(s)$ seems to stay bounded. For the inelasticity η , the assumption (2.9) corresponds to an approach to 1 with a rate determined by the relation

$$\frac{1 - \eta^2(s)}{4g(s)\operatorname{Im} f(s)} \underset{s \rightarrow \infty}{\sim} 1 - \frac{1}{R_0} (\ln s)^{-\gamma}, \quad \gamma \geq 0. \quad (2.9')$$

This quantity is restricted, by the unitarity relation, to be between 0 and 1.

With the assumptions (2.9) or (2.9'), unitarity gives

$$C (\ln s)^{-\alpha} \underset{s \rightarrow \infty}{\sim} R_0 (\ln s)^\gamma \frac{C^2(1+m)^2}{\pi^2(\alpha-1)^2} (\ln s)^{2-2\alpha},$$

which implies

$$\alpha = 2 + \gamma, \quad C = \frac{\pi^2 (1+\gamma)^2}{R_0 (1+m)^2}, \quad (2.10)$$

that is, a completely fixed leading asymptotic term :

$$\operatorname{Im} f(s) \underset{s \rightarrow \infty}{\sim} \frac{\pi^2 (1+\gamma)^2}{R_0 (1+m)^2} (\ln s)^{-2-\gamma}; \quad \operatorname{Re} f(s) \underset{s \rightarrow \infty}{\sim} \frac{\pi (1+\gamma)}{R_0 (1+m)} (\ln s)^{-1-\gamma} \quad (2.11)$$

We shall call any solution fulfilling Eq. (2.11) a class 1 solution.

These solutions will be studied in great detail in the following, but first we want to see what other solutions there might be. Therefore, let us assume that

$$C s^{-\beta-\varepsilon} < \operatorname{Im} f(s) < C s^{-\beta+\varepsilon} \quad \text{for } s > s_a, \quad (2.12)$$

with C , β and ε positive constants, $\beta - \varepsilon > 0$.

Up to now, the large s behaviour of $\operatorname{Re} f(s)$ has been determined by the large s behaviour of $\operatorname{Im} f(s)$. However, for β large enough in Eq. (2.12) this is no longer true. Take the dispersion relation A.2 and let $s \gg s_a$. Then the contribution to $\operatorname{Re} f(s)$ from $\operatorname{Im} f(s')$ for $s' < s_a$ is

$$\frac{1}{\pi} \int_4^{s_a} \left[\frac{1}{s'-s} + \frac{m}{s'+s-4} \right] \operatorname{Im} f(s') ds' = \frac{C_a}{s} + O\left(\frac{s_a}{s^2}\right), \quad (2.13)$$

where C_a is the constant

$$C_a = \frac{m-1}{\pi} \int_4^{s_a} \operatorname{Im} f(s') ds', \quad (2.14)$$

which is different from zero, except when $m = 1$.

If we take the dispersion relation A.1 the situation is slightly more complicated. On the left-hand cut the discontinuity at $s = s_b$ is given by

$$\text{Im} f(s_b) = -\frac{2}{4-s_b^4} \int_{s_b}^{4-s_b} \text{Im} f(s') ds', \quad s_b < 0. \quad (2.15)$$

For fast decreasing $\text{Im} f(s')$, say $\beta > 1+\epsilon$ in Eq. (2.12), the left-hand cut discontinuity will not decrease faster than s_b^{-1} ,

$$\text{Im} f(s_b) = \frac{\pi C_b}{s_b} + o\left(\frac{1}{s_b}\right) \text{ as } s_b \rightarrow -\infty, \quad (2.16)$$

with a strictly positive constant C_b ,

$$C_b = \frac{2}{\pi} \int_4^{\infty} \text{Im} f(s') ds'. \quad (2.17)$$

No matter how fast $\text{Im} f(s')$ decreases as $s' \rightarrow \infty$, the left-hand cut will thus contribute to $\text{Re} f(s)$ with a term

$$C_b \int_{-\infty}^{-s_b} \frac{ds_b}{s_b(s_b-s)} = \frac{C_b \ln s}{s} + o\left(\frac{1}{s \ln s}\right) \quad (2.18)$$

as $s \rightarrow \infty$, according to Eq. (A.1.11).

Thus, for $\beta > 1+\epsilon$, Eq. (2.12) implies that

$$\text{Re} f(s) \underset{s \rightarrow \infty}{\sim} \begin{cases} \frac{C_b \ln s}{s} & \text{in the case A.1} \\ \frac{C_m}{s} & \text{" " " A.2 for } m \neq 1 \\ o\left(\frac{1}{s^2}\right) & \text{" " " A.2 for } m = 1 \end{cases} \quad (2.19)$$

But then unitarity forces upon $\text{Im} f(s)$ the behaviour

$$\text{Im} f(s) \underset{s \rightarrow \infty}{\sim} \begin{cases} \frac{R_0 C_b^2 (\ln s)^{2+\gamma}}{s^2} & \text{in the case A.1} \\ \frac{R_0 C_m^2 (\ln s)^\gamma}{s^2} & \text{" " " A.2 for } m \neq 1 \\ o\left(\frac{(\ln s)^\gamma}{s^4}\right) & \text{" " " A.2 for } m = 1 \end{cases} \quad (2.20)$$

Any solution satisfying Eqs. (2.19) and (2.20) we shall call a class 2 solution. In contrast to class 1 solutions, the coefficient of its leading asymptotic term contains low energy contributions, as we have seen from its construction. Since the constants C_a and C_b are different from zero (except for $m = 1$), the leading term has a non-zero coefficient, and no solution can exist, having a faster decrease than the class 2 solutions.

What about the possibility of solutions which decrease faster than class 1 but slower than class 2? Assume again that Eq. (2.12) is valid, with some $\beta \in (0, 2)$. Then the dispersion relation A.2 gives, with the help of Eq. (A.1.11), that for sufficiently large s ,

$$\frac{(1+m)|C_b|C}{\pi s^{\beta+\varepsilon}} + O\left(\frac{1}{s}\right) < |\operatorname{Re} f(s)| < \frac{(1+m)|C_b|C}{\pi s^{\beta-\varepsilon}} + O\left(\frac{1}{s}\right). \quad (2.21)$$

We have here chosen ε such that $\beta + \varepsilon$ and $\beta - \varepsilon$ are non-integers. On the other hand, the unitarity relation gives that

$$|\operatorname{Re} f(s)| = \left[\frac{\operatorname{Im} f(s)}{g(s)R(s)} \right]^{1/2} \left[1 - g(s)R(s)\operatorname{Im} f(s) \right]^{1/2} \underset{s \rightarrow \infty}{\sim} \left[\frac{\operatorname{Im} f(s)}{R(s)} \right]^{1/2}, \quad (2.22)$$

which, together with Eq. (2.12) implies that

$$R_0^{-1/2} C^{1/2} s^{-\frac{\beta+\varepsilon}{2}} (\ln s)^{-\frac{\gamma}{2}} < |\operatorname{Re} f(s)| < R_0^{-1/2} C^{1/2} s^{-\frac{\beta-\varepsilon}{2}} (\ln s)^{-\frac{\gamma}{2}}. \quad (2.23)$$

The allowed ranges (2.21) and (2.23) of values for $|\operatorname{Re} f(s)|$ will not overlap if $3\varepsilon < \beta < 1 + \varepsilon$, which shows that there are no solutions with $\beta > 0$ except the class 2 solutions.

The only possibility now left of having a solution outside the classes 1 and 2 is a solution decreasing slower than any inverse power of s , but faster than any inverse power of $\ln s$,

$$C s^{-\varepsilon} < \operatorname{Im} f(s) < C (\ln s)^{-N} \text{ as } s \rightarrow \infty, \quad (2.24)$$

for any $\varepsilon > 0$ and $N < \infty$. We saw above that for such a solution, the asymptotic form of $\operatorname{Re} f(s)$ is determined by the asymptotic form of $\operatorname{Im} f(s)$, without finite energy contributions. But then, as we saw in Appendix 1, the leading contribution to the dispersion integral is

$$\operatorname{Re} f(s) \underset{s \rightarrow \infty}{\sim} C_1 \int_s^\infty \frac{ds'}{s'} \operatorname{Im} f(s'), \quad C_1 = \frac{1+m}{\pi} > 0. \quad (2.25)$$

Now, if Eq. (2.24) is valid,

$$\max_{s \leq s' < \infty} \{(\ln s')^2 \operatorname{Im} f(s')\} = (\ln s)^2 \operatorname{Im} f(s) \quad (2.26)$$

for sufficiently large s . Thus

$$\operatorname{Re} f(s) \underset{s \rightarrow \infty}{\leq} C_1 (\ln s)^2 \operatorname{Im} f(s) \int_s^\infty \frac{ds'}{s' (\ln s')^2} = C_1 \ln s \operatorname{Im} f(s). \quad (2.27)$$

Putting this into Eq. (2.22), squared, gives

$$C_1^2 (\ln s)^2 [\operatorname{Im} f(s)]^2 \underset{s \rightarrow \infty}{\gtrsim} \frac{\operatorname{Im} f(s)}{R(s)}, \quad (2.28)$$

that is

$$\operatorname{Im} f(s) \underset{s \rightarrow \infty}{\gtrsim} \frac{1}{C_1^2 (\ln s)^2 R(s)}. \quad (2.29)$$

This is certainly not in the interval (2.24), but is a class 1 behaviour.

Thus we can conclude that there are no solutions to dispersion relation

A.1 or A.2 and unitarity condition B outside the classes 1 and 2.

3. - ASYMPTOTIC SERIES FOR CLASS 1 SOLUTIONS

We find the fact that class 1 solutions have a fixed leading asymptotic term quite interesting. Therefore, we shall in this section derive their complete asymptotic series in inverse powers of $\ln s$.

Consider any class 1 solution, and assume that it has an asymptotic expansion

$$\text{Im } f(s) \sim \sum_{n=0}^{\infty} \frac{c_n}{(\ln s)^{2+\gamma+n}} ; \quad \text{Re } f(s) \sim \sum_{n=0}^{\infty} \frac{d_n}{(\ln s)^{1+\gamma+n}}. \quad (3.1)$$

As before, $\gamma \geq 0$ comes from the asymptotic form of the inelasticity ratio $R(s)$, for which we also assume an asymptotic series

$$R(s) \sim \sum_{i=0}^{\infty} R_i (\ln s)^{\gamma-i}. \quad (3.2)$$

To the assumption that an asymptotic expansion (3.1) exists we will have reason to come back later. In particular, if γ were non-integer, we would have to include into the sum also terms with $m\gamma$, m integer > 1 , in the exponent of $\ln s$. For simplicity, we therefore assume γ integer in this Section.

Let us write the dispersion relation A.1 in the form ($s > 0$)

$$\text{Re } f(s) = \frac{1}{\pi} P \int_4^{\infty} \frac{\text{Im } f(s') ds'}{s' - s} + \frac{1}{\pi} \int_{-\infty}^0 \frac{\text{Im } f(s') ds'}{s' - s}, \quad (3.3)$$

where for $s' < 0$

$$\text{Im } f(s') = -\frac{2}{4-s'} \int_4^{4-s'} \text{Im } f(s'') ds'' \sim -\frac{2}{4-s'} \sum_{n=0}^{\infty} \frac{c_n}{4} \frac{ds''}{(\ln s'')^{2+\gamma+n}}. \quad (3.4)$$

The rest term of the last series will contain contributions from

- (i) finite $s'' \Rightarrow O\left(\frac{1}{s}\right)$
- (ii) rest term of (3.1) $\Rightarrow \frac{1}{s} \int_4^{4-s'} \frac{ds''}{(\ln s'')^N} = O\left(\frac{1}{(\ln s')^N}\right)$, N arbitrary.

Now

$$\int_4^{4-s'} \frac{ds''}{(\ln s'')^{2+\gamma+m}} \sim \sum_{k=0}^{\infty} \frac{(m+k+1+\gamma)!}{(m+1+\gamma)!} \frac{4-s'}{[\ln(4-s')]^{2+\gamma+m+k}}, \quad (3.5)$$

as is easily obtained by partial integrations. This gives for $\text{Im } f(s')$ on the left-hand cut the asymptotic series

$$\text{Im } f(s') \sim \sum_{n,k=0}^{\infty} \frac{(m+k+1+\gamma)!}{(m+1+\gamma)!} \frac{c_n}{[\ln(4-s')]^{2+\gamma+m+k}}$$

or

$$\text{Im } f(s') \sim \sum_{n=0}^{\infty} \frac{c_n}{[\ln(4-s')]^{2+\gamma+m}}, \quad (3.6)$$

with

$$c_n = \sum_{m=0}^n \frac{(m+1+\gamma)!}{(n-m+1+\gamma)!} 2 c_{n-m}. \quad (3.7)$$

Now we go on by substituting the asymptotic series for $\text{Im } f(s')$ on the right and left-hand cuts, respectively, into the two integrals of Eq. (3.3). Besides the question of controlling the rest term of the asymptotic expansion, we have to be careful with the singularity of the Cauchy integral.

In Appendix 1 we study this problem in detail, and show that for each term in the asymptotic series (3.1) or (3.6) we get an asymptotic expansion for the corresponding integral in Eq. (3.3).

Substitution of the asymptotic series (3.1) into the first integral of Eq. (3.3) gives

$$\frac{1}{\pi} P \int_4^{\infty} \frac{\text{Im } f(s') ds'}{s'-s} \sim \sum_{n=0}^{\infty} c_n \frac{1}{\pi} P \int_4^{\infty} \frac{ds'}{(s'-s)(\ln s')^{2+\gamma+m}}. \quad (3.8)$$

The rest term of this series contains contributions from

- (i) finite $s' \Rightarrow \mathcal{O}(\frac{1}{s})$
- (ii) rest term of (3.1) $\Rightarrow \int_4^{\infty} \frac{ds'}{(s'-s)(\ln s')^N} = \mathcal{O}(\frac{1}{(\ln s)^{N-1}})$, N arbitrary.

Using Eq. (A.1.7) for each of the integrals on the right-hand side of Eq. (3.8), and rearranging the terms, we now get the asymptotic series

$$\frac{1}{\pi} p \int_4^{\infty} \frac{\text{Im} f(s') ds'}{s'-s} \sim \sum_{n=0}^{\infty} \left[\frac{c_n}{1+\gamma+n} - \sum_{k=0}^{\left[\frac{n-\gamma}{2}\right]} \frac{2(n+\gamma)! p_k^+ c_{n-2k-2}}{(n+\gamma-2k-1)!} \right] \frac{1}{\pi (\ln s)^{1+\gamma+n}} \quad (3.9)$$

with a rest term that is $\mathcal{O}((\ln s)^{-n})$, any n , as $s \rightarrow \infty$. The constants $p_k^{\pm}$ are defined in Appendix 1.

For the second integral of Eq. (3.3), we use Eq. (3.6), change the integration variable, and then use Eq. (A.1.3) to get an expression very similar to Eq. (3.9). This expression, together with Eq. (3.9), gives the following result for the coefficients d_n of the asymptotic series for $\text{Re} f(s)$:

$$d_n = \frac{1}{\pi} \left[\frac{c_n + e_n}{1+\gamma+n} - \sum_{k=0}^{\left[\frac{n-\gamma}{2}\right]} \frac{2(n+\gamma)!}{(n+\gamma-2k-1)!} (p_k^+ c_{n-2k-2} - p_k^- e_{n-2k-2}) \right] \quad (3.10)$$

This formula gives the connection between the asymptotic forms of $\text{Im} f(s)$ and $\text{Re} f(s)$, implied by the dispersion relation A.1. For the case of dispersion relation A.2, the same formula is obtained, but with $e_n = m c_n$ instead of the expression (3.7).

The unitarity condition B also gives a connection between the asymptotic forms of $\text{Im} f(s)$ and $\text{Re} f(s)$. This connection is ($c_{-1} = 0$)

$$c_n = \sum_{i=0}^n \sum_{j=0}^{n-i} (d_i d_{n-i-j} + c_{i-1} c_{n-i-j-1}) R_j. \quad (3.11)$$

For given R_j , we could, hopefully, solve Eqs. (3.10) and (3.11) for c_n and d_n . For $n = 0$ we get

$$c_0 = \frac{\pi^2(1+\gamma)^2}{9R_0}, \quad d_0 = \frac{\pi(1+\gamma)}{3R_0}, \quad (3.12)$$

as we had in Section 2 for class 1 solutions. Similarly, we get for $n = 1$

$$c_1 = -\frac{\gamma+2}{\gamma} \left[\frac{4\pi^2(1+\gamma)^3}{27R_0} + \frac{\pi^2(1+\gamma)^2 R_1}{9R_0^2} \right]. \quad (3.13)$$

Provided $\gamma \neq 0$, this is a well-defined solution and we can go on to $n = 2, 3$, etc., finding

$$c_n = \frac{n+\gamma+1}{n-\gamma-1} \left[\sum_{j=1}^n d_0 d_{n-j} R_j + \sum_{i=1}^{n-1} \sum_{j=0}^{n-i} (d_i d_{n-i-j} + c_{i-1} c_{n-i-j}) R_j + \right. \\ \left. + \sum_{i=1}^n \frac{4(1+\gamma) c_{n-i} (n+\gamma)!}{3(n+\gamma-i+1)!} - \sum_{h=0}^{\left[\frac{n-2}{2}\right]} \frac{4(1+\gamma)(n+\gamma)!}{3(n+\gamma-2h-1)!} (P_h^+ c_{n-2h-2} - P_h^- c_{n-2h-2}) \right], \quad (3.14)$$

until $n = 1+\gamma$. For this value of n , however, no solution exists for c_n , unless the expression in square brackets happens to be zero! This gives a condition on one (but only one!) of the coefficients R_j of the asymptotic series (3.2). The condition is that $R_{1+\gamma}$ should be the following combination of $R_0, R_1, \dots, R_\gamma$:

$$R_{1+\gamma} = -\frac{1}{d_0^2} \left[\sum_{j=0}^{\gamma} d_0 d_{1+\gamma-j} R_j + \sum_{i=1}^{\gamma} \sum_{j=0}^{1+\gamma-i} (d_i d_{1+\gamma-i-j} + c_{i-1} c_{\gamma-i-j}) R_j + \right. \\ \left. + \sum_{i=1}^{1+\gamma} \frac{2 c_{1+\gamma-i} (2+2\gamma)!}{3(2+2\gamma-i)!} - \sum_{h=0}^{\left[\frac{\gamma-1}{2}\right]} \frac{2(2+2\gamma)!}{3(2\gamma-2h)!} (P_h^+ c_{\gamma-2h-1} - P_h^- c_{\gamma-2h-1}) \right] \quad (3.15)$$

For $\gamma = 0$, this simplifies to $R_1 = -\frac{4}{3}R_0$.

If $R_{1+\gamma}$ is given by this expression, the corresponding coefficient $c_{1+\gamma}$ is a free parameter. The higher coefficients, c_n for $n > 1+\gamma$, are given by Eq. (3.14) and contain the free parameter $c_{1+\gamma}$.

Provided Eq. (3.15) is fulfilled, we have thus found the complete asymptotic series (3.1) for an arbitrary class 1 solution, subject only to the condition that an expansion of the form (3.1) exists. The asymptotic series contains one arbitrary parameter, but is otherwise fixed. We will later be able to connect this parameter to the scattering length.

We should at this point not overstress the importance of the condition (3.15), because, as will become clear in the next section, it is more connected to the existence of an asymptotic expansion (3.1) than to the question of the existence for the exact solution.

For the case of dispersion relation A.2, the same reasoning as above applies, but the formulae differ somewhat. For $n = 0$ we get

$$c_0 = \frac{\pi^2 (1+\gamma)^2}{R_0 (1+m)^2}, \quad d_0 = \frac{\pi (1+\gamma)}{R_0 (1+m)}, \quad (3.16)$$

and for $n > 0$ we get Eq. (3.14), but without the third sum. Also in Eq. (3.15) the third sum drops out, and the condition for $\gamma = 0$ becomes simply $R_1 = 0$.

4. - SOLUTION OF THE RIGHT-HAND CUT EQUATION

As is well known, our non-linear problem could be easily solved with the Castillejo-Dalitz-Dyson ⁸⁾ method if the left-hand cut were absent, that is for $m = 0$ in the dispersion relation A.2. This exact solution is of great interest to us in this work, since, as we will prove, the gross features of it cannot be changed by the left-hand cut.

The exact solution for $m = 0$ is

$$\frac{1}{f(s)} = \frac{1}{a_0} - \beta_0(s-4) - \sum_{n=1}^{\infty} \frac{\beta_n(s-4)}{(\alpha_n-4)(\alpha_n-s)} - \frac{s-4}{\pi} \int_4^{\infty} \frac{\rho(s')R(s')ds'}{(s'-4)(s'-s)}, \quad (4.1)$$

where a_0 is the scattering length, and α_n is the position and β_n the residue of the n^{th} CDD pole, β_0 corresponding to a CDD pole at infinity. These constants have to be real and satisfy

$$\begin{cases} 0 < a_0 < \infty \\ 4 < \alpha_n < \infty, \quad n = 1, 2, \dots \\ 0 \leq \beta_n < \infty, \quad n = 0, 1, 2, \dots \end{cases} \quad (4.2)$$

Let us first look at the asymptotic behaviour of the solutions (4.1). If $\beta_0 \neq 0$ it is obvious that the term $-\beta_0 s$ dominates for large s , i.e., that

$$\text{Re} \frac{1}{f(s)} = \frac{\text{Re} f(s)}{|f(s)|^2} \underset{s \rightarrow \infty}{\sim} -\beta_0 s. \quad (4.3)$$

This clearly corresponds to a class 2 solution

$$\text{Re} f(s) \underset{s \rightarrow \infty}{\sim} -\frac{1}{\beta_0 s}, \quad \text{Im} f(s) \underset{s \rightarrow \infty}{\sim} \frac{R_0 (\ln s)^{\gamma}}{\beta_0^2 s^2}, \quad (4.4)$$

according to Eqs. (2.19) and (2.20), case A.2 with $m = 0$.

If $\beta_0 = 0$ the integral will dominate the right-hand side of Eq. (4.1) for large s . According to Appendix 1, Eq. (A.1.9), we have

$$-\frac{s-4}{\pi} \rho \int_4^{\infty} \frac{\rho(s')R(s')ds'}{(s'-4)(s'-s)} \underset{s \rightarrow \infty}{\sim} \frac{R_0}{\pi(1+\gamma)} (\ln s)^{1+\gamma}. \quad (4.5)$$

This corresponds to a class 1 solution with

$$\operatorname{Re} f(s) \underset{s \rightarrow \infty}{\sim} \frac{\pi(1+\gamma)}{R_0} (\ln s)^{-1-\gamma}, \quad \operatorname{Im} f(s) \underset{s \rightarrow \infty}{\sim} \frac{\pi^2(1+\gamma)^2}{R_0} (\ln s)^{-2-\gamma}, \quad (4.6)$$

in accordance with Eq. (2.11) with $m = 0$. Thus the absence or presence of a CDD pole at infinity determines whether the solution is of class 1 or class 2.

Let us examine in some detail the asymptotic form of class 1 solutions, in order to connect to the asymptotic analysis of the preceding section. Taking $\gamma = 0$ for simplicity, we obtain from Eq. (A.1.9) the next term of Eq. (4.5) :

$$-\frac{s-4}{\pi} P \int_4^\infty \frac{g(s') R(s') ds'}{(s'-4)(s'-s)} = \frac{R_0}{\pi} \ln s + \frac{R_1}{\pi} \ln(\ln s) + O(1) \text{ as } s \rightarrow \infty. \quad (4.5')$$

Thus, for $R_1 \neq 0$ we would get

$$\operatorname{Re} f(s) = \frac{\pi}{R_0} (\ln s)^{-1} - \frac{\pi R_1}{R_0^2} \frac{\ln(\ln s)}{(\ln s)^2} + O\left(\frac{1}{(\ln s)^2}\right) \text{ as } s \rightarrow \infty. \quad (4.7)$$

This clearly shows why a condition like Eq. (3.15) is necessary in order for an asymptotic expansion like Eq. (3.1) to exist. As we mentioned at the end of Section 3, this condition is simply $R_1 = 0$ for the case studied here.

The last term of Eq. (4.5') contains among other contributions the constant $1/a_0$. This constant will, therefore, influence the coefficient of $(\ln s)^{-2}$ in the expansion for $\operatorname{Re} f(s)$, that is d_1 in Eq. (3.1). Through the unitarity condition it is then connected to the coefficient c_1 . Without going into details, this shows that there is a connection between the scattering length and the arbitrary parameter of the asymptotic expansions of Section 3.

For elastic unitarity, we can easily trace the variation of the phase shift for the solution (4.1) through the formula

$$\cot \delta(s) = \frac{1}{g(s)} \operatorname{Re} \frac{1}{f(s)}. \quad (4.8)$$

Assuming first that all $\beta_n = 0$ for $n \geq 1$, we find from Eq. (4.1) that if $\beta_0 > 0$, $\text{Re}(1/f)$ is positive at threshold, decreases through zero and reaches $-\infty$ at $s = \infty$. This corresponds to an increase of $\delta(s)$ from zero through $\pi/2$ to π at $s = \infty$. If instead $\beta_0 = 0$, $\text{Re}(1/f)$ stays positive and approaches $+\infty$ at $s = 0$. This is so since for elastic unitarity, the integral can be easily computed, and shown to give a positive contribution for all s . For $\gamma > 0$, on the other hand, the integral may change sign, depending on the actual form of $R(s)$, so we cannot draw such definite conclusions.

For elastic unitarity, and the sum in Eq. (4.1) absent, we thus have

$$\delta(\infty) - \delta(4) = \begin{cases} 0 & \text{for } \beta_0 = 0, \beta_n = 0, n \geq 1 \\ \pi & \text{" } \beta_0 \neq 0, \beta_n = 0, n \geq 1. \end{cases} \quad (4.9)$$

Adding now CDD poles, we find that each pole contributes one zero and one infinity to $\text{Re}(1/f)$. Since the residue is negative, this means that each CDD pole adds π to $\delta(\infty)$.

Consider now a particular solution, for which

$$\delta(4) = 0, \delta(\infty) = \pi\nu, \nu \text{ integer.} \quad (4.10)$$

Then the analysis above shows that this solution has exactly ν CDD poles (counting also the one at infinity), and that its multiplicity κ , that is the number of arbitrary constants in it, is given by

$$\kappa = \begin{cases} 2\nu + 1 & \text{for class 1 solutions} \\ 2\nu & \text{for class 2 solutions} \end{cases} \quad (4.11)$$

This result is now proved for elastic unitarity and $m = 0$, and we go on to generalize it to inelastic unitarity, to $m \neq 0$ and to the case of dispersion relation A.1.

5. - BANACH SPACE FORMULATION OF THE NON-LINEAR PROBLEM

The non-linear problem of finding solutions to the dispersion relation A.1 or A.2 and the unitarity condition B is best discussed within the framework of mappings of Banach spaces with suitable norms ^{5),9)}.

We have chosen to work in a Banach space of Hölder continuous functions on $4 \leq s < \infty$, which, however, in order to allow for class 1 solutions, do not have to obey the Hölder condition at $s = \infty$. As $s \rightarrow \infty$ they are only requested to approach a limiting value at least as fast as $(\ln s)^{-k}$, $k \geq 1$.

To be more explicit, we put

$$\mu = 1 - \frac{4}{s} \quad (5.1)$$

and let $H(\mu, k)$ be the space of Hölder continuous functions $\varphi(u)$ on $u \in [0, 1)$, including $u = 0$ but excluding $u = 1$, with the norm

$$\|\varphi\| = \sup_{u \in [0, 1]} |\varphi(u)| + \sup_{u_1, u_2 \in [0, 1]} \frac{|\varphi(u_1) - \varphi(u_2)|}{\left| \frac{u_1 - u_2}{1 - u_-} \right|^\mu} \left(\ln \frac{4}{1 - u_-} \right)^k, \quad (5.2)$$

where $0 < \mu < 1$, $k \geq 1$ and $u_- = \min(u_1, u_2)$. That this is a Banach space, and even a normed ring, is proved in Appendix 2.

We shall also use the subspace $H_0(\mu, k)$, consisting of all the functions of $H(\mu, k)$ that are zero at $u = 1$, $\varphi(1) = 0$.

In $H_0(\mu, k)$ we use the simpler norm

$$\|\varphi\| = \sup_{u_1, u_2 \in [0, 1]} \frac{|\varphi(u_1) - \varphi(u_2)|}{\left| \frac{u_1 - u_2}{1 - u_-} \right|^\mu} \left(\ln \frac{4}{1 - u_-} \right)^k, \quad (5.3)$$

since if this norm is finite and $\varphi(1) = 0$, it follows that $\sup |\varphi(u)|$ is finite.

Let finally $C(k)$ be the space of continuous functions on $[0,1]$ with the norm

$$\|\varphi\| = \sup_{u \in [0,1]} \left\{ |\varphi(u)| \left(\ln \frac{4}{1-u} \right)^k \right\}, \quad k \geq 0. \quad (5.4)$$

For $k = 0$ this is the usual supremum norm, for $k > 0$ it also demands φ to approach zero as $u \rightarrow 1$ at least as fast as $(\ln 4/1-u)^{-k}$.

Since Hölder continuity of the scattering amplitude follows from unitarity and analyticity, the discussion of Section 2 shows that any solution to the dispersion relation A.1 or A.2 and the unitarity condition B will have

$$\operatorname{Re} f(u) \in H_0(\mu, 1+\gamma), \quad \operatorname{Im} f(u) \in H_0(\mu, 2+\gamma), \quad (5.5)$$

with $\mu \leq \frac{1}{2}$ and γ defined by Eq. (2.9).

Let us now write the dispersion relations in the form

$$g(u) \operatorname{Re} f(u) = [(K_R + K_L) \operatorname{Im} f](u), \quad (5.6)$$

where

$$(K_R \varphi)(u) = \frac{g(u)}{\pi} \rho \int_0^1 \frac{(1-u) \varphi(u') du'}{(1-u')(u'-u)} \quad (5.7)$$

and

$$(K_L \varphi)(u) = (K_2 \varphi)(u) = \frac{g(u)}{\pi} \int_0^1 \frac{2(1-u)}{u(1-u')^2} \ln \left[1 + \frac{u}{1-u} (1-u') \right] \varphi(u') du' \quad (5.8)$$

for the case of dispersion relation A.1, and

$$(K_L \varphi)(u) = (\bar{K}_m \varphi)(u) = \frac{g(u)}{\pi} \int_0^1 \frac{m(1-u) \varphi(u') du'}{(1-u')(1-u'u)}, \quad m \geq 0, \quad (5.9)$$

for the case of dispersion relation A.2.

K_R is a singular linear integral operator from $H_0(\mu, k+1)$ to $H_0(\mu, k)$, any $k > 0$. It is bounded in the norm (5.3), as shown in Appendix 2. K_2 and $\bar{K}_m$ are non-singular linear integral operators, defining mappings of $C(k+1)$ into $H_0(\mu, k)$, which are also bounded in the norm (5.3).

K_2 and $\bar{K}_2$ have a similar behaviour near $u = 1$, so that their difference

$$\Delta K = K_2 - \bar{K}_2 \quad (5.10)$$

is more well-behaved than either of them. It is shown in Appendix 2 that ΔK defines a bounded map of $C(k)$ into $H_0(\mu, k)$.

Now $H(\mu, k)$ is a normed ring, multiplication being a well-defined operation in it. Therefore, the unitarity condition B is a non-linear relation in $H(\mu, 1+\gamma)$, provided we assume

$$\eta(u) \in H(\mu, 1+\gamma), \quad R(u) \left(\ln \frac{4}{1-u} \right)^{-\gamma} \in H(\mu, 1). \quad (5.11)$$

Introduce the phase shift $\delta(u) \in H(\mu, 1+\gamma)$, and put

$$\operatorname{Re} f(u) = \frac{\eta(u)}{2g(u)} \sin 2\delta(u), \quad \operatorname{Im} f(u) = \frac{1}{2g(u)} \{1 - \eta(u) \cos 2\delta(u)\}. \quad (5.12)$$

Assuming also $\delta(0) = 0$ and $\eta(0) = 1$, so that we get normal threshold behaviour, Eqs. (5.12) are well-defined relations in the normed ring $H(\mu, 1+\gamma)$.

Since further $\operatorname{Re} f(1) = 0$, $\operatorname{Im} f(1) = 0$, $\eta(1) = 1$ for any solution according to Section 2, we must have $\delta(1) = \pi\nu$, $\nu = 0, \pm 1, \pm 2, \dots$. In fact, we have as $u \rightarrow 1$

$$\delta(u) = \begin{cases} \pi\nu + \pi\sigma \left(\ln \frac{4}{1-u} \right)^{-1-\gamma} + O\left(\left(\ln \frac{4}{1-u} \right)^{-2+\gamma} \right) & \text{for class 1 solutions} \\ \pi\nu + O((1-u)^\mu) & \text{for class 2 solutions,} \end{cases} \quad (5.13)$$

where

$$\sigma = \frac{1+\gamma}{R_0(1+m)}. \quad (5.14)$$

With the phase shift, the dispersion relation can now be written

$$P(\delta(u)) \equiv \frac{1}{2} \eta(u) \sin 2\delta(u) - \left[(K_R + K_L) \frac{1 - \eta \cos 2\delta}{2g} \right](u) = 0. \quad (5.15)$$

This is a non-linear relation in $H(\mu, 1+\gamma)$, which is equivalent to the unitarity condition B combined with the dispersion relation A.1 or A.2. In the following, it is the solutions of this equation that we shall study.

Since $H(\mu, 1+\gamma) \subset H(\mu, 1)$ for $\gamma \geq 0$, we shall treat Eq. (5.15) as a relation in $H(\mu, 1)$, thereby including all cases.

We are interested in the local multiplicity of the solutions. Therefore, we assume $\delta(u)$ to be a solution to Eq. (5.15), of class 1 or of class 2, and make a small variation $\delta\delta(u)$ of it. Let P'_η be the Fréchet derivative of P , that is

$$P'_\eta(m, \delta) \delta\delta(u) = \eta(u) \cos 2\delta(u) \delta\delta(u) - \left[(K_R + K_L) \frac{\eta \sin 2\delta}{g} \delta\delta \right](u). \quad (5.16)$$

The index η is to indicate that η has been held constant during the differentiation. The m dependence comes from K_L of Eq. (5.9). When we occasionally use Eq. (5.8) for K_L we just leave out the argument m . The Fréchet derivative is a linear operator in $H(\mu, 1)$, or in $H_0(\mu, 1)$ if we assume $\delta\delta(1) = 0$.

In order to compare with Section 4 we also need the derivative with respect to $\delta(u)$ when $R(u)$ is kept fixed and $\eta(u)$ is allowed to vary accordingly. The unitarity condition gives the following relation between R and η :

$$\eta^2 - 2\eta \frac{R-1}{R} \cos 2\delta + \frac{R-2}{R} = 0 \quad (5.17)$$

Thus, for R constant we have the following connection between $\delta\eta$ and $\delta\delta$:

$$\delta\eta = - \frac{2\eta \frac{R-1}{R} \sin 2\delta}{\eta - \frac{R-1}{R} \cos 2\delta} \delta\delta \quad (5.18)$$

This gives for the derivative with constant R the expression

$$P'(m, \delta) \delta\delta(u) = \frac{\eta(u)}{s(u)} \frac{\eta(u) \cos 2\delta(u) - \frac{R(u)-1}{R(u)}}{\eta(u) - \frac{R(u)-1}{R(u)} \cos 2\delta(u)} \delta\delta(u) - \left[(K_R + K_L) \frac{\eta^2 \sin 2\delta}{\eta - \frac{R-1}{R} \cos 2\delta} \frac{\delta\delta}{s} \right] (u). \quad (5.19)$$

The operator defined by this expression is also a linear operator in $H_0(\mu, 1)$, provided $\delta\delta \in H_0(\mu, 1)$ and

$$\eta(u) \neq \frac{R(u)-1}{R(u)} \cos 2\delta(u), \quad 0 < u < 1. \quad (5.20)$$

This condition corresponds to the absence of a double root of Eq. (5.17).

From Eq. (1.6) we see that such a double root can occur if $R(s)$ is allowed to become ≥ 2 , and for some s

$$|\sin 2\delta(s)| = \frac{1}{R(s)-1}. \quad (5.21)$$

In fact, as soon as $R(s) > 2$, the variation of the phase shift is restricted since Eq. (1.6) then implies

$$|\sin 2\delta(s)| \leq \frac{1}{R(s)-1}.$$

This condition forces the phase shift to stay in the interval S_j ,

$$S_j = (-\delta_{\max} + j\pi, \delta_{\max} + j\pi), \quad \delta_{\max} = \frac{1}{2} \arcsin \frac{1}{R(s)-1}, \quad j=0, \pm 1, \dots \quad (5.22)$$

As long as $R(s)$ stays > 2 , $\delta(s)$ has to remain in that interval S_j , where it was when R became > 2 .

The condition (5.20) corresponds to δ not touching the borders of the allowed region. When it does, it is quite natural that making an arbitrary variation $\delta\delta$ causes a singularity in Eqs. (5.18) and (5.19).

In the following, when R is given and ≥ 2 , we shall invariably assume the condition (5.20) satisfied.

Provided we can show that $P'(m, \delta)$ is a $\emptyset$ operator, i.e., that it is closed, has a closed range, and that its null space and defect space are finite dimensional, we can follow Lovelace ⁵⁾ and employ the "implicit variable" theorem of Vainberg and Trenogin ⁶⁾. Putting α and β equal to the dimensions of the null space and the defect space, respectively, of $P'(m, \delta)$, the implicit variable theorem says that in the neighbourhood of the solution $\delta(u)$, the equation (5.14) is exactly equivalent to β non-linear scalar equation in α real variables.

The index $\kappa = \alpha - \beta$ of the $\emptyset$ operator $P'(m, \delta)$ is thus the excess of the number of variables over the number of equations, or, in general, the dimensionality of the manifold of solutions to Eq. (5.14) containing $\delta(u)$. Thus, $\delta(u)$ should contain κ arbitrary parameters.

We now set out to prove that $P'(m, \delta)$ is a $\emptyset$ operator, and to determine its index κ . For a smaller Banach space, containing only the class 2 solutions, this has already been done by Lovelace ⁵⁾. We shall try to extend his proof to include all solutions. Because of the very slow decrease of the class 1 solutions this is a very delicate task, as we shall see.

We decompose P' and P'_η into a sum of a right-hand cut contribution

$$P'_R(\delta) = P'(0, \delta) \quad (5.23)$$

and a left-hand cut contribution

$$P'_L(m, \delta) = P'(m, \delta) - P'(0, \delta). \quad (5.24)$$

The right-hand cut operator is examined in the next section and in Section 7 the influence of the left-hand cut contribution is analyzed.

6. - THE INDEX OF THE RIGHT HAND CUT FRÉCHET DERIVATIVE

Combining the result of Section 4 with the implicit variable theorem immediately tells us that for elastic unitarity and $m = 0$, that is for the right-hand cut Fréchet derivative $P'_R(\delta)$, the index is κ , where

$$\kappa = \begin{cases} 2\nu + 1 & \text{if } \delta \text{ is a class 1 solution with } m = 0 \\ 2\nu & \text{if } \delta \text{ is a class 2 solution with } m = 0 \end{cases} \quad (6.1)$$

However, in order to be allowed to use the implicit variable theorem, we should first prove that $P'_R(\delta)$ is a $\emptyset$ operator. We would also like to cover inelastic unitarity, and study $P'_R(\delta)$ when δ is a solution with $m > 0$. Therefore, we shall now solve explicitly the equation

$$P'_R(\delta) \delta \delta(u) = \delta f(u), \quad (6.2)$$

that is the singular integral equation

$$A(u) \delta \delta(u) - \frac{1}{i\pi} P \int_0^1 \frac{(1-u)B(u')}{(1-u')(u'-u)} \delta \delta(u') du' = \frac{\delta f(u)}{\delta(u)}. \quad (6.3)$$

Here, A and B are the functions

$$\begin{cases} A(u) = \frac{\eta(u)}{\delta(u)} \frac{\eta(u) \cos 2\delta(u) - \frac{R(u)-1}{R(u)}}{\eta(u) - \frac{R(u)-1}{R(u)} \cos 2\delta(u)} \\ B(u) = i \frac{\eta^2(u)}{\delta(u)} \frac{\sin 2\delta(u)}{\eta(u) - \frac{R(u)-1}{R(u)} \cos 2\delta(u)} \end{cases} \quad (6.4)$$

which are Hölder continuous on $(0,1)$, excluding the end points, provided the condition (5.20) is satisfied. But then Eq. (6.3) is of the type studied in Muskhelishvili's book ¹⁰⁾, except that for class 1 solutions we must relax the Hölder condition at $u = 1$. The function ϕ ,

$$\phi(z) = \frac{1}{2\pi i} \int_0^1 \frac{(1-z)B(u') \delta \delta(u') du'}{(1-u')(u'-z)}, \quad (6.5)$$

is analytic, except for a cut $(0,1)$. On the cut its limiting values $\phi^\pm(u)$, from above and below, respectively, belong to $H_0(\mu,1)$ if $\delta\delta \in H_0(\mu,1)$. This follows from the proof of the boundedness of K_R in Appendix 2 and the fact that $B(0) = 0$ and

$$B(u) \underset{u \rightarrow 1}{\sim} i\pi\sigma R_0 \left(\ln \frac{4}{1-u}\right)^{-1} \quad (6.6)$$

when δ is a class 1 solution with

$$\delta(u) \underset{u \rightarrow 1}{\sim} \pi\nu + \pi\sigma \left(\ln \frac{4}{1-u}\right)^{-1-\gamma}. \quad (6.7)$$

Equation (6.2') can now be written

$$A(u)\delta\delta(u) - \phi^+(u) - \phi^-(u) = \frac{\delta f(u)}{g(u)}, \quad (6.8)$$

which, since $\phi^+ - \phi^- = B\delta\delta$, is equivalent to the Hilbert problem

$$\phi^+(u) = \frac{A(u)+B(u)}{A(u)-B(u)} \phi^-(u) + \frac{B(u)}{A(u)-B(u)} \frac{\delta f(u)}{g(u)}. \quad (6.9)$$

Here

$$\frac{A(u)+B(u)}{A(u)-B(u)} = e^{4i\delta(u)} \frac{1 - \frac{R-1}{\gamma R} e^{-2i\delta}}{1 - \frac{R-1}{\gamma R} e^{2i\delta}}, \quad (6.10)$$

which is finite and non-zero on $0 \leq u \leq 1$. Taking the logarithm thus gives a finite function, $\Delta(u)$, which near $u = 1$ can be shown to have the asymptotic form

$$\Delta(u) = \frac{1}{2i} \ln \frac{A+B}{A-B} = 2\pi\nu + 2\pi\sigma R_0 \left(\ln \frac{4}{1-u}\right)^{-1} + O\left(\left(\ln \frac{4}{1-u}\right)^{-2}\right) \quad (6.11)$$

for any class 1 solution with the asymptotic form (6.7).

For class 2 solutions one gets instead

$$\Delta(u) = 2\pi\nu + O((1-u)^\mu), \quad (6.11')$$

that is, a Hölder condition also at $u = 1$.

Introduce now the function

$$\Gamma(z) = \frac{1}{\pi} \int_0^1 \frac{\Delta(u') du'}{u' - z}, \quad (6.12)$$

which is analytic except for a cut $(0,1)$, on which $\Gamma^\pm(u)$ is Hölder continuous, excluding $u = 1$, where

$$\Gamma^\pm(u) = \begin{cases} 2\nu \ln(1-u) - 2\sigma R_0 \ln\left(\ln \frac{4}{1-u}\right) + O(1) & \text{for } \delta \text{ class 1} \\ 2\nu \ln(1-u) + O(1) & \text{for } \delta \text{ class 2.} \end{cases} \quad (6.13)$$

This follows as for $\phi^\pm$ above, using for the asymptotic form of the integral (6.12) also the results of Appendix 1.

The function $\exp(\Gamma)$ is now a solution to the homogeneous Hilbert problem, and a solution to the inhomogeneous problem (6.8) can be constructed as in Ref. 9). However, we shall follow Lovelace, and use the Dashen-Frautschi method ¹¹⁾. The D function is, since we have assumed no bound states, given by

$$D(u) = e^{-\frac{1}{2}\Gamma^+(u)}. \quad (6.14)$$

It has the asymptotic form

$$D(u) \underset{u \rightarrow 1}{\sim} \begin{cases} \text{const } (1-u)^{-\nu} \left(\ln \frac{4}{1-u}\right)^{\sigma R_0} & \text{for } \delta \text{ class 1} \\ \text{const } (1-u)^{-\nu} & \text{for } \delta \text{ class 2,} \end{cases} \quad (6.15)$$

or in the s variable

$$D(s) \underset{s \rightarrow \infty}{\sim} \begin{cases} \text{const } s^\nu (\ln s)^{\sigma R_0} & \text{for } \delta \text{ class 1} \\ \text{const } s^\nu & \text{for } \delta \text{ class 2,} \end{cases} \quad (6.15')$$

where the constants ν and σ are defined by Eqs. (6.7) or (5.13) and (5.14).

According to the Dashen-Frautschi method, we next write a dispersion relation for the function $D^2\phi$, which is analytic, except for a cut $(0,1)$ in u or $(4,\infty)$ in s . On the cut

$$\text{Im}\{D^2(s)\phi^+(s)\} = |D(s)|^2 \sin \Delta(s) \frac{\delta f(s)}{g(s)}, \quad (6.16)$$

as follows from Eqs. (6.8), (6.12) and (6.14). The asymptotic form of this is

$$\text{Im}\{D^2(s)\phi^+(s)\} \underset{s \rightarrow \infty}{\sim} \begin{cases} \text{const } s^{2\nu} (\ln s)^{-2+2\sigma R_0} & \text{for } \delta \text{ class 1} \\ \text{const } s^{2\nu-\mu} & \text{for } \delta \text{ class 2,} \end{cases} \quad (6.17)$$

provided we assume the perturbation to go to zero as $s \rightarrow \infty$, that is $\delta f \in H_0(\mu, 1)$.

If $2\sigma R_0 \geq 1$ we obviously have to make $2\nu+1$ subtractions in the dispersion relation for $D^2\phi$, otherwise it is sufficient with 2ν subtractions. Since the non-linear problem of Section 4 had no solutions with $\nu < 0$, we assume here that $\nu \geq 0$. Putting

$$x = \begin{cases} 2\nu+1 & \text{for class 1 and } 2\sigma R_0 \geq 1, \\ 2\nu & \text{otherwise,} \end{cases} \quad (6.18)$$

the dispersion relation for $D^2\phi$ can be written

$$D^2(s)\phi(s) = Q_{x-1}(s) + \frac{s^x}{\pi} \int_4^\infty \frac{|D(s')|^2 \sin \Delta(s') \delta f(s') ds'}{s'^x (s'-s) 2g(s')}, \quad (6.19)$$

where $Q_{x-1}(s)$ is an arbitrary polynomial of degree $\leq x-1$. This is obviously the most general analytic function fulfilling Eqs. (6.16), (6.17) and the bound

$$|D^2(s)\phi(s)| \underset{|s| \rightarrow \infty}{\leq} \begin{cases} \text{const } |s|^{2\nu} (\ln |s|)^{-1+2\sigma R_0} & \text{for } \delta \text{ class 1} \\ \text{const } |s|^{2\nu} (\ln |s|)^{-1} & \text{" } \delta \text{ " 2,} \end{cases} \quad (6.20)$$

which follows from Eq. (6.15') and the fact that $\phi^\pm \in H_0(\mu, 1)$.

The expression for $\delta\delta$ corresponding to Eq. (6.19) can now be obtained with the help of Eq. (6.8) :

$$\delta\delta(s) = \frac{\delta f(s)}{\eta(s)} \cos 2\delta(s) + \frac{2g(s)}{\eta(s)|D(s)|^2} \left[Q_{\kappa-1}(s) + \frac{s^\kappa}{\pi} P \int_4^\infty \frac{|D(s')|^2 \sin \Delta(s') \delta f(s') ds'}{s'^\kappa (s'-s) 2g(s')} \right] \quad (6.21)$$

This is the explicit solution of Eq. (6.3). It contains exactly κ arbitrary parameters, the coefficients of the polynomial $Q_{\kappa-1}(s)$. Note that

$$\frac{2g(s)}{\eta(s)|D(s)|^2} Q_{\kappa-1}(s) \underset{s \rightarrow \infty}{\sim} \begin{cases} \text{const } s^{\kappa-2\nu-1} (\ln s)^{-2\sigma R_0} & \text{for } \delta \text{ class 1} \\ \text{const } s^{\kappa-2\nu-1} & \text{for } \delta \text{ class 2,} \end{cases} \quad (6.22)$$

so that if we, e.g., had chosen $\kappa = 2\nu + 1$ for $2\sigma R_0 < 1$, $\delta\delta$ would not have belonged to $H_0(\mu, 1)$. Therefore, we are forced to choose κ according to Eq. (6.18).

Using the asymptotic form (6.11) for Δ it is easy to see (cf. Appendix 2) that the last term of Eq. (6.21), like the first, gives contributions to $\delta\delta \in H_0(\mu, 1)$ for any $\delta f \in H_0(\mu, 1)$. [Provided $\eta(s) \neq 0$, which we assume.] Thus the defect space of $P'_R(\delta)$ is empty, and its range is the whole Banach space $H_0(\mu, 1)$ thus closed. Since $P'_R(\delta)$ is bounded, it is a closed operator. Since further, according to Eq. (6.21), its null space has dimension κ , we have now proved $P'_R(\delta)$ to be a $\emptyset$ operator with index κ given by Eq. (6.18).

Let $\mathcal{N}_\kappa$ be the null space of $P'_R(\delta)$, and define the factor space

$$\mathcal{B} = H_0(\mu, 1) / \mathcal{N}_\kappa. \quad (6.23)$$

Let further G_R be the restriction of $P'_R(\delta)$ to $\mathcal{B}$. Then G_R defines a 1 to 1 mapping of $\mathcal{B}$ onto $H_0(\mu, 1)$, and its inverse G_R^{-1} exists. From Eq. (6.21) we obtain

$$G_R^{-1}(\delta) \delta f(s) = \frac{\delta f(s)}{\eta(s)} \cos 2\delta(s) + \frac{g(s)s^\kappa}{\pi \eta(s)|D(s)|^2} P \int_4^\infty \frac{|D(s')|^2 \sin \Delta(s') \delta f(s') ds'}{s'^\kappa (s'-s) g(s')}. \quad (6.24)$$

This is a bounded operator from $H_0(\mu, 1)$ to $\mathcal{B}$, [provided $\eta(s) \neq 0$] as proved in Appendix 2. G_R and G_R^{-1} are $\emptyset$ operators with index zero.

The Fréchet derivative of $P(\delta)$ with respect to δ with η fixed is simpler than the one we have studied above, with R fixed. For the case with η fixed we get simply

$$\Delta(u) = 2\delta(u) \sim_{u \rightarrow 1} 2\nu\pi + 2\pi\sigma \left(\ln \frac{4}{1-u}\right)^{-1-\delta} \quad (6.25)$$

which results in

$$D(s) \sim_{s \rightarrow \infty} \begin{cases} \text{const } s^\nu (\ln s)^\sigma & \text{for } \delta \text{ class 1 and } \gamma = 0 \\ \text{const } s^\nu & \text{otherwise.} \end{cases} \quad (6.26)$$

This gives with the same arguments as above that the index of $P'_{\eta, R}(\delta)$ is

$$\kappa = \begin{cases} 2\nu + 1 & \text{for } \delta \text{ class 1, } \gamma = 0 \text{ and } 2\sigma \geq 1, \\ 2\nu & \text{otherwise.} \end{cases} \quad (6.27)$$

For class 1 solutions, the index thus may depend on how we formulate the problem, with η or R kept fixed.

We shall finally take advantage of the fact that for $m = 1$ the left-hand cut discontinuity is just the mirror image in $s = 2$ of the one of the right-hand cut. This means that in the variable

$$t = (s-2)^2, \quad (6.28)$$

the amplitude is a real analytic function with just a right-hand cut $(4, \infty)$. As a result of this, we can solve the equation

$$P'(1, \delta) \delta \delta(u) = \delta f(u) \quad (6.29)$$

with the method used above for $P'_R(\delta)$.

In the variable t we have

$$P'(1, \delta) \delta \delta(t) = g(t) A(t) \delta \delta(t) - \frac{g(t)}{\pi i} P \int_4^\infty \frac{B(t') \delta \delta(t') dt'}{t' - t}, \quad (6.30)$$

where A and B are given in Eq. (6.4) as functions of

$$u = 1 - \frac{4}{s} = 1 - \frac{4}{2 + \sqrt{t}}. \quad (6.31)$$

From Eq. (6.6) we then get the asymptotic form

$$B(t) \underset{t \rightarrow \infty}{\sim} 2i\pi\sigma R_0 (\ln t)^{-1} \quad (6.32)$$

for any class 1 solution. The factor 2 here comes from the fact that $s \sim \sqrt{t}$. It comes in also in the asymptotic form of $\Delta(t)$:

$$\Delta(t) \underset{t \rightarrow \infty}{\sim} \begin{cases} 2\pi\nu + 4\pi\sigma R_0 (\ln t)^{-1} & \text{for } \delta \text{ class 1} \\ 2\pi\nu & \text{for } \delta \text{ class 2.} \end{cases} \quad (6.33)$$

In the asymptotic form of the D function, defined as above, this extra 2 also enters :

$$D(t) \underset{t \rightarrow \infty}{\sim} \begin{cases} \text{const } t^\nu (\ln t)^{2\sigma R_0} & \text{for } \delta \text{ class 1} \\ \text{const } t^\nu & \text{for } \delta \text{ class 2.} \end{cases} \quad (6.34)$$

But this means that the rule for the number of subtractions changes compared to Eq. (6.18) :

$$n = \begin{cases} 2\nu + 1 & \text{for } \delta \text{ class 1 and } 4\sigma R_0 \geq 1 \\ 2\nu & \text{otherwise.} \end{cases} \quad (6.35)$$

Let now $\mathcal{N}_\infty$ be the null space of $P'(1, \delta)$, and $G(1, \delta)$ be the restriction of $P'(1, \delta)$ to the factor space (6.23). Then the inverse G^{-1} is a bounded operator from $H_0(\mu, 1)$ to $\mathcal{B}$ and has the explicit form

$$G^{-1}(1, \delta) \delta f(t) = \frac{\delta f(t)}{\eta(t)} \cos 2\delta(t) + \frac{g(t) t^\nu}{\pi \eta(t) |D(t)|^2} P \int_4^\infty \frac{|D(t')|^2 \sin \Delta(t') \delta f(t') dt'}{t'^2 (t' - t) g(t')}. \quad (6.36)$$

As a last point of this section we point out the differences in the last derivation for the case that the Fréchet derivative is taken with η fixed.

For Δ we would then have obtained

$$\Delta(t) \underset{t \rightarrow \infty}{\sim} \begin{cases} 2\pi\nu + 4\pi\sigma (\ln t)^{-1-\sigma} & \text{for } \delta \text{ class 1} \\ 2\pi\nu & \text{for } \delta \text{ class 2,} \end{cases} \quad (6.37)$$

which results in the following number of subtractions :

$$\kappa = \begin{cases} 2\nu+1 & \text{for } \delta \text{ class 1, } \sigma = 0 \text{ and } 4\sigma \geq 1 \\ 2\nu & \text{otherwise.} \end{cases} \quad (6.38)$$

Also for this case we can define the restriction G_η of P'_η to the factor space $\mathcal{B}$. This operator has the inverse

$$G_\eta^{-1}(1, \delta) \delta f(t) = \frac{\delta f(t)}{\eta(t)} \cos 2\delta(t) + \frac{g(t) t^\kappa}{\pi \eta(t) |D(t)|^2} P \int_4^\infty \frac{|D(t')|^2 \sin 2\delta(t') \delta f(t') dt'}{t'^\kappa (t'-t) g(t')}, \quad (6.39)$$

which is a bounded operator from $H_0(\mu, 1)$ to $\mathcal{B}$. This proves that G_η and G_η^{-1} are $\emptyset$ operators with index zero, and $P'_\eta(1, \delta)$ is a $\emptyset$ operator with index κ given by Eq. (6.38) [see Ref. 7), p. 193 and Lemma 2.2].

7. - THE LEFT-HAND CUT CONTRIBUTION

In this section we shall discuss whether or not the addition of the left-hand cut operator $P'_L(m, \delta)$ to the $\emptyset$ operator $P'_R(\delta)$ may change its index. The tools we have at disposal for this task are some theorems on $\emptyset$ operators from Ref. 7).

First we shall employ Theorem 2.3 of Ref. 7) [= Theorem 5A of Lovelace ⁵⁾], which says that if $P'_L(m, \delta)$ is a compact operator, the sum

$$P'(m, \delta) = P'_R(\delta) + P'_L(m, \delta) \quad (7.1)$$

is a $\emptyset$ operator with the same index as $P'_R(\delta)$.

We recall from Section 5 that we had two different forms of K_L , Eqs. (5.8) and (5.9), respectively, and that their difference ΔK , defined in Eq. (5.10), is asymptotically smaller than either of them, so that ΔK defines a bounded map from $C(k)$ to $H_0(\mu, k)$ (cf. Appendix 2).

For the corresponding Fréchet derivatives P'_L we correspondingly define the difference

$$\Delta P'_\eta(\delta)\delta\delta(u) = [P'_\eta(\delta) - P'_\eta(R, \delta)]\delta\delta(u) = -\left(\Delta K \frac{\eta^2 \sin 2\delta}{\delta}\delta\delta\right)(u) \quad (7.2)$$

for the derivative with constant η , and

$$\Delta P'_R(\delta)\delta\delta(u) = -\left(\Delta K \frac{\eta^2 \sin 2\delta}{\delta(\eta - \frac{R-1}{R}\cos 2\delta)}\delta\delta\right)(u) \quad (7.2')$$

for the derivative with constant R .

Provided the condition (6.4) is satisfied in the latter case, both these operators are compact operators in $H_0(\mu, 1)$, as is seen from the following decomposition into a series of successive mappings :

- (i) a compact map from $H_0(\mu, 1)$ to $C(0)$: this embedding is compact by the Arzelà-Ascoli Theorem (see Appendix 2) ;
- (ii) a bounded map from $C(0)$ to $C(1)$: multiplication by

$$\frac{\eta \sin 2\delta}{\delta} \quad \text{or} \quad \frac{\eta^2 \sin 2\delta}{\delta(\eta - \frac{R-1}{R}\cos 2\delta)},$$

respectively ;

- (iii) a bounded map from $C(1)$ to $H_0(\mu, 1)$: the action of the operator ΔK .

Since the product of a compact operator and a bounded operator is compact [Ref. 12), VI.5.4], it follows that the entire mapping is compact.

That the operators (7.2) are compact implies by the above theorem that the Fréchet derivative P' has the same index for the case of dispersion relation A.1 as for dispersion relation A.2 with $m = 2$, and that if P' is a $\emptyset$ operator for one of those cases it is so for the other also. Therefore, we can in the following concentrate on the case of dispersion relation A.2.

Next, we discuss the case of fixed η , that is the operator $P'_{\eta,L}$ with the explicit form

$$P'_{\eta,L}(m, \delta) \delta \delta(u) = - \left(K_L \frac{\eta \sin 2\delta}{g} \delta \delta \right)(u) = -m \frac{g(u)}{\pi} \int_0^1 \frac{(1-u') \eta(u') \sin 2\delta(u')}{(1-u')(1-u'u) g(u')} \delta \delta(u') du'. \quad (7.3)$$

If δ is a class 2 solution, or class 1 solution when $\gamma > 0$, we can follow Lovelace and decompose $P'_{\eta,L}$ as above for $\Delta P'$. We write

$$P'_{\eta,L}(m, \delta) \delta \delta(u) = - \left[K_L \left(\ln \frac{4}{1-u'} \right)^{\gamma-1} \frac{\eta(u') \sin 2\delta(u')}{g(u')} \left(\ln \frac{4}{1-u'} \right)^{1-\gamma} \delta \delta(u') \right](u). \quad (7.4)$$

The successive mappings are :

- (i) a bounded map from $H_0(\mu, 1)$ to $H_0(\mu, \gamma)$: multiplication with $(\ln(4/1-u'))^{1-\gamma}$;
- (ii) a compact map from $H_0(\mu, \gamma)$ to $C(0)$: this embedding is compact by the Arzelà-Ascoli theorem, provided $\gamma > 0$;
- (iii) a bounded map from $C(0)$ to $C(2)$: multiplication with [cf. Eqs. (5.5) and (5.12)]

$$\left(\ln \frac{4}{1-u'} \right)^{\gamma-1} \frac{\eta(u') \sin 2\delta(u')}{g(u')}.$$

- (iv) a bounded map from $C(2)$ to $H_0(\mu, 1)$: the action of the operator K_L .

The net result is thus a compact map of $H_0(\mu, 1)$ into itself, proving that for $\gamma > 0$ $P'_{\eta,L}(m, \delta)$ is a compact operator in $H_0(\mu, 1)$.

For $\gamma = 0$ and δ a class 2 solution, the same result follows by simply putting $\gamma = \varepsilon > 0$ in Eq. (7.4) and repeating the argument. However, if δ is a class 1 solution, this trick would not work, since the map (iii) would not be bounded.

Employing again Theorem 2.3 of Ref. 7), we now have that for δ a class 2 solution, or a class 1 solution for $\gamma > 0$, $P'_{\eta}(m, \delta)$ is a $\emptyset$ operator with the index fixed by Eq. (6.27) to $\kappa = 2\nu$.

For δ a class 1 solution and $\gamma = 0$, however, we cannot prove the compactness of $P'_{\eta, L}$. If the limit $\gamma \rightarrow 0$ is taken in Eq. (5.16) we formally get

$$P'_{\eta}(m, \gamma, \delta) \xrightarrow{\gamma \rightarrow 0} P'_{\eta}(m, 0, \delta), \quad (7.5)$$

but since this limit in Eq. (5.16) is not approached uniformly in u , Eq. (7.5) must be understood in a weak sense, not in the uniform operator topology. [If the limit had been uniform, $P'_{\eta, L}$ would have been compact by Ref. 12), VI.5.3.]

Instead of using Theorem 2.3, we shall now use the limit (7.5) to prove that P'_{η} is a $\emptyset$ operator also for $\gamma = 0$. Because of the non-uniformness of this limit, the proof becomes rather long.

We define operators G_{η}^{-1} by the relations

$$G_{\eta}^{-1}(0, \gamma, \delta) \delta f(s) = \frac{\delta f(s)}{\eta(s)} \cos 2\delta(s) + \frac{g(s) \cdot s^{2\nu+1}}{\pi \eta(s) |D(s)|^2} P \int_0^{\infty} \frac{|D(s')|^2 \sin 2\delta(s') \delta f(s') ds'}{s^{2\nu+1} (s'-s) g(s')} \quad (7.6)$$

[cf. Eq. (6.24)], where we have made an extra subtraction compared to what is necessary for $\gamma > 0$ when α is given by Eq. (6.27), and

$$G_{\eta}^{-1}(m, \gamma, \delta) = [1 + G_{\eta}^{-1}(0, \gamma, \delta) P'_{\eta, L}(m, \gamma, \delta)]^{-1} G_{\eta}^{-1}(0, \gamma, \delta). \quad (7.7)$$

For $\gamma > 0$ these operators are well defined on the whole of $H_0(\mu, 1)$, except for certain isolated m values, corresponding to spontaneous symmetry breaking⁵⁾. These m values must be isolated since they correspond to the compact operator

$$G_{\eta}^{-1}(0, \gamma, \delta) P'_{\eta, L}(m, \gamma, \delta) \quad (7.8)$$

having one of its discrete eigenvalues equal to -1 , so that the operator in square brackets in Eq. (7.7) ceases to have a bounded inverse. The extra subtraction in Eq. (7.6) is made because then the operator becomes uniformly bounded for all γ , so that the formal limit $\gamma \rightarrow 0$ can be taken. Just as for Eq. (7.5) this limit is not uniform, but to be understood in a weak sense.

That $P'_\eta(m, 0, \delta)$ is bounded, and thus closed, we know already from Section 5. To prove that it has a closed range, we prove that any convergent set of elements δf_k in the range $\mathcal{R}$,

$$\delta f_k = P'_\eta(m, 0, \delta) \delta \delta_k \in \mathcal{R}, \quad \delta f_k \xrightarrow{k \rightarrow \infty} \delta f, \quad (7.9)$$

converges to an element in the range, $\delta f \in \mathcal{R}$.

Choose an m value such that

$$G_n^{-1} \equiv G'_\eta(m, \frac{1}{n}, \delta), \quad n = 1, 2, 3, \dots, \quad (7.10)$$

are well defined by Eq. (7.7). Let further

$$\delta \delta^n = G_n^{-1} \delta f_k, \quad \delta \delta^n = G_n^{-1} \delta f \quad (7.11)$$

Since G_n^{-1} is defined on the whole of $H_0(\mu, 1)$, and is uniformly bounded¹³⁾ all the $\delta \delta$ are uniformly bounded elements of $H_0(\mu, 1)$. As $n \rightarrow \infty$ they form Cauchy sequences, since, e.g.,

$$\begin{aligned} \|\delta \delta^{n_1} - \delta \delta^{n_2}\| &= \left\| \frac{g(s)}{\pi \eta(s)} P \int_0^\infty \frac{s^{2\nu+1} \delta f(s') ds'}{s^{2\nu+1} (s'-s) g(s')} \left\{ \left[\frac{|D(s')|^2}{|D(s)|^2} \sin 2\delta(s') \right]_{\gamma=\frac{1}{n_1}} - \left[\frac{|D(s')|^2}{|D(s)|^2} \sin 2\delta(s') \right]_{\gamma=\frac{1}{n_2}} \right\} \right\| \leq \\ &\leq \text{const } \|\delta f\| \left\| [\sin 2\delta]_{\gamma=\frac{1}{n_1}} - [\sin 2\delta]_{\gamma=\frac{1}{n_2}} \right\| \xrightarrow{n_1, n_2 \rightarrow \infty} 0. \end{aligned} \quad (7.12)$$

The last norm goes to zero because the two functions approach each other pointwise, and are bounded by

$$\text{const } (\ln s)^{-1 - \min(\frac{1}{n_1}, \frac{1}{n_2})}, \quad (7.13)$$

so the norm (5.3) with $k = 1$ goes to zero.

Since $H_0(\mu, 1)$ is a complete space, it follows that

$$\delta\delta^n \xrightarrow{n \rightarrow \infty} \delta\delta \in H_0(\mu, 1) \text{ and } \delta\delta_h^n \xrightarrow{n \rightarrow \infty} \delta\delta_h^0 \in H_0(\mu, 1) \quad (7.14)$$

in the norm of $H_0(\mu, 1)$.

Next, consider

$$\delta f_h^n = P'_\eta(m, 0, \delta) \delta\delta_h^n \text{ and } \delta f^n = P'_\eta(m, 0, \delta) \delta\delta^n, \quad (7.15)$$

which are uniformly bounded elements of $H_0(\mu, 1)$. Since $\delta f_k \rightarrow \delta f$, we have

$$\delta f_h^n \xrightarrow{n \rightarrow \infty} \delta f^n \quad (7.16)$$

in the norm of $H_0(\mu, 1)$. Moreover,

$$\|\delta f_h^{n_1} - \delta f_h^{n_2}\| \leq \|P'_\eta(m, 0, \delta)\| \|\delta\delta_h^{n_1} - \delta\delta_h^{n_2}\| \xrightarrow{n_1, n_2 \rightarrow \infty} 0, \quad (7.17)$$

in analogy with Eq. (7.12), and thus

$$\delta f_h^n \xrightarrow{n \rightarrow \infty} \delta f_h^0 \in H_0(\mu, 1) \quad (7.18)$$

in the norm of $H_0(\mu, 1)$, and similarly

$$\delta f^n \xrightarrow{n \rightarrow \infty} \delta f^0 \in H_0(\mu, 1). \quad (7.19)$$

The latter limit function can be written

$$\delta f^0 = P'_\eta(m, 0, \delta) \left[\lim_{n \rightarrow \infty} \delta \delta^n \right] = P'_\eta(m, 0, \delta) \delta \delta \quad (7.20)$$

with the help of Eq. (7.14), since the limits involved are in the norm.

Finally we note that

$$P'_\eta(m, \frac{1}{n}, \delta) \delta \delta_h^n = \delta f_h \quad (7.21)$$

since according to Eq. (7.7)

$$P'_\eta(m, \frac{1}{n}, \delta) G_\eta^{-1}(m, \frac{1}{n}, \delta) \varphi = \varphi \quad (7.22)$$

for any φ in the range of $P'_\eta(m, \frac{1}{n}, \delta)$, that is for any $\varphi \in H_0(\mu, 1)$.
It follows that

$$|\delta f_h(u) - \delta f_h^0(u)| \leq |P'_\eta(m, \frac{1}{n}, \delta) (\delta \delta_h^n - \delta \delta_h^0)| + |P'_\eta(m, \frac{1}{n}, \delta) \delta \delta_h^0 - P'_\eta(m, 0, \delta) \delta \delta_h^0|. \quad (7.23)$$

Here, the first term on the right-hand side goes to zero as $n \rightarrow \infty$ because of Eq. (7.14), and the second because the limit (7.5) is approached pointwise. Thus

$$\delta f_h^0(u) = \delta f_h(u) \text{ pointwise,} \quad (7.24)$$

but since they are both elements of $H_0(\mu, 1)$, they must then be identical,

$$\|\delta f_h - \delta f_h^0\| = 0. \quad (7.25)$$

Putting now all the pieces together, we have that

$$\|\delta f - \delta f^0\| \leq \|\delta f - \delta f_h\| + \|\delta f_h - \delta f_h^0\| + \|\delta f_h^0 - \delta f_h^{\sim}\| + \|\delta f_h^{\sim} - \delta f^{\sim}\| + \|\delta f^{\sim} - \delta f^0\|. \quad (7.26)$$

By Eqs. (7.9), (7.25), (7.18), (7.16), (7.19) and (7.20) this implies that

$$\delta f = \delta f^0 = P_\eta'(m, 0, \delta) \delta f, \quad (7.27)$$

which shows that δf is in the range of $P_\eta'(m, 0, \delta)$, that is $\delta f \in \mathcal{R}$, which is what we wanted to show. It moreover follows that $\mathcal{R} = H_0(\mu, 1)$.

We do not consider a serious limitation the condition that m should be such that the operators (7.10) are well defined, since either one could choose another set of G_n^{-1} , or one could work with a range $\mathcal{R}_\gamma \neq H_0(\mu, 1)$ for $P_\eta'(m, \gamma, \delta)$. In this way it is probably possible to show that the range $\mathcal{R}$ is closed for all m . However, we do not want to go into the details also of this point here. To check what happens for a particular m value one would anyway have to make detailed numerical calculations. Even for compact left-hand cut contribution such calculations are needed to say whether symmetry breaking occurs or not.

From the above proof, it also follows that the null space of $P_\eta'(m, 0, \delta)$ is at most $2\nu + 1$ dimensional. This concludes the proof that $P_\eta'(m, 0, \delta)$ is a $\emptyset$ operator (for almost all m).

What is the index of this $\emptyset$ operator? From Section 6 we have that if

$$\delta(s) = \pi\nu + \pi\sigma (\ln s)^{-1} + O((\ln s)^{-2}) \quad (7.28)$$

then the index of

$$P_\eta'(0, 0, \delta) \text{ is } \kappa = \begin{cases} 2\nu + 1 & \text{if } 2\sigma \geq 1 \\ 2\nu & \text{if } 2\sigma < 1 \end{cases} \quad (7.29)$$

and

$$P'_\eta(1,0,\delta) \text{ is } \kappa = \begin{cases} 2\nu+1 & \text{if } 4\sigma \geq 1 \\ 2\nu & \text{if } 4\sigma < 1. \end{cases} \quad (7.30)$$

From Section 2 we further know that in order for δ to be a solution to the non-linear problem, we must couple σ and m . But then the conditions of Eqs. (7.29) and (7.30) on κ become

$$\sigma = \frac{1+\delta}{R_0(1+m)} \Rightarrow \kappa = \begin{cases} 2\nu+1 & \text{if } R_0 \leq 2 \\ 2\nu & \text{if } R_0 > 2. \end{cases} \quad (7.31)$$

We shall now prove that this rule for κ is valid not only for $m = 0$ and $m = 1$, but for (almost) all m . We can see the mechanism: if m is increased, the phase shift δ decreases faster as $s \rightarrow \infty$, tending to decrease the index of $P'_\eta(0,0,\delta)$, but adding the (non-compact!) left-hand cut contribution to get $P'_\eta(m,0,\delta)$ tends to increase the index, so that the final rule for the index, Eq. (7.31), does not contain m !

Gokhberg and Krein ⁷⁾ have some theorems on the stability of the index, which we shall use. Theorem 2.2 tells us that adding a bounded operator with sufficiently small norm to a $\emptyset$ operator does not change its index. Thus there is a neighbourhood of $m = 0$ and $m = 1$, where the index is given by Eq. (7.31).

Theorem 3.5 says that if $P'_\eta(m,0,\delta)$ is a holomorphic operator function for m in a certain region Ω , and is a $\emptyset$ operator for each $m \in \Omega$, then the index is constant throughout the region Ω .

Take the non-linear equation (5.15) :

$$P(m, \delta(u)) = \frac{1}{2} \eta(u) \sin 2\delta(u) - \left[(K_R + m \bar{K}_1) \frac{1 - \eta \cos 2\delta}{2\delta} \right](u) = 0 \quad (7.32)$$

The only m dependence of $P(m, \delta)$ is the factor m in front of the left-hand cut operator. Thus it can be trivially extended to a holomorphic operator function, by just allowing complex multipliers in the space $H(\mu, 1)$, so that it consists of complex-valued Hölder continuous functions on $u \in (0, 1)$.

But then it can be proved from the implicit variable theorem that each solution $\delta(u)$ of Eq. (7.32) has an analytical dependence on the complex variable m . The region of analyticity, Ω , is moreover only bounded by points where the Fréchet derivative $P'_\eta(m, \delta)$ ceases to be continuous, or to be a $\emptyset$ operator, or possibly where κ , α or β for this $\emptyset$ operator change. Since we have shown that the last points are isolated, and that $P'_\eta(m, \delta)$ is a $\emptyset$ operator for almost all m , we can conclude that Ω is a connected region in the complex m plane containing the whole real positive m axis, except possibly for some isolated points (see, however, Appendix 3).

The Fréchet derivative $P'_\eta(m, \delta)$ depends on m explicitly, and implicitly through δ . As just discussed, both these dependences are analytical in the region Ω , so we can employ the theorem 3.5 given above. Thus κ is given by Eq. (7.31) for almost all real positive m . QED.

We finally discuss the Fréchet derivative when R is held fixed. For any class 1 solution δ this case is similar to the case $\gamma = 0$ just treated, because of the asymptotic form (6.6) of the function B in the kernel of the integral operator.

We shall treat this case by approximating the functions A and B by functions with a nicer asymptotic behaviour, corresponding to a $\gamma > 0$, say $\gamma = 1$. Let thus

$$\begin{cases} A_n(u) = \frac{\eta_n(u)}{g(u)} \cos 2\delta_n(u) \in H(\mu, 2) \\ B_n(u) = i \frac{\eta_n(u)}{g(u)} \sin 2\delta_n(u) \in H_0(\mu, 2) \\ \eta_n(u) \in H(\mu, 2), \delta_n(u) \in H(\mu, 2), \end{cases} \quad (7.33)$$

and let

$$A_n(u) \xrightarrow{n \rightarrow \infty} A(u), \quad B_n(u) \xrightarrow{n \rightarrow \infty} B(u) \text{ pointwise,} \quad (7.34)$$

where A and B are the functions given by Eq. (6.4). Such a set of functions can always be chosen, since $H_0(\mu, 2)$ is dense in $H_0(\mu, 1)$.

But then, in fact, operators G_n^{-1} corresponding to the ones of Eq. (7.10) can be constructed by substituting η_n, δ_n for η, δ in Eqs. (7.6) and (7.7). These operators act as inverses of $P'_\eta(m, 1, \delta_n)$, and

$$P'_\eta(m, 1, \delta_n) \delta \delta(n) \xrightarrow{n \rightarrow \infty} P'(m, \delta) \delta \delta(n) \text{ pointwise.} \quad (7.35)$$

Thus we have the same situation as above for $P'_\eta(m, \frac{1}{n}, \delta_n)$ as $n \rightarrow \infty$, and we can draw the same conclusion as there, namely that the operator $P'(m, \delta)$ obtained in the limit is a $\emptyset$ operator for almost all m .

About the index of this $\emptyset$ operator we know from Section 6 that

$$P'(0, \delta) \text{ has index } \kappa = \begin{cases} 2\nu+1 & \text{for } 2\sigma R_0 \geq 1 \\ 2\nu & \text{for } 2\sigma R_0 < 1 \end{cases} \quad (7.36)$$

and

$$P'(1, \delta) \text{ has index } \kappa = \begin{cases} 2\nu+1 & \text{for } 4\sigma R_0 \geq 1 \\ 2\nu & \text{for } 4\sigma R_0 < 1. \end{cases} \quad (7.37)$$

However, when σ and m are connected by Eq. (7.31), that is when δ is a solution to the non-linear problem, only the case $\kappa = 2\nu+1$ comes into question, for any $\gamma \geq 0$.

For $m \neq 0$ or 1 we repeat the above argument about analytical dependence on m . The result is that $P'(m, \delta)$ has index $\kappa = 2\nu+1$ for almost all real positive m .

8. - RESULTS AND DISCUSSION

We have studied the solutions to the dispersion relation A.1 or A.2 and the unitarity condition B, given in the Introduction. The main results about those solutions are the following.

- I - The solutions fall into two distinct classes, according to their large s behaviour. The asymptotic behaviour is like an inverse logarithm for class 1 solutions [Eq. (2.11)] and like an inverse power for class 2 solutions [Eqs. (2.19) and (2.20)].

The two classes can also be characterized with the integral

$$\gamma = \int_4^{\infty} \text{Im} f(s) ds. \quad (8.1)$$

Class 2 consists of those solutions for which this integral is convergent. They are well behaved, and their asymptotic form contains information about the absorptive part at low energy. Class 1, on the other hand, consists of those solutions for which the integral (8.1) is divergent. They are dominated for large s by their own asymptotic form, so that this form can be determined self-consistently, and does not contain any information about the absorptive part at low energy.

The next result is :

- II - Class 1 solutions have a complete asymptotic expansion in inverse powers of $\ln s$. This expansion has exactly one free parameter, which does not enter the leading term, however. The parameter is related to the scattering length, but contains in general also other contributions.

About the local multiplicity of the solutions to our non-linear problem we get for class 2 solutions the same result as Lovelace, presented in the Introduction. For class 1 solutions, we get :

- III - If in our non-linear problem we change dispersion relation A.1 to dispersion relation A.2 with $m = 2$, all other characteristics of the problem remaining unchanged, the local multiplicity does not change.

IV - The local multiplicity may depend on whether we ask for solutions with the inelasticity $\eta(s)$ given, or the inelasticity ratio $R(s)$ given in the unitarity condition B.

For

$$R(s) \underset{s \rightarrow \infty}{\sim} R_0 (\ln s)^\gamma, \quad \gamma \geq 0, \quad R_0 > 0 \quad (R_0 \geq 1 \text{ for } \gamma = 0) \quad (8.2)$$

and $\mathcal{J}(s)$ a class 1 solution satisfying

$$\mathcal{J}(s) \underset{s \rightarrow \infty}{\sim} \pi \nu + \mathcal{O}((\ln s)^{-1-\gamma}), \quad (8.3)$$

we get the following more specific results :

V - If $\eta(s)$ is prescribed, the local multiplicity of the class 1 solution $\mathcal{J}(s)$ to our non-linear problem is, for almost all $m \geq 0$, given by

$$\kappa = \begin{cases} 2\nu + 1 & \text{if } \gamma = 0 \text{ and } R_0 \leq 2 \\ 2\nu & \text{" } \gamma = 0 \text{ " } R_0 > 2 \\ 2\nu & \text{" } \gamma > 0. \end{cases} \quad (8.4)$$

VI - If $R(s)$ is prescribed, the local multiplicity of the class 1 solution $\mathcal{J}(s)$ to our non-linear problem is, for almost all $m \geq 0$, given by

$$\kappa = 2\nu + 1 \quad \text{for any } \gamma \text{ and } R_0. \quad (8.5)$$

We have not been able to exclude the possibility that for some isolated m values the above result might not hold. This fact is related to the possibility of spontaneous symmetry breaking, discussed by Lovelace⁵⁾. We propose to resolve this difficulty for each particular interesting case by numerical calculations.

A numerical solution of our non-linear problem with the Newton-Kantorovich ¹⁴⁾ method would, at the same time as it resolved this difficulty, also provide an existence proof for the solution of that particular case.

As a final result we formulate what we have to do, besides using analyticity, unitarity and crossing symmetry, in order to get a unique solution for the S wave amplitude :

- (i) decide whether we want a class 1 or a class 2 solution ;
- (ii) fix the total variation of the phase shift (to $\pi \nu$) ;
- (iii) determine a number of parameters $[2\nu]$ for class 2 solutions, 2ν or $2\nu+1$ according to Eqs. (8.4) and (8.5) for class 1 solutions] from auxiliary conditions.

If other partial waves or coupled channels are included, the number of arbitrary parameters increases, as discussed by Lovelace ⁵⁾.

About the general question of the existence of solutions to our non-linear problem we have, unfortunately, very little to say. Fixed point theorems ⁹⁾ give information about small solutions, but do not seem to be directly applicable to class 1 solutions.

The class 2 solution with $\nu = 0$ is locally unique ($\alpha = 2\nu = 0$), which for elastic unitarity, or for $R(s)$ given, only means that the solution $f \equiv 0$ is isolated. If $\eta \neq 1$ is given, the fixed point theorem would presumably give a unique class 2 solution $f \neq 0$ with $\nu = 0$ for either of the dispersion relations A.1 or A.2, but no information about the other solutions (see, however, Atkinson and Warnock ⁹⁾).

We believe the Newton-Kantorovich method to be the best to prove the existence of class 1 solutions or class 2 solutions with $\nu > 0$. The present work contains some necessary prerequisite for such a study, since one must know the multiplicity, and work in the factor space $\mathcal{B}$ of Eq. (6.23) in order to have a well-defined inverse operator G^{-1} , as necessary in the Newton-Kantorovich method. We hope to be able to present such a proof in a future publication.

ACKNOWLEDGEMENTS

I want to thank Professor A. Martin for suggesting this study, for taking part in the earlier parts of it, and for reading the manuscript. I also want to acknowledge the hospitality extended to me at the Theoretical Study Division at CERN, where this work was done, and thank the Swedish Atomic Research Council for a grant.

APPENDIX 1

Our principal aim in this Appendix is to derive asymptotic expansions in $(\ln s)^{-n}$ for large s of integrals of the type

$$I(A) = \int_4^{\infty} \frac{ds'}{(s'+A)(\ln s')^{j+2}} \quad (\text{A.1.1})$$

for integer $j \geq 0$ and $|A| = \mathcal{O}(s)$ as $s \rightarrow \infty$.

Beginning with positive A , $I(A)$ is obviously convergent, approaches zero as $A \rightarrow \infty$, and we have

$$I(A) = \frac{1}{A} \int_4^A \frac{ds'}{(\ln s')^{j+2}} \left(1 + \frac{s'}{A}\right)^{-1} + \int_A^{\infty} \frac{ds'}{s'(\ln s')^{j+2}} \left(1 + \frac{A}{s'}\right)^{-1},$$

which gives

$$I(A) = \sum_{i=0}^{\infty} \frac{(-1)^i}{A^{i+1}} \int_4^A \frac{s'^i ds'}{(\ln s')^{j+2}} + \sum_{i=0}^{\infty} \frac{(-A)^i}{A} \int_A^{\infty} \frac{ds'}{s'^{i+1}(\ln s')^{j+2}}. \quad (\text{A.1.2})$$

One of those integrals is easy,

$$\int_A^{\infty} \frac{ds'}{s'(\ln s')^{j+2}} = \frac{1}{(j+1)(\ln A)^{j+1}},$$

and for the other ones we use partial integrations to get

$$\int_A^{\infty} \frac{ds'}{s'^{i+1}(\ln s')^{j+2}} = - \sum_{k=0}^N \frac{(j+k+1)!}{(-i)^{k+1}(j+1)!} \frac{A^{-i}}{(\ln A)^{j+k+2}} + R_N(A),$$

where (N is an arbitrary positive integer)

$$R_N(A) = \frac{(j+N+2)!}{(-i)^{N+1}(j+1)!} \int_A^{\infty} \frac{ds'}{s'^{i+1}(\ln s')^{j+N+3}} = \mathcal{O}\left(\frac{A^{-i}}{(\ln A)^{j+N+3}}\right) \text{ as } A \rightarrow \infty.$$

Similarly

$$\int_4^A \frac{s'^i ds'}{(\ln s')^{j+2}} = \sum_{h=0}^N \frac{(j+h+1)!}{(i+1)^{h+1} (j+1)!} \frac{A^{i+1}}{(\ln A)^{j+h+2}} + \mathcal{O}\left(\frac{A^{i+1}}{(\ln A)^{j+N+3}}\right)$$

as $A \rightarrow \infty$.

Substituting these expressions for the integrals into Eq. (A.1.2), we get the following asymptotic expression for $I(A)$:

$$I(A) = \frac{1}{(j+1)(\ln A)^{j+1}} + \sum_{i=0}^{\infty} \sum_{h=0}^N \frac{(-1)^i (j+h+1)!}{(i+1)^{h+1} (j+1)!} \frac{1 + (-1)^{h+1}}{(\ln A)^{j+h+2}} + \mathcal{O}\left(\frac{1}{(\ln A)^{j+N+3}}\right)$$

Here, the summation over i can be performed to get the asymptotic series

$$I(A) = \frac{1}{(j+1)(\ln A)^{j+1}} + \sum_{h=0}^{\infty} \frac{2(j+2h+2)!}{(j+1)!} \frac{\pi_h^-}{(\ln A)^{j+2h+3}} + o((\ln A)^{-n}) \quad (\text{A.1.3})$$

as $A \rightarrow \infty$. The small ordo symbol is here used to indicate that the rest term of the series is asymptotically smaller than any inverse power of $\ln A$.

The constants $p_k^{\pm}$ are defined by

$$\pi_h^{\pm} = \sum_{i=0}^{\infty} \frac{(\pm 1)^i}{(i+1)^{2h+2}}, \quad h=0, 1, 2, \dots, \quad (\text{A.1.4})$$

and are connected to the Bernoulli numbers B_k ,

$$\pi_h^+ = \frac{\pi^{2h+2} 2^{2h+1}}{(2h+2)!} B_{h+1}, \quad \pi_h^- = \frac{\pi^{2h+2} (2^{2h+1} - 1)}{(2h+2)!} B_{h+1}. \quad (\text{A.1.5})$$

The first few constants are

$$\begin{aligned} \pi_0^+ &= \frac{\pi^2}{6}, \quad \pi_1^+ = \frac{\pi^4}{90}, \quad \pi_2^+ = \frac{\pi^6}{945}, \\ \pi_0^- &= \frac{\pi^2}{12}, \quad \pi_1^- = \frac{7\pi^4}{720}, \quad \pi_2^- = \frac{31\pi^6}{30240}. \end{aligned}$$

For negative $A < -4$ the integral (A.1.1) is defined by its Cauchy principal value. The contribution to the integral from near the singularity is then

$$\begin{aligned} & \int_{|A|-\varepsilon}^{|A|+\varepsilon} \frac{ds'}{(s'+A)(\ln s')^{j+2}} = \int_{-\varepsilon}^{\varepsilon} \frac{dx}{x [\ln|A| + \ln(1 - \frac{x}{A})]^{j+2}} = \\ &= (\ln|A|)^{-j-2} \left[\int_{-\varepsilon}^{\varepsilon} \frac{dx}{x} - (j+2) \int_{-\varepsilon}^{\varepsilon} \frac{dx}{x} \left\{ \frac{\ln(1 - \frac{x}{A})}{\ln|A|} + \mathcal{O}\left(\frac{\ln^2(1 - \frac{x}{A})}{\ln^2|A|}\right) \right\} \right] = \\ &= (\ln|A|)^{-j-2} \left[\int_{-\varepsilon}^{\varepsilon} \frac{dx}{x} + \frac{j+2}{A \ln|A|} \int_{-\varepsilon}^{\varepsilon} dx \left\{ 1 + \mathcal{O}\left(\frac{x}{A}\right) \right\} \right] = \mathcal{O}\left(\frac{\varepsilon}{A (\ln|A|)^{j+3}}\right) \end{aligned} \quad (\text{A.1.6})$$

as $\varepsilon/A \rightarrow 0$. This proves that the contribution from the region near the singularity can be included in the rest term of an asymptotic series like Eq. (A.1.3). But this also means that it is unnecessary to subtract the integral (A.1.6) explicitly from $I(A)$ before doing the calculations leading to such an asymptotic series. We only have to take care of the different sign of A that we now have. This is easily done, and the result is

$$I(A) = \frac{1}{(j+1)(\ln|A|)^{j+1}} - \sum_{h=0}^{\infty} \frac{2(j+2h+2)!}{(j+1)!} \frac{\pi_h^+}{(\ln|A|)^{j+2h+3}} + \mathcal{O}((\ln A)^{-n}) \quad (\text{A.1.7})$$

as $A \rightarrow -\infty$.

The results (A.1.3) and (A.1.7) generalize trivially to any real $j > -1$. For $j \leq -1$ we must introduce a subtraction, and define

$$I(A) = \int_4^{\infty} \frac{A \cdot ds'}{s'(s'+A)(\ln s')^{j+2}}. \quad (\text{A.1.8})$$

For this case we will only need the leading asymptotic behaviour, which is easily obtained from the appropriately modified Eq. (A.1.2). The result is

$$I(A) = \begin{cases} \ln(\ln|A|) + \mathcal{O}(1) & \text{for } j = -1 \\ \frac{1}{(-j-1)} (\ln|A|)^{-j-1} + \mathcal{O}((\ln|A|)^{-j-2}) & \text{for } j < -1 \end{cases} \quad (\text{A.1.9})$$

as $A \rightarrow \pm\infty$.

Another integral, of which we will also need the leading asymptotic form is

$$I_{\beta}(A) = \int_4^{\infty} \frac{ds'}{(s'+A)s'^{\beta}(\ln s')^{j+2}}, \quad \beta > 0. \quad (\text{A.1.10})$$

The above method gives for $A \rightarrow \pm\infty$

$$I_{\beta}(A) = \begin{cases} \frac{(-1)^{\beta}}{(j+1)A^{\beta}(\ln|A|)^{j+1}} + \mathcal{O}\left(\frac{1}{A^{\beta}(\ln|A|)^{j+3}}\right) & \text{for integer } \beta \\ \frac{C_{\beta}}{A^{\beta}(\ln|A|)^{j+2}} + \mathcal{O}\left(\frac{1}{A^{\beta}(\ln|A|)^{j+3}}\right) & \text{for noninteger } \beta. \end{cases} \quad (\text{A.1.11})$$

Here, C_{β} is a finite but complicated β dependent constant, the value of which is not of importance to the present work.

APPENDIX 2

In this Appendix we have collected some proofs of Banach space properties. We begin by proving that $H_0(\mu, k)$, defined in Section 5, is a complete space.

If the norm (5.3) is finite, it follows that

$$|\varphi(u) - \varphi(1)| \leq \|\varphi\| \left(\ln \frac{4}{1-u} \right)^{-k} < \|\varphi\|, \quad (\text{A.2.1})$$

which, since $\varphi(1) = 0$ for $\varphi \in H_0(\mu, k)$, implies that

$$H_0(\mu, k) \subset C(k) \subset C(0), \quad k > 0. \quad (\text{A.2.2})$$

Assume that $\{\varphi_n\}$ is a Cauchy sequence in $H_0(\mu, k)$, that is

$$\|\varphi_{n+p} - \varphi_n\| \leq \varepsilon_n, \quad \text{any } n, \quad \varepsilon_n \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (\text{A.2.3})$$

Then Eq. (A.2.1) implies that it is a Cauchy sequence also in $C(0)$, which is known to be complete. But then there exists a φ in $C(0)$, such that

$$|\varphi(u) - \varphi_n(u)| \leq \varepsilon_n. \quad (\text{A.2.4})$$

Moreover, Eq. (A.2.3) implies that

$$|\varphi_{n+p}(u_1) - \varphi_{n+p}(u_2) - \varphi_n(u_1) + \varphi_n(u_2)| \leq \varepsilon_n \left| \frac{u_1 - u_2}{1 - u_2} \right|^\mu \left(\ln \frac{4}{1 - u_2} \right)^{-k}. \quad (\text{A.2.5})$$

Since the right-hand side of this equation is independent of p , we can take the limit $p \rightarrow \infty$. Then according to Eq. (A.2.4) we get

$$|\varphi(u_1) - \varphi(u_2) - \varphi_n(u_1) + \varphi_n(u_2)| \leq \varepsilon_n \left| \frac{u_1 - u_2}{1 - u_2} \right|^\mu \left(\ln \frac{4}{1 - u_2} \right)^{-k}, \quad (\text{A.2.6})$$

which implies that

$$\|\varphi\| < \infty, \quad \|\varphi - \varphi_n\| \leq \varepsilon_n \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (\text{A.2.7})$$

Thus $H_0(\mu, k)$ is complete.

The proof above is almost identical to the one, for a smaller space, of Muskhelishvili [p. 132 of Ref. 10]. It works equally well for $H(\mu, k)$ and $C(k)$.

Next we prove that the Banach space $H(\mu, k)$ is also a normed ring. This follows if we can prove that the product of two elements φ and ψ of $H(\mu, k)$ is again an element of $H(\mu, k)$, and that

$$\|\varphi\psi\| \leq \|\varphi\| \|\psi\|. \quad (\text{A.2.8})$$

Let us distinguish the two parts of the norm (5.2) by indices

$$\|\varphi\| = \|\varphi\|_c + \|\varphi\|_0. \quad (\text{A.2.9})$$

Then $\varphi\psi$ is Hölder continuous, and

$$\|\varphi\psi\|_0 \leq \sup_{u_1, u_2} \frac{[|\varphi(u_1) - \varphi(u_2)| |\psi(u_1)| + |\varphi(u_2)| |\psi(u_1) - \psi(u_2)|]}{\left| \frac{u_1 - u_2}{1 - u_2} \right|^\mu \left(\ln \frac{4}{1 - u_2} \right)^{-k}} \leq \|\varphi\|_0 \|\psi\|_c + \|\varphi\|_c \|\psi\|_0. \quad (\text{A.2.10})$$

Thus

$$\|\varphi\psi\| \leq \|\varphi\|_c \|\psi\|_c + \|\varphi\|_0 \|\psi\|_c + \|\varphi\|_c \|\psi\|_0 \leq \|\varphi\| \|\psi\|. \quad \text{Q.E.D.} \quad (\text{A.2.11})$$

In Section 7 we need the fact that the embedding (A.2.2) of $H_0(\mu, k)$ in $C(0)$ is a compact map. To prove this, we prove that the bounded set $\|\varphi\| \leq C$ in $H_0(\mu, k)$ is a compact set in $C(0)$.

Any function of this set is uniformly bounded by Eq. (A.2.1), and further satisfies

$$|\varphi(u_1) - \varphi(u_2)| \leq C \frac{\delta^k}{(1-u_2)^k} \left(\ln \frac{4}{1-u_2} \right)^{-k}, \quad \delta = u_1 - u_2. \quad (\text{A.2.12})$$

Now

$$\max_{0 \leq u_2 \leq 1-\delta} \frac{\delta^k}{(1-u_2)^k} \left(\ln \frac{4}{1-u_2} \right)^{-k} = \left(\ln \frac{4}{\delta} \right)^{-k} \xrightarrow{\delta \rightarrow 0} 0 \quad (\text{A.2.13})$$

for any $k > 0$. Thus for any $\varepsilon > 0$ there exists a $\delta > 0$ such that

$$|\varphi(u_1) - \varphi(u_2)| \leq \varepsilon \quad \text{for } |u_1 - u_2| \leq \delta = 4 \exp \left[- \left(\frac{4}{\varepsilon} \right)^{\frac{1}{k}} \right] > 0. \quad (\text{A.2.14})$$

for any function of the set $\|\varphi\| \leq C$. But this is the statement that this set of functions is equicontinuous. Then the Arzelà-Ascoli theorem [Ref. 12) IV.6.7] tells us that the set is compact. Q.E.D.

Next, we want to prove that the operator K_R ,

$$(K_R \varphi)(u) = - \frac{g(u)}{\pi} \rho \int_0^1 \frac{(1-u') \varphi(u') du'}{(1-u')(u'-u)}, \quad (\text{A.2.15})$$

is a bounded operator from $H_0(\mu, k+1)$ to $H_0(\mu, k)$, $k > 0$, $0 < \mu < \frac{1}{2}$.

We write

$$(K_R \varphi)(u) = \mathcal{Y}_1(u) - \mathcal{Y}_2(u), \quad (\text{A.2.16})$$

where

$$\mathcal{Y}_1(u) = \frac{g(u)}{\pi} \int_0^1 \frac{\varphi(u') du'}{u'-1}, \quad \mathcal{Y}_2(u) = \frac{g(u)}{\pi} \rho \int_0^1 \frac{\varphi(u') du'}{u'-u}. \quad (\text{A.2.17})$$

Since $\varphi(u) = \sqrt{u}$ and the integral is convergent for any $\varphi \in H_0(\mu, k+1)$, $k > 0$, $J_1(u)$ is obviously a Hölder continuous function on $0 \leq u \leq 1$ with $\mu \leq \frac{1}{2}$. Further, as proved in Muskhelishvili [Ref. 10], pp. 46-48, the integral in $J_2(u)$ is a Hölder continuous function on the open interval $0 < u < 1$, with the same Hölder index μ as φ . At $u = 0$ the integral has a logarithmic singularity, but since it is to be multiplied with $\sqrt{u}$, we obtain that $J_2(u)$ is Hölder continuous on $0 \leq u < 1$ with index $\mu < \frac{1}{2}$. This implies that

$$|(K_R \varphi)(u_1) - (K_R \varphi)(u_2)| \leq C_R(\varepsilon) C \left| \frac{u_1 - u_2}{1 - u_1} \right|^\mu, \quad 0 \leq u_1 \leq u_2 \leq 1 - \varepsilon, \quad \varepsilon > 0. \quad (\text{A.2.18})$$

as soon as $\varphi \in H_0(\mu, k+1)$, $\mu < \frac{1}{2}$, $k > 0$, with norm $\|\varphi\| \leq C$. For the constant $C_R(\varepsilon)$, which may be ε dependent, one can, with some extra work, following Muskhelishvili, obtain a finite, explicit value. However, here we only need the fact that it is finite for any fixed $\varepsilon > 0$.

What happens when $\varepsilon \rightarrow 0$? For the behaviour of the integral (A.2.15) when $u \rightarrow 1$ we can use the analysis of Appendix 1. Equation (A.1.9) gives

$$(K_R \varphi)(u) \underset{u \rightarrow 1}{\sim} \frac{C}{\pi h} \left(\ln \frac{4}{1-u} \right)^{-h} \quad (\text{A.2.19})$$

if

$$\varphi(u) \underset{u \rightarrow 1}{\sim} C \left(\ln \frac{4}{1-u} \right)^{-h-1}. \quad (\text{A.2.20})$$

But then

$$|(K_R \varphi)(u_1) - (K_R \varphi)(u_2)| \leq C'_R C \left(\ln \frac{4}{1-u_1} \right)^{-h}, \quad 0 \leq u_1 \leq u_2 \leq 1, \quad (\text{A.2.21})$$

for some finite constant C'_R , since, besides fulfilling Eq. (A.2.19), $(K_R \varphi)(u)$ is bounded on $0 \leq u < 1$. This shows that if we let $\varepsilon \rightarrow 0$, $u_2 \rightarrow 1$ and hold u_1 constant in Eq. (A.2.18), the equation is still valid, with a finite $C_R(0)$. Finally, we put

$$\bar{C}_R = \max \{ C_R(0) (\ln 4)^h, C'_R \} \quad (\text{A.2.22})$$

and combine Eqs. (A.2.18) and (A.2.21) to get

$$|(K_R \varphi)(u_1) - (K_R \varphi)(u_2)| \leq \bar{C}_R C \left| \frac{u_1 - u_2}{1 - u_1} \right|^h \left(\ln \frac{4}{1 - u_1} \right)^{-h}, \quad 0 \leq u_1 \leq u_2 \leq 1, \quad (\text{A.2.23})$$

which is to say that

$$\|K_R\| = \sup_{\varphi} \frac{\|K_R \varphi\|_h}{\|\varphi\|_{h+1}} \leq \bar{C}_R. \quad \text{Q.E.D.} \quad (\text{A.2.24})$$

The operator K_L has, for the case of dispersion relation A.2, the form

$$(K_L \varphi)(u) = -\frac{m \varphi(u)}{\pi} \int_0^1 \frac{(1-u') \varphi(u') du'}{(1-u')(1-u'u')} = \frac{m \varphi(u)}{\pi} \mathcal{J}_3(u). \quad (\text{A.2.25})$$

For any $\varphi \in C(k+1)$, $k > 0$, the integral $\mathcal{J}_3(u)$ is convergent for $0 \leq u < 1$. As $u \rightarrow 1$, we obtain from Appendix 1 the asymptotic form

$$\mathcal{J}_3(u) \sim_{u \rightarrow 1} \frac{C}{h} \left(\ln \frac{4}{1-u} \right)^{-h} \text{ if } \varphi(u) \sim_{u \rightarrow 1} C \left(\ln \frac{4}{1-u} \right)^{-h-1}, \quad (\text{A.2.26})$$

which implies that in analogy with Eq. (A.2.21) we have

$$|\mathcal{J}_3(u_1) - \mathcal{J}_3(u_2)| \leq C_L' C \left(\ln \frac{4}{1-u_1} \right)^{-h}, \quad 0 \leq u_1 \leq u_2 \leq 1, \quad (\text{A.2.27})$$

for some finite constant C_L' . Moreover, we get directly from Eq. (A.2.25) that

$$\begin{aligned} |\mathcal{J}_3(u_1) - \mathcal{J}_3(u_2)| &= \left| (u_1 - u_2) \int_0^1 \frac{\varphi(u') du'}{(1-u_2 u')(1-u_1 u')} \right| \leq \\ &\leq C_L(\varepsilon) C \left| \frac{u_1 - u_2}{1 - u_1} \right| \text{ for } 0 \leq u_1 \leq u_2 \leq 1 - \varepsilon, \varepsilon > 0. \end{aligned} \quad (\text{A.2.28})$$

Combining the last two equations gives that $\mathcal{J}_3(u)$ has a finite norm in $H_0(1, k)$ for any finite element φ of $C(k+1)$. This implies for K_L that its norm as an operator from $C(k+1)$ to $H_0(\mu, k)$, $\mu \leq \frac{1}{2}$, is bounded by $\bar{C}_L$, where

$$\bar{C}_L = \frac{m}{\pi} \max \{ C_L(0) \left(\ln 4 \right)^h, C_L' \}. \quad (\text{A.2.29})$$

For the case of dispersion relation A.1, the form of K_L is

$$(K_L \varphi)(u) = -\frac{2g(u)}{\pi} \int_0^1 \frac{1-u}{u(1-u')^2} \ln\left[1 + \frac{u}{1-u}(1-u')\right] \varphi(u') du' = \frac{2g(u)}{\pi} \mathcal{J}_4(u). \quad (\text{A.2.30})$$

This integral is also convergent for $0 \leq u < 1$ for any $\varphi \in C(k+1)$, $k > 0$. As $u \rightarrow 1$, the integrand has the same asymptotic form as the integrand of Eq. (A.2.25), and the asymptotic analysis of Appendix 1 shows that

$$\mathcal{J}_4(u) \sim \frac{C}{h} \left(\ln \frac{4}{1-u}\right)^{-h} \text{ if } \varphi(u) \sim C \left(\ln \frac{4}{1-u}\right)^{-h-1}. \quad (\text{A.2.31})$$

This implies that Eq. (A.2.27) is valid also for $\mathcal{J}_4(u)$. Moreover, $\mathcal{J}_4(u)$ has a continuous derivative on $\varepsilon < u < 1-\varepsilon$, since

$$\mathcal{J}_4'(u) = \frac{1}{u} \int_0^1 \left\{ \frac{1}{1-uu'} - \frac{1}{u(1-u')} \ln\left[1 + \frac{u}{1-u}(1-u')\right] \right\} \frac{\varphi(u') du'}{1-u'} \quad (\text{A.2.32})$$

is a uniformly convergent integral on this interval. Thus Eq. (A.2.28) is valid for $\mathcal{J}_4(u)$, provided we cut off the interval at $u_1 = \varepsilon$. However, this cut-off is not necessary, since from the fact that

$$x - \frac{1}{2}x^2 < \ln(1+x) < x, \quad 0 \leq x < 1, \quad (\text{A.2.33})$$

it follows that

$$|\mathcal{J}_4(u_1) - \mathcal{J}_4(u_2)| \leq \frac{u_2}{2(1-u_2)} \int_0^1 |\varphi(u')| du' < \text{const } \varepsilon \text{ for } 0 \leq u_1 \leq u_2 \leq \varepsilon. \quad (\text{A.2.34})$$

But then the boundedness of K_L follows just as for the previous case.

Consider now the difference ΔK ,

$$(\Delta K \varphi)(u) = \frac{2g(u)}{\pi} \int_0^1 \left\{ \frac{1}{1-uu'} - \frac{1}{u(1-u')} \ln\left[1 + \frac{u}{1-u}(1-u')\right] \right\} \frac{(1-u)\varphi(u') du'}{1-u'}, \quad (\text{A.2.35})$$

for $\varphi \in C(k)$, $k > 0$. Let first $u \leq 1-\varepsilon$, $\varepsilon > 0$, and divide the integration interval into two, expanding the logarithm in the second term:

$$\begin{aligned}
 (\Delta K \varphi)(u) &= \frac{2g(u)}{\pi} \int_0^{1-\varepsilon} \left\{ \frac{1}{1-uu'} - \frac{1}{u(1-u')} \ln \left[1 + \frac{u}{1-u} (1-u') \right] \right\} \frac{(1-u)\varphi(u') du'}{1-u'} + \\
 &+ \frac{2g(u)}{\pi} \int_{1-\varepsilon}^1 \left\{ \frac{u}{1-uu'} + O\left(\frac{u}{1-u}\right) \right\} \varphi(u') du' \quad (A.2.36)
 \end{aligned}$$

This shows that there appear no convergence problems at $u' = 1$ when we take $\varphi \in C(k)$ instead of $C(k+1)$, as long as $u \leq 1 - \varepsilon$. Thus $(\Delta K \varphi)(u)$ is Hölder continuous on $0 \leq u \leq 1 - \varepsilon$, just as $K_L \varphi$.

To see what happens when $u \rightarrow 1$ we can go back to Eq. (3.7), which shows that the asymptotic form of the discontinuity on the left-hand cut is the same to leading order for dispersion relations A.1 and A.2 with $m = 2$, but differs in the next order. Since in Eq. (A.2.35) we take the difference between these two cases, the leading term in $K_L \varphi$ is cancelled, but the next appears, that is

$$(\Delta K \varphi)(u) = O\left(\left(\ln \frac{4}{1-u}\right)^{-k-1}\right) \text{ if } \varphi(u) \underset{u \rightarrow 1}{\sim} C \left(\ln \frac{4}{1-u}\right)^{-k-1}. \quad (A.2.37)$$

It follows from Appendix 1, by analogy with Eq. (A.1.9), that this equation is in fact valid for all $k \neq -1$. Thus

$$(\Delta K \varphi)(u) \in H_0(\mu, k) \text{ for any } \varphi \in C(k). \text{ Q.E.D.} \quad (A.2.38)$$

Finally, we want to prove that the operator G_R^{-1} of Eq. (6.24) is a bounded operator in $H_0(\mu, 1)$. We can write

$$G_R^{-1}(\delta) \delta f(s) = \frac{\cos 2\delta(s)}{\eta(s)} \delta f(s) - \frac{s^{\infty}}{\eta(s)|D(s)|^2} K_R \left\{ \frac{|D(s')|^2 \sin \Delta(s')}{s'^{\infty} g(s')} \delta f(s') \right\}(s). \quad (A.2.39)$$

The factor multiplying δf inside the brackets in the last term has the asymptotic form [cf. Eqs. (6.11), (6.15'), (6.18)]

$$\frac{|D(s')|^2 \sin \Delta(s')}{s'^{\infty} g(s')} \underset{s' \rightarrow \infty}{\sim} \begin{cases} \text{const } \frac{1}{s'} (\ln s')^{2\sigma R_0 - 1} & \text{for } \delta \text{ class 1} \\ \text{const } s'^{-\mu} & \text{for } \delta \text{ class 2} \end{cases} \quad (A.2.40)$$

and is finite for all s' , since the zero at threshold of $\varphi(s')$ is cancelled by a similar zero of $\Delta(s')$. Thus, since all the factors are Hölder continuous, K_R operates on a quantity with a finite norm in $H_0(\mu, 2)$. Thus it gives a function with a finite norm in $H_0(\mu, 1)$, which moreover has the asymptotic form

$$K_R \left\{ \frac{|D(s')|^2 \sin \Delta(s')}{s'^2 \varphi(s')} \delta f(s') \right\}(s) \underset{s \rightarrow \infty}{\sim} \begin{cases} \text{const } \frac{1}{s} (\ln s)^{2\sigma R_0 - 1} & \text{for } \delta \text{ class 1} \\ \text{const } s^{-\mu} & \text{for } \delta \text{ class 2.} \end{cases} \quad (\text{A.2.41})$$

Thus

$$G_R^{-1}(\delta) \delta f(s) \in H_0(\mu, 1) \text{ if } \delta f(s) \in H_0(\mu, 1) \text{ and } \eta(s) \neq 0. \quad (\text{A.2.42})$$

With some extra work, an explicit bound for the norm of G_R^{-1} as an operator in $H_0(\mu, 1)$ can be obtained, but here we only need the fact that it is finite.

APPENDIX 3

In this Appendix we shall draw some conclusions about the spectrum of the operator (7.8) for $\gamma = 0$, that is

$$\mathcal{G}(m, \delta) = G_{\eta}^{-1}(0, 0, \delta) P'_{\eta, L}(m, 0, \delta). \quad (\text{A.3.1})$$

Since G_{η}^{-1} is bounded uniformly in δ and m independent, we first study $P'_{\eta, L}$. Decompose it into an asymptotic and a compact part :

$$P'_{\eta, L}(m, 0, \delta) = Q_A(m, \delta) + Q_C(m, \delta) \quad (\text{A.3.2})$$

Here

$$Q_A(m, \delta) \delta \delta(u) = -2m\sigma \int_0^1 \frac{(1-u) \delta \delta(u') du'}{(1-u')(1-u'm) \ln \frac{4}{1-u'}} \quad (\text{A.3.3})$$

and

$$Q_C(m, \delta) \delta \delta(u) = -\frac{m}{\pi} \int_0^1 \left[\frac{\eta(u')}{g(u')} \sin 2\delta(u') - 2\pi\sigma \left(\ln \frac{4}{1-u'} \right)^{-1} \right] \frac{(1-u) \delta \delta(u') du'}{(1-u')(1-u'm)}, \quad (\text{A.3.4})$$

defining bounded operators according to the considerations of Appendix 2.

Q_C is compact also for $\gamma = 0$, since the asymptotic behaviour of the expression in square brackets is $(\ln 4/1-u')^{-2}$. Q_C contains the finite energy contributions, and is approximately proportional to m , even though δ is m dependent. The discrete spectrum of Q_C thus moves with m , the eigenvalues being almost proportional to m .

Q_A , on the other hand, is non-compact in the limit $\gamma \rightarrow 0$, and is strictly proportional to $m\sigma$. We expect it to have a continuous spectrum, which is, however, bounded since Q_A is a bounded operator. If σ is held constant, this spectrum grows without bound when m grows. However, if σ is coupled to m by letting $\delta(u)$ be a solution to our non-linear problem, that is

$$\sigma = \frac{1}{R_0(1+m)}, \quad (\text{A.3.5})$$

the continuous spectrum of Q_A is restricted to a finite interval for all m .

Using the identity

$$[1 + G_\eta^{-1}(Q_A + Q_C)]^{-1} = [1 + G_\eta^{-1}Q_C]^{-1} \{1 + G_\eta^{-1}Q_A [1 + G_\eta^{-1}Q_C]^{-1}\}^{-1} \quad (\text{A.3.6})$$

we can now draw conclusions about the spectrum of the operator $\mathcal{G}$ of Eq. (A.3.1). First, it has a discrete spectrum analogous to the spectrum of Q_C ; in particular the eigenvalues move almost proportional to m . Second, avoiding the discrete eigenvalues so that

$$[1 + G_\eta^{-1}Q_C]^{-1} \quad (\text{A.3.7})$$

is a well-defined operator, we see that it has also a continuous spectrum, like Q_A . In particular, the continuous spectrum is restricted to a finite interval for all m if σ is given by Eq. (A.3.5).

In Section 7 we assumed that the point -1 is not in the spectrum of the operator $\mathcal{G}$. That it is not in the discrete spectrum can obviously be avoided by excluding a number of isolated m values, as done in Section 7. That it is not in the continuous spectrum, that $\mathcal{G}$ is supposed to acquire in the limit $\gamma \rightarrow 0$, seems a reasonable assumption due to the boundedness of this spectrum. However, this is a very intricate point, and the argument given is not a complete proof. That we still believe in this assumption rests on two facts. First, for $m = 1$ we solved the problem exactly in Section 6, and that solution shows that for this m value, the continuous spectrum does not contain -1 . Second, the exact results for $m = 0$ and $m = 1$ are consistent with no m dependence in, e.g., the rule

$$\kappa = \begin{cases} 2\nu + 1 & \text{for } R_0 \leq 2 \\ 2\nu & \text{" } R_0 > 2, \end{cases} \quad (\text{A.3.8})$$

which in turn can be motivated by the balance between the two effects of the left-hand cut, to increase the index and to decrease the asymptotic form. This would correspond to the continuous spectrum being proportional to $(1+m)\sigma$ instead of $m\sigma$, which is reasonable, since the right-hand cut also contributes a term with the same asymptotic form as Eq. (A.3.3), but without the m .

REFERENCES

- 1) O. Brander - Nuovo Cimento 6A, 371 (1971) [the sign in front of the last logarithm in Eq. (6) of this paper is incorrect. This does not, however, influence any of the conclusions of the paper].
- 2) A. Martin - Nuovo Cimento 47A, 265 (1967) [see also the reference list of Ref. 1] ;
S.M. Roy - Phys.Letters 36B, 353 (1971).
- 3) J.L. Basdevant, C.D. Froggatt and J.L. Petersen - Phys.Letters 41B, 173, 178 (1972).
- 4) M. Cini and S. Fubini - Ann.Phys. 3, 352 (1960) ;
C. Lovelace - Nuovo Cimento 21, 305 (1961) ;
D.V. Shirkov, V.V. Serebryakov and V.A. Meshcheryakov - "Dispersion Theories of Strong Interactions at Low Energy", North Holland, Amsterdam (1969).
- 5) C. Lovelace - Commun.Math.Phys. 4, 261 (1967).
- 6) M.M. Vainberg and V.A. Trenogin - Uspekhi Math.Nauk 17, 13 (1962).
- 7) I.C. Gokhberg and M.G. Krein - Uspekhi Math.Nauk 12, 43 (1957) [Am. Math.Soc.Translations 13, 185 (1960)].
- 8) L. Castillejo, R. Dalitz and F. Dyson - Phys.Rev. 101, 543 (1956).
- 9) D. Atkinson - Nuclear Phys. B7, 375 (1968) ; B8, 377 (1968) ; B13, 415 (1969) ; B23, 397 (1970) ;
D. Atkinson and R.L. Warnock - Phys.Rev. 188, 2098 (1969).
- 10) N.I. Muskhelishvili - "Singular Integral Equations", Groningen (1953).
- 11) R.F. Dashen and S.C. Frautschi - Phys.Rev. 135, B1190 (1964).
- 12) N. Dunford and J.T. Schwartz - "Linear Operators", Part 1, New York (1958).
- 13) Assuming that also for $\gamma = 0$ the spectrum of the operator (7.8) does not contain the point -1. For a discussion of the spectrum of this operator, see Appendix 3.
- 14) A.Ts. Amatuni - Nuovo Cimento 58A, 321 (1968) ;
L.V. Kantorovich and G.P. Akilov - "Functional Analysis in Normed Spaces", Oxford (1964).