CERN-DATA HANDLING DIVISION
DD/72/7
K.S. Kölbig
April 1972

PROGRAMS FOR COMPUTING THE LOGARITHM OF THE
GAMMA FUNCTION AND THE DIGAMMA FUNCTION
FOR COMPLEX ARGUMENT

K.S. Kölbig

(Submitted to Computer Physics Communications)

PROGRAMS FOR COMPUTING THE LOGARITHM OF THE
GAMMA FUNCTION AND THE DIGAMMA FUNCTION
FOR COMPLEX ARGUMENT

K.S. Kölbig

CERN, Geneva, Switzerland

Geneva - April 1972

PROGRAM SUMMARY

Title of program (32 characters maximum): CLOGAM, CDIGAM
Computer for which the program is designed and others upon which it is operable
Computer: CDC 6600 Installation: CERN, Geneva
Operating system or monitor under which the program is executed: CDC scope
Programming languages used: FORTRAN IV
High speed store required: 334 (CLOGAM), 232 (CDIGAM)
Is the program overlaid? No
No. of magnetic tapes required: None
What other peripherals are used? Logical unit 2 for error message
No. of cards in combined program and test deck: 161
Card punching code: BCD
Keywords descriptive of the problem and method of solution:
gamma function, logarithm of gamma function , psi function,
digamma function, asymptotic expansion.

Nature of the physical problem

The gamma function $\Gamma(z)$, its logarithm $\ln \Gamma(z)$, and its logarithmic derivative $\psi(z) = d \ln \Gamma(z)/dz$ appear in a wide range of physical applications. We mention here only the Veneziano model and its generalizations in high-energy physics, and the Coulomb phase shift for complex energies.

Method of solution

For $\text{Re } z \geq 7$, the asymptotic expansions are used to compute $\ln \Gamma(z)$ and $\psi(z)$. For other regions of the z -plane, suitable functional relations are used. Care is taken that $\text{Im } \ln \Gamma(z)$ is computed correctly, and not merely modulo 2π .

Restrictions on the complexity on the problem

As the tests show, an accuracy of 12-14 significant digits is normally obtained. For details see Long Write-Up.

Running time

The following running times (in microseconds on the CDC 6600) are typical:

	$\ln \Gamma(z)$ (CLOGAM)	$\psi(z)$ (CDIGAM)
$\operatorname{Re} z < -6$	580	570
$-6 \leq \operatorname{Re} z < 0$	790	650
$0 \leq \operatorname{Re} z < 7$	630	390
$7 \leq \operatorname{Re} z$	330	300

Unusual features of the program

None.

LONG WRITE-UP

1. INTRODUCTION

Let $z = x + iy$ be a complex variable. The gamma function (for notation see [1])

$$\Gamma(z) = \int_0^{\infty} t^{z-1} e^{-t} dt \quad (x > 0), \quad (1)$$

its logarithm $\ln \Gamma(z)$, and the repeated derivations of $\ln \Gamma(z)$, namely

$$\psi^{(n)}(z) = \frac{d^{n+1}}{dz^{n+1}} \ln \Gamma(z) \quad (n = 1, 2, 3, \dots) \quad (2)$$

play an important role in several fields of mathematics, physics, and other applications. The $\psi^{(n)}(z)$ are often called polygamma functions. In particular, for $n = 1$,

$$\psi(z) = \Gamma'(z)/\Gamma(z) \quad (3)$$

is called the digamma or psi function.

Several methods have been suggested for the computation of these functions, for real or complex values of the variable [2] - [18]. Algol programs for $\Gamma(x)$ with x real are given in [4,], [5], [8], [10]; for $\ln \Gamma(x)$ in [9], [10]; and for $\psi^{(n)}(x)$ in [11]. Rational Chebyshev approximations for $\ln \Gamma(x)$ are given in [14]. For complex values of the variable z , FORTRAN programs for $\Gamma(z)$ have been published in [18] and [19] ^{*)}. In a recent paper, Luke [13] treated the computation of $\Gamma(z)$ for complex z with the help of Padé approximations. He also announced publication of a program for $\Gamma(z)$ using this technique. Wrench [16] and Spira [17] have developed explicit numerical expressions for the coefficients in the Stirling series of $\Gamma(z)$. For the cases where only a limited accuracy is required, Lanczos [6] gave a remarkably simple formula for $\Gamma(z)$, namely

$$\Gamma(z) \approx (z + 1)^{z-\frac{1}{2}} e^{-(z+1)} \sqrt{2\pi} (0.999779 + 1.084635/z) \quad (4)$$

which has a relative error not exceeding 2.4×10^{-4} everywhere in $\text{Re } z > -1, z \neq 0$.

*) This routine occasionally gives wrong results, a correction will be published soon.

A 12-decimal table of $\ln \Gamma(z)$ for $x = 0(0.1)10$, $y = 0(0.1)10$ has been published by the National bureau of Standards [20]. This table also contains a compilation of properties of $\Gamma(z)$ and a bibliography.

An increasing number of applications require the computation of the functions mentioned above for complex arguments. For example, the Veneziano model and its generalizations in high-energy physics [21], [22], [23], which make extensive use of $\Gamma(z)$, or the computation of Coulomb wave functions for complex energies [19].

Although programs for $\Gamma(z)$ are available, it is often desirable to have a program for $\ln \Gamma(z)$. Occasionally, the derivative of the gamma function is also needed. It is therefore useful to present programs for computation of $\ln \Gamma(z)$ and $\psi(z) = \Gamma'(z)/\Gamma(z)$ for complex arguments z .

2. METHODS OF COMPUTATION

2.1 CLOGAM, the program for $\ln \Gamma(z)$

Since $\ln \Gamma(z)$ is an elementary function of $\Gamma(z)$, one might ask why a separate routine is needed if one has a program for $\Gamma(z)$. The answer to this question is twofold: firstly, because the complex logarithm is a multivalued function, defined for $z = x + iy = |z| e^{i(\theta+2n\pi)}$, by

$$\ln z = \ln |z| + i(\theta + 2n\pi) . \quad (5)$$

The standard FORTRAN function CLOG, however, usually computes only the principal value of $\ln z$, i.e. it assumes $n = 0$ and gives θ between $-\pi$ and π . This is not always sufficient if we wish to take the logarithm of a function. For example, we find from a table [20] that for $z = 1 + 5i$, $\text{Im } \ln \Gamma(z) = 3.82 > \pi$, so that the FORTRAN combination $\text{CLOG}(\text{CGAMMA}(Z))$ would certainly not provide the right answer, assuming that CGAMMA computes $\Gamma(z)$. Unfortunately, the authors of [19] did not take this phenomena into account when using their routine GAMMA as part of their program for calculating the Coulomb phase shift. We note here that Luke [13] has developed a method for finding $\text{Im } \Gamma(z)$ if z and $\Gamma(z)$ are given, which can also be applied to $\ln \Gamma(z)$.

The second reason for introducing a separate routine for $\ln \Gamma(z)$ is that this function is much less affected by overflow than $\Gamma(z)$. It can therefore be used for calculating quotients of gamma functions in cases

where both numerator and denominator would cause overflow, but not the quotient itself.

In order to compute $\ln \Gamma(z)$, we use the Stirling series of $\ln \Gamma(z)$ together with suitable functional relations. Other formulae could also be used, for instance the logarithmic equivalent of the relations given by Lanczos [6], which are reproduced in [12]. In contrast to $\Gamma(z)$, where the Stirling series falls off like $1/z$, this series falls off like $1/z^2$ in the case of $\ln \Gamma(z)$. We then have the following formulae [12] in the different regions of the z -plane:

$$\begin{aligned} \ln \Gamma(z) = & (z - \frac{1}{2}) \ln z - z - \frac{1}{2} \ln 2\pi \\ & + z \sum_{k=1}^K \frac{B_{2k}}{2k(2k-1)} z^{-2k} + R_K(z) \quad (x \geq x_0 > 0) \end{aligned} \quad (6)$$

$$\ln \Gamma(z) = \ln \Gamma(z+n) - \ln \prod_{v=0}^{n-1} (z+v) \quad (0 \leq x < x_0)$$

$$\ln \Gamma(z) = \ln \pi - \ln \Gamma(1-z) - \ln \sin \pi z \quad (x < 0),$$

where $n = [x_0] - [x]$. The B_{2k} are the Bernoulli numbers, $[x]$ is the largest integer $\leq x$. Recently, Spira [17] proved a very simple upper bound of the remainder $R_K(z)$ in (6), namely

$$\begin{aligned} |R_K(z)| & \leq |B_{2K}/(2K-1)/z|^{1-2K} \\ & \leq |B_{2K}/(2K-1)/x|^{1-2K} \quad (x \geq 0). \end{aligned} \quad (7)$$

If we choose $K = 10$ and $x_0 = 7$, we find

$$|R_{10}(7)| < 2.5 \times 10^{-15}$$

so that theoretically these values are suitable for single precision computation on a CDC 6000 computer. When computing the logarithm of the product in formula (6), we must be careful to find the correct imaginary part. This can be achieved by using the relation

$$\ln \prod_{v=0}^{n-1} (z + v) = \frac{1}{2} \ln \left| \prod_{v=0}^{n-1} (z + v) \right| + i \sum_{v=0}^{n-1} \arctan \frac{y}{x + v}. \quad (8)$$

The function $f(z) = \ln \sin \pi z$ in formula (6) is computed in the following way. We set $z = x + iy$ and assume $y \geq 0$. This is no restriction since $\ln \Gamma(\bar{z}) = \overline{\ln \Gamma(z)}$, where $\bar{z} = x - iy$. We introduce a new variable $\xi = x - [x]$ so that $0 \leq \xi < 1$, and obtain

$$f(z) = \frac{1}{2} \ln (\sin^2 \pi \xi + \sinh^2 \pi y) + i \{ \arctan (\cot \pi \xi \tanh \pi y) - [x] \pi \}, \quad (9)$$

where $-\pi < \arctan \omega \leq \pi$. In particular, on the boundaries of the cut along the negative real axis ($y = \pm 0$), we have

$$f(x) = \ln |\sin \pi \xi| \pm [x] \pi i = \ln |\sin \pi \xi| \mp |[x]| \pi i. \quad (10)$$

Finally, in order to avoid possible overflow, we write $\operatorname{Re} f(z)$ in the form

$$\operatorname{Re} f(z) = \pi y + \frac{1}{2} \ln \left[e^{-2\pi y} \sin^2 \pi \xi + \frac{1}{4} (1 - e^{-2\pi y})^2 \right]. \quad (11)$$

2.2 CDIGAM, the program for $\psi(z)$ *)

In order to compute $\psi(z) = d \ln \Gamma(z)/dz$ we differentiate (6), giving

$$\psi(z) = \ln z - \frac{1}{2z} - \sum_{k=1}^K \frac{B_{2k}}{2k} z^{-2k} + R'_K(z) \quad (x \geq x_0 > 0)$$

$$\psi(z) = \psi(z + n) - \sum_{v=0}^{n-1} (z + v)^{-1} \quad (0 \leq x < x_0)$$

$$\psi(z) = \psi(-z) + 1/z + \pi \cot \pi z \quad (x < 0),$$

where $n = [x_0] - [x]$. These formulas are used for the computation of $\psi(z)$ in the regions indicated. We have chosen $x_0 = 7$ and $K = 7$.

*) An earlier version of this program was written by R. Keyser (CERN).

3. THE TESTS

3.1 Tests for $\ln \Gamma(z)$

The program CLOGAM was checked in the following cases.

- i) The values of $\ln \Gamma(z)$ were computed for $z = x + iy$, where $x = 0(0.5)1(10)$, $y = 0(0.1)10$. The results were compared with the 12-decimal table [23]. Occasional discrepancies of 1 unit in the 12th decimal were found.

- ii) The formula [24]

$$\ln \Gamma(2z) = (2z - 1) \ln 2 + \ln \Gamma(z) + \ln \Gamma(z + \frac{1}{2}) - \frac{1}{2} \ln \pi, \quad (13)$$

was "verified" using 500 arguments of the form

$$z = \mu [x - \frac{1}{2} + i(y - \frac{1}{2})],$$

where x and y were random numbers uniformly distributed over $(0,1)$.

The proportionality factor μ was taken to be $\mu = 2, 5, 10, 30, 50$.

For each μ , 100 values z were taken. The two sides of (13) usually agreed to 12-14S (significant digits), occasionally to 11S.

- iii) Calculation of $\ln \Gamma(z)$ for $z = x + iy$, where $y = -3.0(0.1)3.0$, $y = \pm 5, \pm 0.01, \pm 0.00001, 0$. This verifies that the imaginary part is computed correctly.

- iv) Computation of $\ln \Gamma(z)$ for

$$z = \left[(-1)^j x_k + i (-1)^\ell y_k \right] \times 10^m,$$

where $k = 1, 2, 3$; $j = 2, 3$; $\ell = 2, 3$; $m = 1, 2$; $x_k = \{0.1, 0.1, 1\}$; $y_k = \{1, 1, 0.2\}$. The computed values were checked against the values found by direct evaluation of the Stirling series. In particular, for $\text{Re } z < 0$, this check is important as an assurance that the sign of the imaginary part of $\ln \Gamma(z)$ is correct.

3.2 Tests for $\psi(z)$

We have tested the following cases using the program CDIGAM.

- i) The formula

$$\psi(2z) = \frac{1}{2} [\psi(z) + \psi(z + \frac{1}{2})] + \ln 2$$

was checked for the same values of z as formula (13). The two sides agree to 13-14S.

ii) The relations

$$\operatorname{Im} \psi(iy) = 1/(2y) + \frac{1}{2}\pi \coth \pi y$$

$$\operatorname{Im} \psi(1 + iy) = -1/(2y) + \frac{1}{2}\pi \coth \pi y$$

were checked for $y = 0.1(0.1)1(1)100$. There was agreement to 13-14S.

iii) Direct evaluation of [24]

$$\psi(z) = \ln z - 1/(2z) - \int_0^{\infty} \frac{t \, dt}{(t^2 + z^2)(e^{2\pi t} - 1)} \quad (\operatorname{Re} z > 0)$$

for 50 values of z with $0 < \operatorname{Re} z \leq 10$, $0 < \operatorname{Im} z \leq 10$. The upper limit of the integral was replaced by $T = (10 \ln 10)/\pi \approx 7.33$.

There was agreement to 13-14S.

We note here that relative accuracy is necessarily lost near a zero of $\psi(z)$. These zeros are all negative real, except $x_0 = 1.46163\dots$. For arguments z with large negative real part $\operatorname{Re} z \approx -10^n$, about $(14-n)$ significant digits are correct.

4. ERROR EXISTS

4.1 CLOGAM

If the function subprogram CLOGAM is called with an argument $z = -n \pm i0$, ($n = 0, 1, 2, \dots$) an error message

CLOGAM ... ARGUMENT IS NON-POSITIVE INTEGER = (n)

is printed on Logical Unit 2, where (n) denotes the argument. The value of CLOGAM is set to zero in this case.

4.2 CDIGAM

The function subprogram CDIGAM, when called with an argument $z = -n \pm i0$ ($n = 0, 1, 2, \dots$), prints an error message

CDIGAM ... ARGUMENT IS NON-POSITIVE INTEGER = (n)

The value of CDIGAM is set to zero in this case.

5. TEST RUN

The results of the test run are self-explanatory.

Acknowledgement

I wish to thank G.A. Erskine for comments and discussions.

TEST RUN OUTPUT

TEST VALUES FOR LN GAMMA(Z) (Z = X + I*Y)

X	Y	RE LN GAMMA(Z)	IM LN GAMMA(Z)
-13.0	2.0	-26.84992384156715	-37.19840614844497
-9.5	0.0	-12.79589533355426	-31.41592653589794
-3.0	1.0	-2.95350829229571	-9.72641828123693
0.0	8.0	-12.68715285199448	7.83971205351668
3.0	0.0	.69314718055986	0.00000000000000
4.0	2.0	1.25083561935671	2.61019580104885
7.0	4.0	5.41808697187290	7.71810136520472
9.0	16.0	-.27799290829569	39.55316531442281
10.0	0.0	12.80182748008133	0.00000000000000
15.0	5.0	24.34577701569333	13.46736924371726
-13.0	-2.0	-26.84992384156715	37.19840614844497
-9.5	-0.0	-12.79589533355426	31.41592653589794
-3.0	-1.0	-2.95350829229571	9.72641828123693
0.0	-8.0	-12.68715285199448	-7.83971205351668
3.0	-0.0	.69314718055986	-0.00000000000000
4.0	-2.0	1.25083561935671	-2.61019580104885
7.0	-4.0	5.41808697187290	-7.71810136520472
9.0	-16.0	-.27799290829569	-39.55316531442281
10.0	-0.0	12.80182748008133	-0.00000000000000
15.0	-5.0	24.34577701569333	-13.46736924371726

CLOGAM ... ARGUMENT IS NON-POSITIVE INTEGER = -1.00

TEST VALUES FOR PSI(Z) = DIGAMMA(Z)

X	Y	RE PSI(Z)	IM PSI(Z)
-13.0	2.0	2.61375885861489	2.99460095564282
-9.5	0.0	2.30300103429782	0.00000000000000
-3.0	1.0	1.29465032062246	2.87667404746858
0.0	8.0	2.08074567491178	1.63329632679488
3.0	0.0	.92278433509845	0.00000000000000
4.0	2.0	1.39536074614320	.51696112879607
7.0	4.0	2.03269565223019	.55101815665321
9.0	16.0	2.89681672499673	1.08235712929482
10.0	0.0	2.25175258906671	0.00000000000000
15.0	5.0	2.73046382968629	.33195042663378
-13.0	-2.0	2.61375885861489	-2.99460095564282
-9.5	-0.0	2.30300103429782	0.00000000000000
-3.0	-1.0	1.29465032062246	-2.87667404746855
0.0	-8.0	2.08074567491178	-1.63329632679488
3.0	-0.0	.92278433509845	0.00000000000000
4.0	-2.0	1.39536074614320	-.51696112879607
7.0	-4.0	2.03269565223019	-.55101815665321
9.0	-16.0	2.89681672499673	-1.08235712929482
10.0	-0.0	2.25175258906671	0.00000000000000
15.0	-5.0	2.73046382968629	-.33195042663378

CDIGAM ... ARGUMENT IS NON-POSITIVE INTEGER = -1.00

REFERENCES

- [1] M. Abramowitz and I.A. Stegun (Editors), Handbook of Mathematical Functions, with Formulas, Graphs and Mathematical Tables, Nat. Bur. Standards Appl. Math. Ser. 55 (U.S. Government Printing Office, Washington, D.C., 1965).
- [2] H. Werner and R. Collinge, Chebyshev approximations to the gamma function, Math. Comp. 15 (1961) 195-197.
- [3] C.E. Fröberg, Rational Chebyshev approximation of elementary functions, Nordisk Tidskr. Informationsbehandling (BIT) 1 (1961) 256-262.
- [4] P. Wynn, An arsenal of Algol procedures for complex arithmetic, Nordisk Tidskr. Informationsbehandling (BIT) 2 (1962) 232-255.
- [5] C.W. Clenshaw, G.F. Miller and M. Woodger, Algorithms for special functions I, Handbook Series Special Functions, Numer. Math. 4 (1963) 403-419.
- [6] C. Lanczos, A precision approximation of the gamma function, J. SIAM Numer. Anal., Ser. B, 1 (1964) 86-96.
- [7] J.R. Rice, On the L_{∞} Walsh arrays for $\Gamma(x)$ and $\text{Erfc}(x)$, Math. Comp. 18 (1964) 617-626.
- [8] S.J. Cyvin and B.N. Cyvin, Algorithm 225, Gamma function with controlled accuracy, Comm. ACM 7 (1964) 295.
- [9] M.C. Pike and I.D. Hill, Algorithm 291, Logarithm of Gamma function, Comm. ACM 9 (1966) 684.
- [10] A. Machado Souza Filho and G. Schwachheim, Algorithm 309, Gamma function with arbitrary precision, Comm. ACM 10 (1967) 511-512.
- [11] A. Tadeu de Medeiros and G. Schwachheim, Algorithm 349, Polygamma functions with arbitrary precision, Comm. ACM 12 (1969) 213-214.
- [12] Y.L. Luke, The special functions and their approximations, Vol. I and II (Academic Press, New York and London, 1969).
- [13] Y.L. Luke, Evaluation of the gamma function by means of Padé approximations, SIAM J. Math. Anal. 1 (1970) 266-281.
- [14] W.J. Cody and K.E. Hillstrom, Chebyshev approximations for the natural logarithm of the gamma function, Math. Comp. 21 (1967) 198-203.
- [15] J.F. Hart, E.W. Cheney, C.L. Lawson, H.J. Maehly, C.K. Mesztenyi, J.R. Rice, H.C. Thacher and C. Witzgall, Computer Approximations (Wiley and Sons, Inc., New York, 1968).
- [16] J.W. Wrench, Jr., Concerning two series of the gamma functions, Math. Comp. 22 (1968) 617-626.
- [17] R. Spira, Calculation of the gamma functions by Stirlings formula, Math. Comp. 25 (1971) 317-322.
- [18] C.W. Lucas, Jr., and C.W. Terrill, Algorithm 404, Complex gamma function, Comm. ACM 14 (1971) 48-49.

- [19] T. Tamura and F. Rybicki, Coulomb functions for complex energies, Computer Phys. Comm. 1 (1969) 106-112.
- [20] H.E. Salzer, Table of the gamma function for complex arguments, Nat. Bur. Standards Appl. Math. Ser. 34 (U.S. Government Printing Office, Washington, D.C., 1954).
- [21] C. Lovelace, Veneziano theory, Proc. Roy. Soc. Lond. A318 (1970) 321-354.
- [22] Chan Hong-Mo, The generalized Veneziano model, Proc. Roy. Soc. Lond. A318 (1970) 379-399.
- [23] J.F.L. Hopkinson and E. Plahte, Infinite series representation of the N-point function in the generalized Veneziano model, Phys. Letters 28 B (1969) 489-492.
- [24] I.M. Ryshik and I.S. Gradstein, Tafeln-Tables (VEB Deutscher Verlag d. Wissensch., Berlin, 1957).