CM-P00056989

ON THE THEORY OF FERMI INTERACTIONSG. Domokos ^{*)}

CERN - Geneva

A B S T R A C T

The Bethe-Salpeter equation for two fermions interacting through a V-A four-fermion interaction is investigated: the kernel is approximated by a single fermion bubble. The Green function has essential singularity at small relative distances; it is free of arbitrary subtraction constants; at low energies the scattering amplitude tends to its lowest order expression; coupling constant is renormalized by a finite factor. The zero energy result of peratization theory is reproduced to leading order in the coupling constant for vector meson exchange. Generality and possible importance of the results for field theory are discussed.

---

*) On leave from the Joint Institute for Nuclear Research, Dubna, USSR

Permanent address : Central Research
Institute for Physics, Budapest

1. INTRODUCTION

In a recent paper ¹⁾ we started an investigation of the theory of non-renormalizable interactions. The method we used was to calculate the two-particle Green function from a Bethe-Salpeter (BS) equation with a local kernel ²⁾.

In particular, we have calculated the scattering of leptons in the multiperipheral approximation. We found that the scattering amplitude has essential singularities both in energy and momentum transfer, however, in the physical region it decreases with the energy. An interesting feature of our results was that - due to the essential singularity of the Green function at small relative distances - the contribution of subtraction terms vanished identically ; thus, the Green function and the scattering amplitude did not contain any arbitrary constants.

The aim of the present paper is to investigate this point further.

The case we consider is still the uncrossed ladder approximation with a local kernel, however, for a diagram which, in our opinion, resembles reality more closely. In fact, by considering our diagram, we are led to consider a two-channel problem. Decoupling the equations, we find two BS equations of identical structure, the sign of the interaction being only different. In fact, our system of equations is of the same structure as that considered by Feinberg and Pais ³⁾ as a starting point of their peratization procedure and by Sawyer ⁴⁾. Our procedure to be followed is rather close to Sawyer's one and in fact it has been applied with minor modifications already several times by different authors to investigate the qualitative properties of strong interactions ⁵⁾.

In order to facilitate the reading of the present paper, in section 2 we briefly review the method to be applied and we give the formulae necessary in the following.

The model to be investigated is described in section 3, while sections 4 and 5 are devoted to its investigation. In section 6 we consider briefly the W meson ladder of Feinberg and Pais ³⁾. At zero energy and to leading order in the coupling constant the result of Feinberg and Pais is reproduced ; this may serve as a cross-check of both methods. Finally, in section 7, we discuss the results obtained ; the same paragraph contains some speculations concerning their generality and relevance to field theory.

Some parts of this paper overlap with a recent work of P. Dennery and the present author on higher order corrections to weak interactions. Nevertheless, we decided to make both papers self-contained even at the cost of some repetitions.

2. REVIEW OF THE TREATMENT OF BETHE-SALPETER EQUATIONS
WITH LOCAL KERNEL

The BS equation for the two-particle Green function G can be written in symbolic form as follows :

$$G = G_0 + G_0 K G \quad (2.1)$$

where G_0 is the two-particle Green function in the absence of interaction, K - the BS kernel - is usually defined as the sum of contributions of "doubly connected diagrams", i.e., those which cannot be disconnected by cutting two lines of the propagating particle. For our purpose it is important that Eq. (2.1) has a meaning independently of perturbation theory, as emphasized, e.g., by Symanzik ⁶⁾. In fact we shall find that the solution we obtain is in general singular in the coupling constant so a perturbation expansion necessarily diverges. Equation (2.1) can be transcribed into several forms. The one we shall use is obtained by multiplying it by G_0^{-1} from the left ; another one is that used by Feinberg and Pais ³⁾ and Sawyer ⁴⁾. It is obtained by introducing the transition amplitude (T) off the mass shell by the definition

$$G = G_0 + G_0 T G_0 \quad (2.2)$$

which yields

$$T = K + K G_0 T \quad (2.3)$$

Equations (2.1) and (2.3) are equivalent. G or T are in general spinor matrices ; by well-known procedures one can decompose them to obtain a system of equations for the invariant functions. In what follows, we assume that such a decomposition with respect to invariants has been carried out and the resulting

system decoupled. Writing out Eq. (2.1) in momentum representation, we obtain

$$G(p, p' | E) = G_0(p, p' | E) + \int dq' dq'' G_0(p, q' | E) K(q', q'' | E) \times G(q'', p' | E) \quad (2.4)$$

where $p, \dots, p'$ denote the relative momenta of the propagating particles and E their total four momentum.

We write

$$G_0(p, p' | E) = \frac{\delta(p - p')}{F(p, E)}$$

$\frac{1}{F(p, E)}$ being the product of two free one-particle propagators. Assuming that $K(q', q'' | E)$ depends on $q' - q''$ only, (this is our definition of a local kernel), we multiply (2.4) by G_0^{-1} , go over to Euclidean relative momenta ⁷⁾ and expand the equation with respect to hyperspherical harmonics. In the notation used in our previous papers, the latter are :

$$Z_{nl}^m(\psi, \vartheta, \varphi) = \gamma_{nl}(\sin \psi)^l C_{n-l-1}^{l+1}(\cos \psi) \times Y_l^m(\vartheta, \varphi)$$

where the Y_l^m are the usual normalized spherical harmonics, C_{n-l-1}^{l+1} is a Gegenbauer polynomial,

$$\gamma_{nl} = 2^{l+1/2} \left(\frac{n \Gamma(n-l)}{\pi \Gamma(n+l+1)} \right)^{1/2} \Gamma(l+1)$$

We define the quantum number n - the four dimensional angular momentum - such that it ranges from 1 to ∞ through integers. (This is convenient because the resulting radial equations closely resemble ordinary Schrödinger equations.)

In this representation our equation reads

$$\begin{aligned} \sum_{n''} \langle n | F(p^2, E) | n'' \rangle \langle p^2 n'' | G(E) | p'^2 n' \rangle = \\ = \frac{\delta(|p| - |p'|)}{(|p| |p'|)^{3/2}} + \\ + \int_0^\infty q^3 dq \langle p^2 | K(n, E) | q^2 \rangle \langle q^2 | G(E) | p'^2 n' \rangle \end{aligned} \quad (2.5)$$

The notation in (2.5) is self-explaining ; due to the high symmetry of the problem the matrix elements do not depend on l, m ; diagonal labels are written as arguments. The kinematic part $\langle n | F(p^2, E) | n'' \rangle$ for $E_\mu \neq 0$ contains off diagonal elements, which vanish for zero total four momentum. Instead of putting $E_\mu = 0$ we consider the off-diagonal elements of F as perturbations, which can be neglected in the first approximation⁸⁾. Doing so, the whole equation (2.5) becomes diagonal in n . In what follows, we shall work in this "diagonal approximation" only.

If $K(p-q|E)$ possesses a spectral representation in $(q^2 - q)^2$, in the $(np^2|E)$ representation it has the form

$$\begin{aligned} \langle p^2 | K(n, E) | q^2 \rangle = \\ = \int_0^\infty \kappa^3 d\kappa \frac{J_n(p\kappa)}{p\kappa} V(\kappa) \frac{J_n(q\kappa)}{q\kappa} \end{aligned} \quad (2.6)$$

with

$$V(r) = \int_0^\infty \mu^3 d\mu \sigma(\mu) \frac{K_1(\mu r)}{\mu r} \quad (2.7)$$

where $\sigma(\mu)$ is the spectral density of the kernel. Thus a Hankel transformation applied to Eq. (2.5) gives a local differential equation in the radial co-ordinates :

$$D_n(r, E) G(r, r') = \frac{\delta(r - r')}{(rr')^{3/2}} + V(r) G(r, r') \quad (2.8)$$

In the last formula $D_n(r, E)$ means the Hankel transform of $\langle n | F | n \rangle$; it is a local differential operator. We omitted the labels n, E from G . It is important to emphasize that Eq. (2.8) as it stands is valid for $\text{Re } n > N$ only, where N is the number of subtractions necessary in the spectral representation of K . However, as explained in I, we can continue the radial equation (2.8) for $\text{Re } n > N$. Care is needed in the subtracted partial waves only ; here the total Green function is given by the equation (written in a symbolic form)

$$G = G_1 + G_1 \bar{K} G \quad (2.9)$$

where G_1 is the solution of (2.8) while $\bar{K}$ is the projection of the subtraction polynomial onto the partial wave considered.

As Eq. (2.8) is an ordinary differential equation, the Green function $G(r, r')$ can be constructed from the solutions of the homogeneous equation

$$D_n(r, E) \psi(r) = V(r) \psi(r) \quad (2.10)$$

The boundary conditions to be chosen are similar to those in non-relativistic quantum mechanics ; in fact, with the help of $\psi_n(r)$ the off-the-shell scattering amplitude is expressed by the familiar looking formula

$$\langle p' | T_n | p \rangle = \frac{16\pi^2}{p} \int_0^\infty r'^3 dr' \frac{f_n(p'r')}{p'r'} V(r') \psi_n(r') \quad (2.11)$$

The existence of T_n requires to choose a solution which is finite - or vanishes - for $r = 0$. At $r \rightarrow \infty$ the boundary condition is $\psi(r) \sim \exp(-ipr) + S_n \exp(ipr)$ or the corresponding standing wave solution ^{1),4),5)}. S_n is - formally - unitary : $S_n = \exp 2i\delta_n$, where δ_n is real if $V(r)$ is real ; this relation, however, is not necessarily identical to physical unitarity : the two-particle unitarity condition

$$\text{Im} T(E^2, \cos \theta) = \rho(E^2) \int d\Omega'' T(E^2, \cos \theta') T^*(E^2, \cos \theta'')$$

is not diagonal in the basis we use. (The reason of this apparent paradox will be explained later.)

Having found the Green function for the analytic part of the interaction [from Eq. (2.10)], the subtraction equations (2.9) can readily be solved. In fact, the subtraction polynomial, $\bar{K}$ represents a superposition of contact interactions ; after simple manipulations then we arrive at the expression for the complete scattering amplitude ¹⁾, $\bar{T}_n$:

$$\langle p | \bar{T}_n | p' \rangle = \langle p | T_n | p' \rangle + \frac{F_n(p) \bar{K}_n F_n^*(p')}{1 - \bar{K}_n \phi_n} \quad (2.12)$$

8.

where $\langle p | T_n | p' \rangle$ is the amplitude defined by (2.11)

$$F_n(p) = G_0(p)^{-1} \int_0^\infty r^3 dr \frac{f_n(pr)}{pr} G_n(r, 0)$$

$$\phi_n = G_n(0, 0)$$

We see that $\bar{T} = T$ except for partial waves where $\bar{K}_n$ differs from zero. If G can be expanded in the coupling constant, to lowest order we find

$$\langle p | \bar{T}_n | p' \rangle \approx \langle p | T_n | p' \rangle + \bar{K}_n \quad (2.13)$$

After this preparation we turn to the formulation of our model.

3. DESCRIPTION OF THE MODEL

We consider the interaction of massless, but distinguishable fermions ; we call them for the sake of definiteness μ , ν_μ and e , ν_e respectively. They interact via an interaction of the V-A type with charged currents. Except in section 5, we do not assume the existence of elementary W mesons.

We consider both "allowed" transitions, e.g., the reaction

$$\nu_\mu + e \rightarrow \nu_e + \mu \quad (3.1)$$

and "forbidden" ones, e.g.,

$$\nu_\mu + e \rightarrow \nu_\mu + e \quad (3.2)$$

The basic process we are going to consider is an allowed reaction of the type (3.1) taking place through the exchange of a fermion loop (Fig. 1)

Fig. 1

As we neglect fermion masses, the contributions of all the diagrams are equal and we need not specify the fermion loop exchanged ($\mu + \bar{\nu}_\mu$ or $e + \bar{\nu}_e$), but multiply the contribution of each diagram on Fig. 1 by two.

Taking into account time reversal invariance and the conservation of charge and muon number, we are thus left with a two-channel problem, the channels being represented by, e.g., $(\nu_\mu e)$ and $(\nu_e \mu)$.

Equation (2.1) written in matrix form in the channel space thus reads

$$\begin{pmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{pmatrix} = \begin{pmatrix} G_0 & 0 \\ 0 & G_0 \end{pmatrix} + \begin{pmatrix} 0 & G_0 K \\ G_0 K & 0 \end{pmatrix} \begin{pmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{pmatrix} \quad (3.3)$$

($G_{12} = G_{21}$ because of T invariance and $G_{11} = G_{22}$ because of our model). This equation can be diagonalized by the orthogonal transformation

$$U = 2^{-1/2} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$$

whereupon it takes the form

$$G^\pm = G_0 \pm G_0 K G^\pm \quad (3.4)$$

with

$$G^\pm = G_{11} \pm G_{12} \quad (3.5)$$

The reader will immediately recognize that the selection rules and amplitudes of our model correspond to those of Feinberg and Pais³⁾, G_{11} , G_{12} corresponding to their "even" and "odd" diagrams.

The multichannel formulation ⁹⁾ without recourse to diagrams is essential in our case, as we shall find that the final solution is not expandable in powers of the coupling constant.

The spinorial structure of our equation is essentially the same as in Ref. ³⁾; after multiplication by G_0^{-1} from the left, Eq. (3.4) reads in momentum representation

$$\begin{aligned} & \left(\frac{1}{2} \hat{E}^{(1)} + \hat{p}^{(1)} \right) \left(\frac{1}{2} \hat{E}^{(2)} - \hat{p}^{(2)} \right) \gamma_\mu^{(1)} (1 + \gamma_5^{(1)}) \gamma_\nu^{(2)} (1 + \gamma_5^{(2)}) G_{\mu\nu}^\pm(p, p') \\ &= \delta(p - p') \pm \frac{1}{(2\pi)^4 i} \int d^4 q \gamma_5^{(1)} (1 - \gamma_5^{(1)}) \gamma_\sigma^{(2)} (1 - \gamma_5^{(2)}) K_{\rho\sigma}(p, q) \times \\ & \times \gamma_\alpha^{(1)} (1 + \gamma_5^{(1)}) \gamma_\beta^{(2)} (1 + \gamma_5^{(2)}) G_{\alpha\beta}^\pm(q, p'), \end{aligned} \quad (3.6)$$

where the superscripts 1 and 2 refer to the two propagating particles, E is the total four momentum, p, p', q are relative momenta in the final - initial - and intermediate states respectively.

The tensor $G_{\alpha\beta}$ is a linear combination of the tensors $g_{\mu\nu}$, $E_\alpha E_\beta$, $p_\alpha p_\beta$, ... etc., multiplied by invariant functions. In order to get rid of the difficulties connected with coupled equations between invariant functions, we now make the drastic assumption that all the invariant functions are zero except the coefficient of $g_{\mu\nu}$, i.e., we set

$$G_{\alpha\beta} = g_{\alpha\beta} G \quad (3.7)$$

This assumption greatly simplifies the structure of our equations ; intuitively one expects that it may not be too bad for the physical scattering amplitude at low energies ; there does not seem to exist any a priori justification for it for the Green function. We nevertheless make it, because we are now primarily interested in qualitative features of the theory, and we feel that in that respect nothing essential is lost by assuming (3.7).

Inserting (3.7) into (3.6) and taking the trace of the equation, we obtain

$$\begin{aligned} (1/4 E^2 - p^2) G^\pm(p, p') &= \delta(p - p') \pm \\ &\pm \frac{1}{(2\pi)^4 i} \int dq K(p, q) G^\pm(q, p') \end{aligned} \quad (3.8)$$

where $K(p, q) = K_{\epsilon^S}(p, q)$.

K_{ϵ^S} can be calculated in a straightforward way (cf. Fig. 1) : it essentially corresponds to a self-energy diagram in electrodynamics with massless electrons. Omitting the details of the calculation, we only notice that Eq. (3.8), written in Euclidean metric, is diagonal in the $(n|m)$ representation. Expanding $G^\pm$ and K according to the functions

$$\sum_{n \in \mathbb{Z}}^m \frac{J_n(p\epsilon)}{p\epsilon}$$

we arrive at the radial equation in co-ordinate space

$$\left(\frac{d^2}{d\epsilon^2} + \frac{3}{\epsilon} \frac{d}{d\epsilon} + k^2 - \frac{n^2 - 1}{\epsilon^2} \mp V(\epsilon) \right) G^\pm(\epsilon, \epsilon') = \frac{\delta(\epsilon - \epsilon')}{(\epsilon \epsilon')^{3/2}} \quad (3.9)$$

with $k^2 = 1/4 E^2$,

$$V(\tau) = \frac{2f^2}{\pi\tau^6} \int_0^\infty x^4 dx K_1(x) \equiv \lambda^2 \tau^{-6}$$

where $\lambda^2 = 32 \pi^{-1} f^2$. The spectral representation of K needs two subtractions (the fermion loop is quadratically divergent) so the subtracted waves are $n = 1, 2$. (The calculations leading to Eq. (3.9) are described in detail in Ref. ¹⁾). Equation (3.9) is the starting point of our further investigations.

4. INVESTIGATION OF THE GREEN FUNCTION

First of all we set $G^{\pm}(r, r') = (rr')^{-3/2} \Gamma^{\pm}(r, r')$, thereby eliminating the first derivative from Eq. (3.9). According to the procedure sketched in section 2, we write the equation for the radial wave function

$$\left(\frac{d^2}{ds^2} + k^2 - \frac{n^2 - 1/4}{s^2} \mp \frac{\lambda^2}{s^6} \right) u^{\pm}(s) = 0 \quad (4.1)$$

Equation (4.1) is exactly of the form of a non-relativistic Schrödinger equation with a singular potential, $\ell + 1/2$ being now replaced by n , supplemented with boundary conditions which are also the same as in non-relativistic quantum mechanics. In investigating Eq. (4.1) we have to treat the "repulsive" and "attractive" potentials separately.

As far as we know, Eq. (4.1) cannot be integrated in terms of known transcendental functions. We begin by noting that for $k^2 = 0$ the equation can be integrated exactly ^{1), 4)}.

Two independent solutions are

$$u_0^+(s) = \begin{cases} s^{1/2} I_{n/2} \left(\frac{\lambda}{2s^2} \right) \\ s^{1/2} K_{n/2} \left(\frac{\lambda}{2s^2} \right) \end{cases} \quad (4.2)$$

and

$$u_0^-(s) = \begin{cases} s^{1/2} H_{n/2}^{(1)} \left(\frac{\lambda}{2s^2} \right) \\ s^{1/2} H_{n/2}^{(2)} \left(\frac{\lambda}{2s^2} \right) \end{cases} \quad (4.3)$$

For u_0^+ it is obviously only $K_{n/2}$ which satisfies the boundary condition at the origin, while for u_0^- both solutions behave in the same way, they only differ by a phase. In order to proceed further, we set $k^2 > 0$, and calculate $u^\pm(r)$ in the WKB approximation.

a) Repulsive case

There is always one classical turning point. Its position (b) can be easily found as a series in descending powers of k :

$$b = \left(\frac{\lambda}{k}\right)^{1/3} \left[1 + \frac{n^2}{6\lambda^2} \left(\frac{\lambda}{k}\right)^{4/3} + \dots \right] \quad (4.4)$$

The singularity of the potential manifests itself in (4.4) : b is essentially determined by the interaction and does not approach its "free value" - n/k - as $k \rightarrow \infty$. Setting

$$u^+(\kappa) = A(\kappa) \exp i S(\kappa) \\ S(\kappa) = \int_b^\kappa d\xi \left(k^2 - n^2 \xi^{-2} - \lambda^2 \xi^{-6} \right)^{1/2} \quad (4.5)$$

$A(r)$ satisfies the equation

$$A'' + 2iA'S' + iAS'' = 0$$

and dropping A'' , in the first approximation we obtain the standard expression

$$A_1(\kappa) = \left(S'(\kappa) \right)^{-1/2}$$

(Dashes mean differentiation with respect to r .)

Treating A'' as a perturbation, we obtain in the next approximation

$$A_2 = -\frac{i}{4}(S')^{-1/2} \log S'$$

and so on. This calculation shows explicitly that something unexpected does not happen in comparison with the zero energy solution ; one can always choose a solution which vanishes at the origin ; the solution is determined uniquely and in the leading order it behaves as $\exp(-\lambda 2^{-1/2} r^2)$ near the origin. The treatment of the transition region is likewise standard ¹⁰⁾.

b) Attractive case

The effective potential : $U(r) = n^2 r^{-2} - \lambda^2 r^{-6}$ is not monotonous ; it has a maximum at $r_0 = 3^{1/2} n^{-2} \lambda^2$. There are either two real turning points or none, depending on the magnitude of k^2 . To find their positions, we approximate $U(r)$ by a parabola near its maximum :

$$U(r) \approx \frac{3^{-1} 2 n^3}{\lambda} \left[3^{-1/2} - \frac{2n}{\lambda} (r - r_0)^2 \right]$$

This gives for the turning points

$$b_{1,2} = r_0 \pm \frac{\lambda}{2n} \left(3^{-1/2} - \frac{3\lambda k^2}{2n^3} \right)$$

We see that there are real turning points, if

$$k^2 < \frac{2n^3}{3^{1/2}\lambda} \equiv k_0^2 \quad (4.6)$$

If k^2 is less than the limit given by Eq. (4.6), the solution is unique ; there are two real turning points ; under the barrier ($b_2 < r < b_1$) the solution is given in terms of functions of the parabolic cylinder, while for $r < b_2$, $r > b_1$ the solution is given by linear combinations of the functions

$$u^-(r) \sim (S'(r))^{-1/2} \exp \pm i \int_{b_{1,2}}^r p(\xi) d\xi,$$

$$p(\xi) = (k^2 - n^2 \xi^{-2} + \lambda^2 \xi^{-6})^{1/2}$$

The behaviour for $r \rightarrow 0$ to leading order is

$$u^-(r) \sim \exp(\pm i \lambda 2^{-1} r^{-2})$$

in agreement with the zero energy solution (4.3). The situation is different for $k^2 > k_0^2$. (Although we do not believe that the ladder diagram we are considering here gives an adequate description of the scattering amplitude at such extremely high energies, nevertheless, it is instructive to investigate this case from a purely theoretical point of view.) If $k^2 > k_0^2$, there is no real turning point at all, the barrier is too low to prevent the system from "collapsing". In order to get closer to the usual procedure applied in field theory, we rewrite (4.6) as

$$n > 3^{1/6} \lambda^{1/3} (k^2/2)^{1/3} \quad (4.6')$$

For a fixed energy, the partial waves satisfying the inequality (4.6') are well defined. (For $k^2 < k_0^2$ all the physical partial waves are such.) As k^2 increases, the first, second, ... partial waves become divergent.

In fact, let us try to write down the expression for $u^-(r)$ for the "supercritical" domain. Introducing a cut-off ρ at the lower limit in the expression of $S(r)$, we have to leading order

$$u^-(r) \sim \lim_{\xi \rightarrow 0} \exp \pm i \int_{\xi}^r d\xi p(\xi)$$

or for $\xi^2 < r^2 \ll \lambda$,

$$u^-(r) \sim \lim_{\xi \rightarrow 0} \exp \pm i \lambda \int_{\xi}^r \frac{d\xi}{\xi^3}$$

Had we to do with a renormalizable theory, one would begin to separate the divergent part of $u^-(r)$ by one of the usual methods, e.g., by successive differentiation with respect to the external variable ¹¹⁾. It is immediately clear that in this way one arrives at an infinite number of arbitrary subtraction constants - which is clearly nonsense. We propose another way of getting rid of the infinities. One can namely make one subtraction in the exponential, by writing

$$S(r) = \int_0^r d\xi \left\{ (k^2 - n^2 \xi^{-2} + \lambda^2 \xi^{-6})^{1/2} - \lambda \xi^{-3} \right\} - \frac{\lambda}{2r^2} + S_0 \quad (4.7)$$

where S_0 is a real constant, independent of n . The subtraction constant S_0 affects but the phase of the wave functions $u^-(r)$, and it is common to all partial waves. It can be removed by a suitable redefinition of $u^-(r)$, having no effect on any physical consequence of the theory. In order to explore a bit further this result, let us expand the curly bracket in the expression (4.7) for small values of r (corresponding to large relative momenta in momentum representation). A simple calculation gives

$$S(r) = S_0 - \frac{\lambda}{2r^2} + \frac{r^2}{2\lambda} \left(\frac{k^2 r^2}{4} - n^2 \right) - \frac{r^6}{4\lambda^3} \left(\frac{k^4 r^4}{5} - \frac{k^2 r^2 n^2}{2} + \frac{n^4}{3} \right) + \dots \quad (4.8)$$

i.e., one obtains an expansion in descending powers of the coupling constant.

Obviously, such a result cannot be produced by a straightforward application of perturbation theory, as there is a singularity at $\lambda^2 = 0$. In order to clarify the nature of the latter, we put $\lambda \xi^{-2} = y$; this gives

$$S(r) = \frac{1}{2} \int_{\lambda r^{-2}}^{\infty} dy \left\{ \left[\frac{k^2 \lambda}{y^3} - \frac{n^2}{y^2} + 1 \right]^{1/2} - 1 \right\}$$

Hence we can see the singularity at $\lambda^2 = 0$:

$$S(r) \sim \frac{in}{2} \log \lambda / r \quad (\lambda \rightarrow 0) \quad (4.9)$$

or equivalently

$$u^-(r) = O\left(\lambda^{\pm n/2}\right) \quad (\lambda \rightarrow 0)$$

(The same is true for $u^+(r)$.) Let us remark that the same type of singularity in λ^2 is obtained from (4.2) - (4.3), so the nature of the singularity in the λ^2 plane is apparently independent of the energy.

Bearing these considerations in mind, we can unify the treatment of the repulsive and attractive cases by introducing an auxiliary variable σ into the differential equation (4.1) :

$$\left(\frac{d^2}{d\sigma^2} + k^2 - \frac{n^2 - 1/4}{\sigma^2} - \frac{\sigma \lambda^2}{\sigma^6} \right) u(\sigma, r) = 0 \quad (4.1')$$

The solution $u(\sigma, r)$ are analytic in the complex σ plane, cut along the negative real axis. The repulsive interaction corresponds to the solution $u(1, r)$, while the attractive one to $u(-1 \pm i0, r)$; in the attractive case real solutions can be obtained by taking $u(-1 + i0, r) + u(-1 - i0, r)$ and $i^{-1} \sqrt{u(-1 + i0, r) - u(-1 - i0, r)}$, respectively.

One of the most interesting features of the expression obtained for the ladder diagram considered in Ref. ¹⁾ was that the scattering amplitude was free of arbitrary subtraction constants. The reason of this interesting and, in a sense, astonishing result has been that due to the strong singularity of the interaction at $r = 0$, the boundary condition imposed on the wave function was independent of n and so we had $F_n(p) = \phi_n = 0$ in the subtraction equation (Eq. (2.12) in the present paper). Is this still true in the present, more complicated case ?

For the "repulsive" channel (u^+) the equation which the wave function obeys is the same as that in Ref. ¹⁾, so all the results can be carried over without any change. Thus the second term on the right-hand side of Eq. (2.12) vanishes identically and the expression of T^+ is free of arbitrary subtraction constants. The situation is more complicated in the "attractive" channel (u^-), because of the arbitrariness in the boundary condition. In order to decide which solution is to be chosen, we consider the zero energy limit of the scattering amplitude. Although the effective range expansion is divergent (the interaction does not fall off exponentially at large distances) its first few terms can be used as an asymptotic estimate for the amplitude as $k^2 \rightarrow 0$; in particular the scattering length is necessarily finite. At small energies only the lowest partial wave contributes ($n = 1$), the scattering length can be found by expanding the zero energy solution (4.2) and (4.3) at $n = 1$. Thus we find the scattering length of the "+" channel

$$a^+ = \lambda/2 \quad (5.1)$$

Let us observe now that for the lowest partial wave (and only for that !) the "four dimensional" and "ordinary" partial amplitudes coincide. In fact, the coefficient of the lowest wave is $Z_{10}^0(\cos\theta) = P_0(\cos\theta) = 1$, where $\cos\theta$ is the

scattering angle in the c.m.s. ¹⁾. Thus, unitarity requires that the scattering length of the $n = 1$ eigenamplitude be finite and if complex, its imaginary part positive. This is still not sufficient to determine the boundary condition uniquely. We now require the uniqueness of the scattering amplitude (the absence of subtraction terms). This already fixes the solution to be chosen uniquely : in terms of the solutions of (4.1') it is $u(-1+i0, r)$, giving a purely imaginary scattering length in the "-" channel

$$a^- = i\lambda/2 \quad (5.2)$$

One might wonder, why did we obtain complex scattering length. A glance at Fig. 2 makes the reason clear ; as we consider massless particles, all the many particle cuts of the amplitude begin at $k^2 = 0$ together with the "elastic" one, so

Fig. 2

that there is no region, where elastic unitarity would be satisfied. (By the way, that is the explanation of the "paradox", mentioned in section 2.) Nevertheless, it is interesting that the scattering length is proportional to the first power of the Fermi coupling constant ; thus by comparing (5.1) with the first Born approximation to the scattering amplitude, (which obviously can give a real scattering length only) one can see that our ladder reproduces it apart from a finite renormalization of the coupling constant ¹²⁾.

6. ZERO ENERGY LIMIT OF THE W THEORY

In order to compare our method with that of Feinberg and Pais ³⁾ we sketch briefly the treatment of an uncrossed ladder with W meson exchange. We consider the zero energy limit only.

Let us make the same assumption as before, namely that $G_{\mu\nu} = g_{\mu\nu} G$. The kernel $K_{\mu\nu}$ is proportional of course to the vector meson propagator

$$K_{\mu\nu} = g^2 \frac{g_{\mu\nu} - m^{-2} q_\mu q_\nu}{m^2 - q^2}$$

We now proceed as in Ref. ³⁾, i.e., neglect the term proportional to $g_{\mu\nu}$. Then

$$K = K_\mu{}^\mu = - \frac{g^2}{m^2 - q^2} + \frac{g^2}{m^2} \quad (6.1)$$

In co-ordinate representation we find the BS equation (2.8) with the first term only on the right-hand side of (6.1)

$$\left(\frac{d^2}{d\tau^2} + k^2 - \frac{n^2 - 1/4}{\tau^2} \pm m g^2 \frac{K_1(m\tau)}{\tau} \right) u^\pm(\tau) = 0 \quad (6.2)$$

The interaction behaves as τ^{-2} at the origin, so the leading singularity can be included into the centrifugal term. We replace the potential by the following one :

$$V(\tau) = \begin{cases} g^2 \tau^{-2} & (\tau < m^{-1}) \\ 0 & (\tau > m^{-1}) \end{cases} \quad (6.3)$$

7. DISCUSSION

There are two possible attitudes towards the results presented in the previous sections. The first one is to say that in consequence of the very crude approximations we made, they are of no relevance at all. In this respect we refer the reader to the critical discussion of the weak points of our approach given in Ref. ¹⁾.

Hoping now that those sharing this negative opinion have stopped reading the present paper somewhere at the beginning of the third section, we take the other standpoint : supposing that in spite of the approximations made the results still reflect some properties of the - not yet existing - theory of non-renormalizable interactions (or even of nature), what is the lesson we can learn from the present calculation ?

The first and main thing to be underlined is that the annoying phenomenon of having a "theory" with an infinite number of arbitrary constants (i.e., in fact, no theory at all) has disappeared. One can see that the reason of having infinitely many subtraction constants in perturbation theory is that we try to expand a function around a singular point. In fact we have got more than we expected : the final answer we get is completely free of arbitrary constants. So if this were true, weak interactions would be in a sense better than, say, ordinary electrodynamics. To go further, one can speculate whether when treating ordinary strong interactions, should one not include always weak interactions in order to get a natural cut-off in the theory ? (In the present model this is definitely true : if one added to the fermion loop in the kernel any other term, representing strong interactions, the renormalization terms arising from the latter would be annihilated by the strong singularity due to the Fermi interaction.)

The second question one is naturally led to ask : how general this statement is ? We feel that it goes beyond the approximation we employed in the present paper (local kernels). In fact they depend on the strong singularity of the Green function at small distances. Power counting definitely indicates that any Green function involving non-renormalizable interactions shares this property. By now it is, however, unclear how reliable a power counting argument is and whether the Green functions can be made to vanish at small distances. (Power counting indicates the presence of an essential singularity but not the behaviour around the singular point - or the light cone in Minkowskian metric.)

Coming back to reality, we would like to discuss briefly the prediction of the theory for weak interactions. The "Fermi ladder" we considered has the unpleasant property that it predicts equal rate for "allowed" and "forbidden" reactions at low energies. In fact, if we add and subtract Eqs. (51.) and (5.2), we find the scattering lengths of reactions (3.1) and (3.2)

$$\begin{aligned}\bar{a} \text{ allowed} &= \frac{\lambda}{2} (1+i) \\ \bar{a} \text{ forbidden} &= \frac{\lambda}{2} (1-i)\end{aligned}$$

unlike the vector meson ladder, which gives

$$\begin{aligned}\bar{a} \text{ allowed} &= 3/4 g^2/m^2 + O(g^4) \\ \bar{a} \text{ forbidden} &= 0 + O(g^4)\end{aligned}$$

(The physical reason for having obtained complex scattering lengths for the Fermi ladder has been already explained.)

If one believed these results, one would consider this as an argument for a W meson theory against four-fermion interaction. Nevertheless, one should bear in mind that neglecting fermion masses is not likely to be a good approximation at low energies (it is certainly justified if one is interested in the leading singularity). Including the masses of the particles into the calculation could change the situation appreciably, - not only the threshold properties of the Green function.

The author takes pleasure in thanking Professors V.F. Weisskopf and L. Van Hove for the hospitality extended to him at CERN, and Drs. G. Furlan and W. Kummer for some interesting discussions. A grant received from the Joint Institute for Nuclear Research, Dubna, making his stay at CERN possible, is gratefully acknowledged.

REFERENCES AND FOOTNOTES

- 1) G. Domokos, P. Suranyi and A. Vancura, "Fermi Interactions at High Energies", Dubna preprint (1963).
- 2) We recall that we called a BS kernel local, when it depends only on the difference of the relative momenta in the initial and final states.
- 3) G. Feinberg and A. Pais, Phys. Rev. 131, 2724 (1963) ; ibid. 133, B 477 (1964).
- 4) R.F. Sawyer, preprint (1963).
- 5) See, e.g., A. Bastai, L. Bertocchi, S. Fubini, G. Furlan and M. Tonin, Nuovo Cimento 30, 1812 (1963) ;
G. Domokos and P. Suranyi, Dubna preprint (1963) - (I).
These papers contain further references.
- 6) K. Symanzik, Hercegnovi Lecture (1962).
- 7) The special problems connected with the transition to Euclidean momenta in the case of non-renormalizable theories have been discussed in Ref. ¹⁾.
At present we have nothing to add to that discussion.
- 8) In fact, it is not difficult to show that in the problem we are going to investigate here, the neglected off-diagonal elements are bounded ; on the other hand they are proportional to $(E_\mu E_\mu)^{1/2}, E_\mu E_\mu, \dots$, so for sufficiently small energies they are small.
- 9) Mirza A. Baqi Beg, Brookhaven preprint (1963) ;
C. Bouchiat, B. d'Espagnat and J. Prentki, Nuovo Cimento 31, 7587 (1964).
- 10) Cf., e.g., P.M. Morse and H. Feshbach, Methods of Theoretical Physics, (McGraw-Hill, New York 1953), Chapter 9.3.
- 11) N.N. Bogoliubov, D.V. Shirkov, Introduction to the Theory of Quantized Fields, (Interscience, New-York, 1959), Chapter IV.
- 12) This is not trivial, as the subtraction terms vanish.