Phys. Rev. Letters **19**, 1056 (1967); W. D. Walker, Rev. Mod. Phys. **39**, 695 (1967); A. C. Clegg, in Proceedings of the Heidelberg Conference on Elementary Particles, Heidelberg, Germany, 1967 (to be published), where reference to previous analysis can be found.

¹⁴C. Lovelace, R. H. Heinz, and A. Donnachie, Phys. Letters **22**, 332 (1966).

¹⁵L. Brown and P. Singer, Phys. Rev. **133**, B812 (1964).

¹⁶J. P. Baton, G. Laurens, and J. Reigner, Phys. Letters **25**, B419 (1967).

¹⁷See, for example, I. J. R. Aitchison, Nuovo Cimento **51A**, 250, 272 (1967).

¹⁸M. Month, Phys. Letters **18**, 357 (1965); note that the total energy of the 3π system (1836 MeV) is close to the energy (1780 MeV), where the triangular diagram due to the resonance f^0 should have a singularity, with the effect of an enhancement of the population of the Dalitz plot at the intersection of the two f bands.

PHASE CONTOURS OF COLLISION AMPLITUDES*

Charles B. Chiu
Cern, Geneva, Switzerland

and

Richard J. Eden† and Chung-I Tan

Lawrence Radiation Laboratory, University of California, Berkeley, California

(Received 8 December 1967)

Phase contours are curves along which the phase of a collision amplitude is a real constant. In their simplest form they describe the phase as a function of real energy and momentum transfer and provide a convenient summary of experimental results for a scattering process. More generally they can be used to provide consistency tests in the dynamics of strong interactions of elementary particles.

We will give a brief discussion of the following topics: (1) properties of phase contours, (2) phase contours for pion-nucleon scattering below 1.4 GeV, (3) phase contours in a Regge model for pion-nucleon scattering, and (4) the use of phase contours for studying consistency between resonance poles, zeros, and high-energy behavior in a crossing symmetric model.

(1) Properties of phase contours.—The phase $\varphi(s, t)$ of an invariant scattering amplitude $F(s, t)$ is defined by

$$\varphi(s, t) = \text{Im}[\ln F(s, t)], \quad (1)$$

where Im denotes the imaginary part, and s, t denote the invariant energy and momentum transfer variables. It is also necessary to define the phase at an initial point (s_0, t_0) . When the amplitude has zeros or poles on the physical sheet, the phase depends on the route taken from (s_0, t_0) to (s, t) , so we must always specify the route.

A phase contour is defined by

$$\varphi(s, t) = C, \quad (2)$$

where C is a real constant. It is useful to study

phase contours both for real s and t , and for complex s and fixed t , etc. Their properties include the following:

(a) Phase contours, for different values of C , do not meet each other except at zeros and poles of the amplitude and at certain other singularities.

(b) The phase change clockwise in the complex plane round a zero is -2π , and round a pole is 2π .

(c) For fixed t , and $s = s_1 + is_2$, the phase is a harmonic function of s_1 and s_2 , and the phase contours are orthogonal to the modulus contours in this complex s plane.

(d) From the optical theorem,

$$\text{Im}F(s, 0) > 0, \quad \text{for } s > (m+M)^2. \quad (3)$$

Assuming an asymptotically constant total cross section and the Pomeranchuk theorem,

$$F(s, t) \sim isB, \quad \text{as } s \rightarrow +\infty. \quad (4)$$

We choose our initial point (s_0, t_0) as the limit along $(s+i0, t=0)$ as $s \rightarrow +\infty$, and take

$$\varphi(s \rightarrow +\infty, 0) = \frac{1}{2}\pi. \quad (5)$$

(e) The phase contours $\varphi(s, t) = 0$, or π , cannot enter the region $0 \leq t < 4m^2$, when $s > 4m^2$.

(f) If the scattering amplitude has power behavior as $s \rightarrow \infty$, the phase is asymptotically constant. In the particular case of a symmetric amplitude, or a Regge term of even signature, we can write

$$F(s, t) \sim \beta(t)s^{\alpha(t)} \exp[i\pi\{1 - \frac{1}{2}\alpha(t)\}]. \quad (6)$$

If $\beta(t)$ is nonzero in the range $[0, t]$, and $\alpha(t)$ is real, then the asymptotic phase is given by

$$\varphi(s, t) \sim \pi[1 - \frac{1}{2}\alpha(t)], \quad (7)$$

so the asymptotic phase contours are parallel to $t = \text{const.}$

(g) Assuming asymptotic power behavior s^α of $|F|$, the power $\alpha(t)$ can be determined from the phase contours along the real s axis and the zeros of F in the complex s plane (and poles if there are bound states). The location of zeros is of great importance. They are indicated by intersections of phase contours, and their complex locations can be studied by means of phase-contour surfaces when both s and t are complex variables in Eqs. (1) and (2).

(2) Phase contours in pion-nucleon scattering.—We have used the results of phase-shift analysis of pion-nucleon scattering experiments to obtain phase contours for invariant amplitudes,¹ in the energy range

$$0 < T_\pi < 1.4 \text{ GeV}, \quad (8)$$

where T_π denotes the pion kinetic energy. The amplitudes used are the invariant combinations of π^-p and π^+p scattering amplitudes²

$$A^{(+)}, B^{(+)}, A^{(-)}, B^{(-)}. \quad (9)$$

The resulting phase-contour diagrams provide a useful visual aid for comparing different phase-shift solutions and for indicating regions where the resulting scattering amplitude is complicated. In these regions one can conclude that more experiments are desirable, especially in the region around $T_\pi = 0.8 \text{ GeV}$, where there are several resonances and the phase contours for the last three amplitudes listed in (9) indicate unusually rapid phase variations.¹

Phase contours for $A^{(+)}$, derived from the Lovelace phase shifts,³ are shown in Fig. 1. There are two real zeros of this amplitude, indicated by intersections of phase contours, and it is evident that there are some nearby complex zeros. Complex zeros are indicated by a bunching of phase contours, for example, in Fig. 1 between 0.2 and 0.6 GeV. The modulus contours¹ for $A^{(+)}$ show a valley in this region, where the amplitude is very small.

(3) Phase contours in a Regge model for pion-nucleon scattering.—The method of phase contours is useful in studying the relation between high-energy and low-energy scattering. It can be applied either as a consistency meth-

FIG. 1. Phase contours for the $A^{(+)}$ amplitude in pion-nucleon scattering derived from the (1966) Lovelace phase shift solutions.

od, or as an aid in comparing phenomenological solutions for a given scattering process. We illustrate the latter by giving in Fig. 2 the phase contours for $A^{(+)}$ that are obtained by extrapolating from a Regge model for pion-nucleon scattering. Our model is not meant to be realistic at this stage, but it gives surprisingly good agreement with the phase contours shown in Fig. 1 out to 90 deg for $T_\pi > 1 \text{ GeV}$.

Our Regge model is the sum of a Regge solution⁴ for near-forward πN scattering based on P , P' , and ρ exchange, and a Regge solution⁵ for near-backward πN scattering based on N and N^* exchange. We have also assumed that the trajectories fall linearly as t decreases (or as u decreases), and have included effects from zeros of residues.

The modulus contours for $A^{(+)}$, that correspond to Figs. 1 and 2, also show a striking resemblance in the low-energy region.¹ However, the Regge solutions for the other three amplitudes do not resemble the solutions obtained from phase-shift analysis. A comparison of their phase-contour diagrams¹ indicates that a more complicated Regge model is required in this low-energy region.

FIG. 2. Phase contours for $A'^{(\pm)}$ derived from a Regge model for pion-nucleon scattering.

(4) Resonance poles, zeros, and high-energy behavior.—Phase contours provide a method for studying part of the consistency problem in strong interactions, and may help towards understanding whether the bootstrap problem can be formulated in a meaningful approximation. We do not introduce unitarity at this stage, but consider only the consistency that is required by crossing symmetry for a given high-energy behavior, taking into account the associated resonance poles (Regge poles). We consider⁶ a symmetric scattering amplitude for equal-mass spinless bosons, and construct a class of models that is consistent with a high-energy behavior based on Regge terms of the type indicated in Eq. (6). Our model includes effects of zeros of residues below threshold, and resonance poles above threshold on unphysical sheets. We also assume dominance by a leading Regge pole with a continuously rising trajectory.

We find zeros of the scattering amplitude from three sources⁶: (1) From symmetry requirements, there are zeros along $s = u$ in $t < 0$, for s real ($s + i0$) and u real ($u - i0$) above and below their branch cuts, respectively; (2) from zeros of residues, there are zeros of the amplitude that move in from infinity along $\text{Re}(s) = \text{Re}(u)$ as t varies in $t < 0$; and (3) from interference between resonance poles. By studying the phase-contour surfaces for complex

s and t , we find that these three types of zero may be identified as lying on different parts of two-dimensional complex surfaces of zeros in the four-dimensional space.⁶

Further analysis of phase contours is in progress and further details of the above work will be given elsewhere.^{1,6}

We are indebted for hospitality and for helpful discussions to Professor G. F. Chew at the Lawrence Radiation Laboratory and Dr. L. Van Hove at CERN.

*This work was supported by the U. S. Atomic Energy Commission.

†Address after 1 January 1968: Cavendish Laboratory, Cambridge, England.

¹C. B. Chiu, R. J. Eden, and C.-I Tan, University of California Radiation Laboratory Report No. UCRL-17899, 1967 (unpublished).

²These amplitudes are defined, for example, by R. J. Eden, *High Energy Collisions of Elementary Particles* (Cambridge University Press, New York, 1967).

³C. Lovelace, in *Proceedings of the Thirteenth International Conference on High Energy Physics, Berkeley, 1966* (University of California Press, Berkeley, California, 1967).

⁴W. Rarita, R. J. Riddell, Jr., C. B. Chiu, and R. J. N. Phillips, *Phys. Rev.* **165**, 1615 (1967).

⁵C. B. Chiu and J. Stack, *Phys. Rev.* **153**, 1575 (1967).

⁶R. J. Eden and C.-I Tan, University of California Radiation Laboratory Report No. UCRL-17922, 1967 (unpublished).